



**LIBRARIES**  
UNIVERSITY OF WISCONSIN - MADISON

## Meprospan advertisement.


[s.l.]: [s.n.], 1960

<https://digital.library.wisc.edu/1711.dl/LSCZWMHGNTMDR8S>

<http://rightsstatements.org/vocab/InC/1.0/>

The libraries provide public access to a wide range of material, including online exhibits, digitized collections, archival finding aids, our catalog, online articles, and a growing range of materials in many media.

When possible, we provide rights information in catalog records, finding aids, and other metadata that accompanies collections or items. However, it is always the user's obligation to evaluate copyright and rights issues in light of their own use.



## EMOTIONAL RELIEF

FOR THE

## PHYSICALLY ILL



### RELIEVES ANXIETY, APPREHENSION AND TENSION...



*All day long*

... keeps the patient calm,  
and the mind clear.



*All night too*

... aids restful sleep, with  
no barbiturate hangover.

## MEPROSPAN®-400

(MEPROBAMATE 400 MG. SUSTAINED RELEASE)

*Simplified, convenient dosage for emotional relief.*

**Side effects:** 'Meprospan' (meprobamate, sustained release) is remarkably free of untoward reactions. Daytime drowsiness has not been reported. Rare allergic or idiosyncratic reactions may occur, generally developing after 1-4 doses of the drug.

**Contraindications:** Previous allergic or idiosyncratic reactions to meprobamate contraindicate subsequent use.

**Precautions:** Should administration of meprobamate cause drowsiness or visual disturbances, the dose should be reduced. Operation of motor vehicles or machinery or other activity requiring alertness should be avoided if these symptoms are present. Effects of excessive alcohol may possibly be increased by meprobamate. Prescribe cautiously and in small quantities to

patients with suicidal tendencies. Massive overdosage may produce lethargy, stupor, ataxia, coma, shock, vasomotor and respiratory collapse. Consider possibility of dependence, particularly in patients with history of drug or alcohol addiction; withdraw gradually after prolonged use at high dosage.

*Complete product information available in the product package, and to physicians upon request.*

**Usual adult dosage:** One 400 mg. capsule or two 200 mg. capsules at breakfast; repeat with evening meal.

**Supplied:** 'Meprospan'-400 (meprobamate 400 mg.), 'Meprospan'-200 (meprobamate 200 mg.), each in sustained-release capsules. Both potencies in bottles of 30.