



LIBRARIES
UNIVERSITY OF WISCONSIN - MADISON

Fiorinal advertisement.

[s.l.]: [s.n.], 1978

<https://digital.library.wisc.edu/1711.dl/QV4RJLHCPJVPV8A>

<http://rightsstatements.org/vocab/InC/1.0/>

The libraries provide public access to a wide range of material, including online exhibits, digitized collections, archival finding aids, our catalog, online articles, and a growing range of materials in many media.

When possible, we provide rights information in catalog records, finding aids, and other metadata that accompanies collections or items. However, it is always the user's obligation to evaluate copyright and rights issues in light of their own use.



FIORINAL® WITH CODEINE Ⓢ

CAPSULES, in three dosage strengths: Each capsule contains: Codeine phosphate, USP. (*Warning: May be habit forming*) 7.5 mg (1/8 gr) in **No. 1**, 15 mg (1/4 gr) in **No. 2**, 30 mg (1/2 gr) in **No. 3**; Sandoptal® (butalbital, USP) (*Warning: May be habit forming*) 50 mg; caffeine, USP, 40 mg; aspirin, USP, 200 mg; phenacetin, USP, 130 mg.

HAS THE SEDATIVE EXTRA.

The three leading narcotic/analgesics don't.

Analgesic Preparation	Sedative Component
FIORINAL® with CODEINE capsules Ⓢ	SANDOPTAL® (butalbital, USP)
TYLENOL® with Codeine tablets Ⓢ Tablets containing codeine phosphate (<i>Warning: May be habit forming</i>) 7.5 mg (1/8 gr) in No. 1 , 15 mg (1/4 gr) in No. 2 , 30 mg (1/2 gr) in No. 3 , or 60 mg (1 gr) in No. 4 , and acetaminophen 300 mg.	NONE
EMPIRIN® with Codeine Ⓢ Tablets containing aspirin (acetylsalicylic acid) 325 mg, plus codeine phosphate (<i>Warning: May be habit forming</i>) 15 mg in No. 2 , 30 mg in No. 3 , or 60 mg in No. 4 .	NONE
PERCODAN® Ⓢ Tablets containing oxycodone hydrochloride (<i>Warning: May be habit forming</i>) 4.50 mg, oxycodone terephthalate (<i>Warning: May be habit forming</i>) 0.38 mg, aspirin 325 mg.	NONE

FIORINAL® with CODEINE capsules

Contraindications: Hypersensitivity to any of the components.

Precautions: May be habit forming due to presence of codeine and barbiturate.

Side Effects: Nausea, vomiting, constipation, dizziness, skin rash, drowsiness, and miosis. Overdosage is primarily manifested by drowsiness.

Adult Dosage: 1 or 2 capsules, repeated if necessary up to 6 capsules per day, or as directed by physician. *Before prescribing or administering, see package circular for full product information.*

SANDOZ PHARMACEUTICALS, EAST HANOVER, NJ 07936

© 1979 Sandoz, Inc.

