



LIBRARIES
UNIVERSITY OF WISCONSIN - MADISON

Preludin advertisement.

[s.l.]: [s.n.], 1960

<https://digital.library.wisc.edu/1711.dl/3P5YEEJX2KW458H>

<http://rightsstatements.org/vocab/InC/1.0/>

The libraries provide public access to a wide range of material, including online exhibits, digitized collections, archival finding aids, our catalog, online articles, and a growing range of materials in many media.

When possible, we provide rights information in catalog records, finding aids, and other metadata that accompanies collections or items. However, it is always the user's obligation to evaluate copyright and rights issues in light of their own use.

reducing the problems of reducing



Preludin[®] Tablets and
Endurets[®]
brand of phenmetrazine HCl prolonged-action tablets

**an oxazine...
not an amphetamine**

Unsurpassed Effectiveness

In all controlled clinical studies, Preludin has produced impressively greater weight loss than placebo tablets regardless of the degree of enforcement of dietary restriction.

Exceptionally High Tolerance

Reports are numerous of successful use of Preludin in cases intolerant of other anorexians.

Flexibility of Dosage

Available as scored tablets of 25 mg. for b.i.d. or t.i.d. administration and also as Endurets[®], 75 mg., for once daily administration.

Precautions and Contraindications

Although there have been no reports of significant toxic reactions to Preludin, on theoretical grounds it should not be given to patients with severe hypertension, thyrotoxicosis or acute coronary disease.

Preludin may be used with caution in cases of moderate hypertension and cardiac decompensation.

Preludin[®], brand of phenmetrazine hydrochloride.
Under license from C. H. Boehringer Sohn, Ingelheim.



Geigy Pharmaceuticals
Division of Geigy Chemical Corporation
Ardsley, New York

Geigy