



LIBRARIES

UNIVERSITY OF WISCONSIN-MADISON

Menominee Iron Range: [specimens 23260-23362]. No. 176 1895

Weidman, Samuel, 1870-1945

[s.l.]: [s.n.], 1895

<https://digital.library.wisc.edu/1711.dl/AAHZHJP4DHG269D>

<http://rightsstatements.org/vocab/InC/1.0/>

For information on re-use see:

<http://digital.library.wisc.edu/1711.dl/Copyright>

The libraries provide public access to a wide range of material, including online exhibits, digitized collections, archival finding aids, our catalog, online articles, and a growing range of materials in many media.

When possible, we provide rights information in catalog records, finding aids, and other metadata that accompanies collections or items. However, it is always the user's obligation to evaluate copyright and rights issues in light of their own use.

U. S. GEOLOGICAL SURVEY
FIELD SECTION BOOK

9-891

LAKE SUPERIOR DIVISION.

INSTRUCTIONS.

1. Ordinarily at least two pages of this book will be devoted to one section. On the left-hand page, place a map of as much of the section as has *actually been seen*. Denote rivers, lakes, marshes, etc., by the usual topographical signs. Denote the ledges of rock, when no structure is made out, by cross-hatching, making the cross-hatching cover as nearly as possible the areas occupied by the exposures. If the rock is a massive one, but still more or less plainly bedded, use the same sign with a dip arrow and number attached, showing the direction and amount of the dip. Denote a shaly or other very plainly bedded ledge by right parallel lines, and a ledge having a secondary structure by wavy parallel lines running in the direction of the strike, with dip arrow and number attached as before. The greatest care must be taken to avoid confusing slaty or schistose structure with bedding, and in all cases where there is the least doubt about the true bedding direction, indicate it by a query. To each exposure on the face of the map attach the number of the specimen representing it. In mapping the section count each of the spaces between the blue lines as 100 paces, and twenty of these spaces to one mile, or 2,000 paces. Usually the southeast corner will be placed at the bottom of the page, or at the first black line above the bottom of the page, and at the right-hand side. If, however, for any reason, it is desirable to show portions of an adjoining section, the southeast corner may be shifted up, or the map may be turned around and the north placed at the left-hand side of the page. The ruling of the left-hand pages is also arranged so that, if desirable, a larger or a smaller scale can be used, eight inches, two inches, one inch, or one-half inch to the mile. With the two-inch scale, the squares outlined in black represent sections, and those in red, quarter sections and "forties," while the space between the blue lines is 200 paces.

2. On the right-hand page place the notes descriptive of the exposures. Begin in each case with the number of the specimen, placing the number on the left-hand side of the red line, after which give in order on the right of the same red line the position of the ledges as reckoned in paces from the southeast corner of the section and the dip and strike when observable, the latter always being expressed from the north; for instance 4025, 250 N., 300 W., *Strike, N. 78° E., Dip 50° S.* Then follow with a full description of the ledge. When topographical maps are used for locations this paragraph applies only in part.

3. Collect a specimen from every ledge, or wherever there is a change of rock on any one ledge, taking care to get fresh material, unless for a special purpose the weathered surface is desired. In case of trips made on foot or in canoes, for long distances, neighboring ledges, unquestionably of one kind of rock, need not be specimened. The position and extent of the ledges not specimened should be marked on the map, with notes that each is of a rock identical with specimen so-and-so. Under the same conditions small-sized specimens, trimmed to a uniform size of 2 x 2½ x ½ inches will be allowed, but in all other cases *large-sized specimens*, trimmed to a size of 3 x 4 x 1 inches, must be selected, in accordance with section 3, chapter IV, p. 44, Regulations of the U. S. Geological Survey. Specimens should not be placed together without protection in the collecting bag, as the fresh surfaces, important in determining the character of rocks, are thus destroyed. They should be damaged by no temporary mark, but the numbers should be at once marked in at least two places upon the inclosing paper or cloth bags. Specimens may be permanently marked in camp by painting the numbers upon them in white upon a black background, using Silver White and Ivory Black oil tubes for color, with turpentine as a diluent.

4. On the last twenty-five pages of the book give, as may seem desirable, a general account of the examination of the region mapped in the previous pages, correlation of observations, sketches, cross sections, etc.

5. Forward this note book as soon as filled as registered mail matter to C. R. Van Hise, U. S. Geologist, Madison, Wis.


176


176 ✓

S. 28.

T. 42

R. 53






S. 34

T. 42

R. 33



23260 575 N 1025 W 34-42-33

Ledge of slate somewhat ferruginous and not cherty. Strike N 80 W. The slate outcrops along the brow of a long ridge.


23261 800 N 1440 W 34-42-33

Specimen of banded cherty slate taken from the mouth of test pit. The greater part of the material about the pit seems to be like specimen. A good deal of it is a much fractured slate. In this pit the Dip is clearly made out, and is 80° N.E. Strike N 80° W

The strike and dip given is thought to be the true strike and dip of bedding; that is, that the bedding and cleavage is the same

S. 27+22 T. 42

R. 33




S. $\frac{5}{2}$ of 10
and
Sec 13-

T. 42

R. 33

10³

$\frac{11}{11}$ $\frac{11}{11}$



S. $\frac{23}{26}$
25

T. 42

R. 33

22

23³

24

23

24

2 $\frac{1}{2}$

3

3

27

26

28

2 $\frac{1}{2}$

2 $\frac{1}{2}$

2 $\frac{1}{2}$

32

32 $\frac{1}{2}$

36


C.N.W. R.R.

3 $\frac{1}{2}$

S. 11/12/7

T. 42

R. 33/32



0-747

29262 1000 N 1125 W 12-42-32

This place looks as though there had been some test pitting done, fragments of quartz slate are scattered about a little knoll.

at 1000 N 912 and 950 W are test pits in jaspery slate
And also at 875 and 900 W
1000 N are similar test pits in slate formation

29263 1000 N 1100 W 7-42-32

Test pit on top of ridge in black cherty slate,

23264 1165 N 1650 W 8-42-32


Large ledge of black slate, not
much cherty material in it.
Strike of Schistosity N10W
Dip is about vertical

23265 About 20 paces E of where the
S.E. corner of Sec 8-42-32 is
thought to be, Specimen is
from test pit about 25 feet
deep, in sand plain. Rock
is a cherty slate.

S. 17

T. 42

R. 32



23266 70 N 150 W 17-42-32

Ledge of fine grained and amygdaloidal greenstone, much squeezed and stretched in places.

23267 1910 N 300 W -20-42-32

Schistose greenstone, some large outcrop.

At 25°N 675°W 17-42-32 is an outcrop of greenstone conglomerate having a strike slightly W of N.

Strike of the schistosity in other places in this vicinity is slightly W of N.

23268, 200 N 1500 W 17-42-32.

Ledge of greenstone conglomerate, quite fine and quite coarse also. Strike of schistosity slightly W of N

23276 250 N 1740 W 17-42-32

Ledge of greenstone containing what seems to be fragments of black rock like specimen

23277 550 N 575°W 17-42-32

Ledge of fine grained greenstone in places schistose and in places conglomeratic and also amygdaloidal.

23278 720 N 70 W 17-42-32. Ledge of black slate on R.R. Strike of schistosity N20°W Dip 90°

X 23269 1750 N 47° W 19-42-32


Ledge of soft black slate.
Strike of schistosity slightly W of N
Dip about vertical

23270+71 1700 N 54° W 19-42-32

Ledge of cherty slate conglomerate or breccia. Outcrop contains angular fragments of (greenstone?) or slate up to 1 foot in diameter as well as very small fragments.

23272 1760 N 54° W 19-42-32

Going N from above coarse formation like 23270-71 and towards the slate like 23269 is found a finer conglomerate or breccia 23272.



Strike of ^{schistosity} formation slightly W of N Dip 90°


This place was visited by Profs. Van Dine and Bayley who considered it a brecciated slate.

B3

S. 13

T. 42

R. 33



988

23273 1930 N 450 W 24-42-33

Ledge of greenstone conglomerate.

1950 N 5-35-W 24-42-33

Ledge of fine grained amygdaloidal greenstone mingled with which is greenstone conglomerate like 23273

23274 1750 N 1610 W 24-42-33

Specimen taken from mouth of old pit. It looks much like a quartzitic slate but it may be a greenstone.

23275 680 N 760 W 13-42-33.

Ledge of soft black slate.

Strike of schistosity N 20 W Dip 75° N.

But the true dip and strike of the bedding or banding as shown in specimen is -

Strike N 45° E Dip 45° N. W.


The bedding or banding is very much folded and curved so that can be certain as to general Dip and strike, but am certain that bedding and schistosity do not agree.

S. 19

T. 42

R. 32

19



500 N 1480 W. 19-42-32


Edge of black shaly slate

Strike of schistosity $N 10^{\circ} W$ $Dip 90^{\circ}$

S. 24

T. 42

R. 33



23279 500 N 1630 W 24-42-33

Test pit in cherty slate.

Four pits along here in a row.

Delphic Mine

23280 550 N 900 W 24-42-33 (Delphic Mine)

Specimen of black soft shaly slate which is a fair specimen of the greater part of the material about the excavation.

23281 (Delphic Mine) Specimen containing iron sulphide (iron pyrite). The odor of sulphur is easily detected about the dump heap.

23282 (Delphic Mine) Specimens containing considerable iron.

23283 (Delphic Mine) Specimen showing jasper and iron pyrite.


23284 (Delphic Mine) Specimen of schistose slate containing much iron pyrite. This mine apparently has never yielded any ore. There is a large amount of worthless material about the excavation.

Strike and Dip of the slate could not be seen. But the rock is very much shattered & fractured.

S. 30

T. 42

R. 32



ON 200' W 30-42-32

On summit of ridge, fine slate fragments about roots of an old upturned tree; no doubt but what there is slate in situ beneath.

23285 1950 N 1775 W 31-42-32

Ridge of cherty slate.


23287 420 N 1080 W 30-42-32

Outcrop of carbonaceous shaly slate.

S. 25-

T. 42

R. 33




28286 1825-N 150 W 36 - 42 - 33

Specimen of jaspery slate on
large hill. Strike $N 10^{\circ} W$
Dip is vertical. This seems
to be bedding as indicated
by the bands of jasper.

S. 31
6

T. 42
41

R. 32



2328~~28~~ 500 N 1400 W 31-42-32

Specimen from the large ridge of slate, a cherty slate.

Strike $N 45^{\circ} W$ Dip Vertical.

23289 530 N 1660 W 31-42-32

Specimen of jaspery slate showing crumpling. Strike $N 50^{\circ} W$ Dip 90°

At 2650 N 1730 W 31-42-32 the strike is still more to the W. $N. 68^{\circ} W$.

23293 1970 N 1826 W 6-41-32


Specimen is a very coarse quartzite. At this place is a ledge of massive quartzite about 45-feet thick, (across the strike). Strike is

$N 50^{\circ} W$. Dip 90° . This is the true bedding as indicated by thin seams of coarse quartzite like specimen. The rest of the rock is like specimen 23291+2. Thin seams of quartzite schist due to the pressure run along the strike. Fracturing and faulting has taken place

S. 36

T. 42

R. 33



93290 510 N 300 W 36-42-33

Outcropping at edge of bluff
Banded slate. Strike N 65° W
Dip 75° NE

93291 125 N 275 W 36-42-33

Edge of black ferruginous
massive quartzite. Strike and
dip of bedding agrees with
that of the slates to the
north.


93292 125 N 275 W 36-42-33

Schistose quartzite. The bed
of quartzite at this place
shows a thickness of at
least 30 feet.

S. 6

T. 41

R. 32



23294 25°N 1710 W See 31-42-32

Ledge of quartzite schist (ferruginous and black) The quartzite in massive beds is seen in numerous ledges to the S.W. of this and no doubt grades into this. Strike $N65^{\circ}W$. 100 paces north of this is another large exposure showing beds of massive and schistose quartzite.

23295 175°N 1950 W 31-42-32

Ledge of gneiss slates interbedded with thin seams of quartzite as shown by the three specimens numbered 23295;

23296 1300 N 1000 W 6-41-32

Ledge of slate and quartzite. This is a small ledge and immediately to the north and south of it is quartzite again.

Further S.E. along the Brule as far east at least as 130 in sec 5-41-32 several ledges of massive ferruginous quartzite was seen.

Clements says he finds no quartzite along the N and S. $\frac{1}{4}$ line near ~~in~~ the river in Sec 5.

The ledge along the river near the west side of Sec 5. Strike N-63°-70°W

In this whole area of quartzite the strike varies from N 45°W ~~toward~~ the slate area to N 65°-70°W further S.E. near B rule in Sec 5.

The dip is usually vertical sometimes apparently N. as well as S. passing from the one to the other in either direction across the strike.


The quartzite is usually in massive beds although a very large percentage of it is more or less schistose. The massive beds are often as thick as two feet and usually enclose between them a thin layer of very fine quartzite schist indicating excessive adjustment in these narrow ~~beds~~ seams. The transition of the quartzite in this area into the jaspery slates to the N.E.

is gradual. To the south
of the blue line on the plat
it is mainly slates and almost
wholly slate to the N E. To
the south of the line is the
quartzite with here and there
narrow beds of cherty slate

S. $\frac{22}{28}$
34

T. 41

R. 14 W. 12.




23297 940 N 1040 W 27-41-14 W is
Outcrop of pink coarse massive
granite at edge of swamp

23298 400 N 865 W 34-41-14
Ledge of coarse granite.

S. 35

T. 41

R. 14



23299 40 N 165°0 W 35-41-14

Ledge of rather coarse greenstone or diabase which is cut by many small veins of granite like that to the west. Specimen shows contact of granite with the diabase.

23300 40 N 165°0 W 35-41-14

Specimen showing granite vein and diabase. Diabase is stretched in places. The granite seems to tail out and it lies in patches and veins in the diabase. Am not certain which is the cutting rock but it may be the granite.

23301 1835 N 102°0 W 35-41-14

Small ledge of greenstone formation composed of two kinds. 23301 is the finer grained and shows fluxion structure and is amygdaloidal.


23302

Same location. Specimen is of the coarser grained kind and looks much like 23299. Specimen contains fragments of the finer grained. It is difficult to tell which is the older but no fragments of the coarse grained was seen in the finer grained rock.

S. 26

T. 41

R. 14 E Wis



23303 560 N 1020 W 26-41-14

Ledge of fine grained massive
amygdaloidal greenstone, much
like 23301.

25/30
S. 36/31

T. 41
2

R. 11/15

26²

30

2

Brele River


38^{1/2}

31

S.

T. 41


R. 15+16 Niz



S. $\frac{27}{34} \mid \frac{26}{35} \mid \frac{25}{36}$

T. 41

R. / 5 -




23304 90° N 85° W 26 - 41 - 15

Sedge of reddish greenstone. Its redness is probably due to the weathering of calcite for it is filled with veins of calcite and also of quartz. On weathered surface it is grey with black spots.

S. 28

T. 41

R. 16 E Wis



28305 900 N 1600 W 28-41-16

Large ledge of pink quartzite in place conglomerate.

28306 875 N 1550 W 28-41-16

Specimen of dolomite or dolomitic quartzite in same ledge

28307 925 N 1620 W 28-41-16

specimen of quartzite conglomerate.

This quartzite ridge extends in a NW and SE direction with vertical walls on NE side and covered with drift or other material on the SW side. The pink quartzite 28305 and the conglomerate 28307 are very similar to the quartzite near the Brule River, in Range 35 on the Michigan side. But the dolomitic phase is new at this place.

The Dolomite forms the south east $\frac{1}{2}$ of the ridge and the quartzite and conglomerate is found only on the NW end.

No bedding structure could be made out. The rock is much shattered.

S. 31-32-33 T. 41

R. 16

6-1747

→

31

→

→

32

→


33

→

→

S. 34, 35, 36 T. 41


R. 16 E. N. 15



S. 33²

T. 41

R. 32




23308 965 N 235 W 3-41-32

Ledge of micaceous quartzite.

13 and
S. 5/8 of 8.24 " T. 42

R. 32 Mich



28309 1500 N 680 W 24-42-32

Ledge of dark grey micaceous quartzite. There is a system of fracturing running $N 40^{\circ} E$, Dip of fractured planes about vertical and a little to the S.E.

28310 1750 N 1720 W 24-42-32

Ledge of quartzite like the above but in this bedding can be made out as shown by banding of 28310. Strike of bedding $N 45^{\circ} W$ Dip about vertical as near as could be made out. Same fracturing as shown in the other ledge.

28311 230 N 580 W 13-42-32


Ledge of quartzite coarse grained like 28309 and one half of ledge very fine grained and shaly like 28311 which has undergone much crumpling Strike $N 70^{\circ} W$ Dip $66^{\circ} NE$

Ledges at 1000 and 1200 W on 300 N are like the other quartzite ledges, being coarse and fine grained the latter very much crumpled

14 and
S. $\frac{1}{4}$ of 23

T. 42

R. 32



28

23312 1000 N 730 W 14-42-32

Ledge of quartzite extending across the Paint River at the dam. This quartzite appears to have a great deal of chlorite and iron oxide in it. Weathers to a bright red. Strike $N 40^{\circ} W$. This place was visited by Van Hise and Bayley in July.

S. 18

T. 42


R. 31

18



23331

Granite




23331 200 N 1000 W 18-42-31

Large knot of greyish fine grained granite, with small flakes of mica. Ledge essentially like specimen.

S. 17

T. 42

R. 31



28325 975 N 75 W 17-42-31

The large ledge at this place is principally like specimen which contain a large quantity of hornblende. Gabbro or coarse diabase.

28382 975 N 1030 W 17-42-31

Ledge of gabbro. Its easternmost extension in this area.

28383 650 N 250 W 17-42-31

This large knot is made up of fine grained greenstone (gabbro? like specimen. Do not think it quartzite, - looks more like an eruptive. Average strike of schistosity N 55° W. Dip about 90°

28384 From same ledge as 28383.


The specimen looks like a breccia or conglomerate. In places the rock is crumpled across the strike.



S. 16

T. 42

R. 31



23313 1500 N 0 W 16-42-31

Ledge of fine grained micaceous quartzite, much squeezed and crumpled. 23313 is fairly coarse

23314 1500 N 0 W 16-42-31

Schistose variety, very micaceous and chloritic. There are a few squeezed veins of chert-like quartz in the schists. Strike of the schistosity which may be or may not be bedding N80°W. Dip about vertical.

23324 770 N 1500 W 16-42-31

The ledge at this place contains quite a large quantity of hornblende gabbro? But further south the large outcrop is a finer grained greenstone. Strike of schistosity ~~is~~ slightly N of W with a dip about 90° sometimes to the S but usually to the north.


23335 850 N 1800 W 16-42-31

The rock at this place is intermediate in texture between 23332 and 23333. Near this place what was taken to be greenstone bombs were found ~~Do Do~~

S. 15

T. 42

R. 31



23315 1400 N 1980 W 15-42-31

Continuation of the ledge of mica schists on the river. These fine schists alternate with the coarser quartzite schists but there is a predominance of the mica schists.

23316 1300 N 1750 W 15-42-31

Specimen is from a narrow strip of pyriteiferous and cherty rock. Opposite 1350 N 1750 W there is a working test pit known as Dickie's Exploration. Now down to a depth of 55 feet.

23317 1000 N 1500 W 15-42-31

The rock after alternating from that like 23315 and 23313 passes into a more massive variety like 23317. The specimen looks like a gabbro and if it is it must be an intrusion parallel to the schists or bedding. Strike of schistosity here is little South of West.

23318 1000 N 1480 W 15-42-31

This is finer grained than 23317 and contains more quartz.

- 28319 475-N 1200 W See 15-42-31
Ledge of gabbro. Specimen is
fine grained.
- 28820 450 N 1200 W 15-42-31
The greater part of the ledge is
coarse gabbro like specimen
- 28821 400 N 1025 W 15-42-31
A dyke of granite at the east
end of the above ledge. This
dyke is three or four feet in
thickness and cuts across the
gabbro in a slightly E of N direction
- 28822 500 N 875 W 15-42-31
Ledge of rock much like granite
yet much like 28820. This ledge
is on the river bank and seems
to be connected with the ledge
which contains 28319-20-21 but
no relations could be made
out.

Going north from this place
along the river the rock occurs
in high and rugged exposures
It is wholly the sedimentary
formation, alternating quartzite and
mica schist. Strike of schistosity or

bedding at 75°0 N 95°0 W is
N 80°0 E Dip 70° S.

28823 1750 N 2000 W 15-42-31

The northern part of this
large ledge is like specimen
a fine grained micaceous
quartzite.

28826 425-N 1050 W 15-42-31

From the dyke of granite in
the gabbro showing contact
between the two.

Specimens number 28832 to 28841
were gathered one forenoon with the
idea of working out in a more
satisfactory manner the relations
of the rock in this locality, in
sections 16 and 17 and northern part
of 21 and 20. Nos 28832-35 have
already been described and located.

28836. 600 N 1700 W 16-42-31

At this place a dyke of coarse gabbro
cuts the fine grained rock like
-35 and -33.

23337 400 N 1700 W 16-42-31

The ledge at this place is like specimen, fine grained and schistose. The planes of the schistosity curve about but have a general strike of $N 70^{\circ} W$.

23338 150 N 1650 W 16-42-31

The large high ledge at this place is much like -37. Strike is $N 70^{\circ} W$. Dip somewhat to the north.

23339 100 N 1600 W 16-42-31

At this place a dyke cuts the rock, running parallel to the strike of schistosity or bedding. This dyke is 5 or 6 feet in thickness in places. In places the dyke shows a zone of alteration minerals with the dyke rock sometimes on both sides of zone and sometimes the country rock on one side.

About 500 paces west of this dyke of diabase there is another similar dyke also running parallel to

the schistosity. There is also near this a dyke of granite which cuts very close in places to the greenstone dyke. The granite dyke cuts across the schistosity but the greenstone dyke runs parallel to the schistosity.



It appears to me that the granite dyke is the youngest rock and

that it cuts both the schists and the other dyke.


23340 Specimen from the granite dyke. About 200 W of SE cor of 17-42-31

23341 200 W of SE cor of 17-42-31 from near where 23340 was taken. At this place the ledge shows a large quantity of gabbro fragments squeezed into the schistose greenstone or quartzite. Specimen is a gabbro fragment in the schist. ~~Sketch of gabbro fragments~~. (Bayley ran an section line between 17+20 and 16+21.)

S. 36

T. 42

R. 32



23327 1200 N 1670 W 36-42-39


North end of rapids. Quite fine grained as well as coarse grained quartzite compose the ledges at this place.

Strike N 45° W Dip 75° S

S. 35

T. 42

R. 32



28828 560 N 500 W 36-42-32

At the south end of rapids
The quartzite is much like that
at the north end. Strike of
bedding N 70° W Dip about
 75° S.W. as near as could be
made out. No folding of
the beds back upon each
other was noticed.

S. 31

T. 42

R. 81


31

{ $\frac{11}{11}$ $\frac{11}{11}$ $\frac{11}{11}$ }

S. 32

T. 42

R. 31



23329 70 N 400 W 32-42-31

Continuous outcrops of mica schists on river bank. Strike is $N 75^{\circ} E$ Dip $75^{\circ} N$

23330 450 N 800 W 32-48-31


The rock at this place is more quartzose. At this place there is a high precipitous cliff on the river bank. The rapids known as Beaver Falls are at this place and further up the river. The rapids consist of rock alternating between 23329 and 23330

S. 1/4 of 20
and
Sec 29

T. 42

R. 31

20



28353 1260 N 200 W 29-42-31

On bank of river is outcrop of weathered pyroxenic gabbro like specimen.

28354 1500 N 200 W 29-42-31


The greater part of the knob is of coarse gabbro like the specimen

28355 Same location. Occurs in narrow vein in the above. It is a coarse gabbro containing much quartz. Shear zone?

28356 Same location. A fragment of quartzite included in the pyroxenic gabbro.

S. 21+28 T. 42

R. 31



23342 1450 N 600 W 28-42-31
Ledge of fine grained greenstone

23343 1460 N 350 W 28-42-31
Passing eastward over same
ledge come to a quite coarse
mass of gabbro.

23344 1340 N 200 W 28-42-31
At this place the rock is diabase
again like 23342.

23345 1325 N 150 W 28-42-31
At this place is a mass of
granite. The relations of the
granite, gabbro and fine grained
greenstone could not be made
out. Much the larger part
of ledge is fine grained
greenstone like 23342 and -44
The ledge is covered with drift
in places.

23349 325 W 1975 N 28-42-31
Ledge of quite coarse gabbro



23350 1975 N 350 W 28-42-31

Further west along same ridge
find fine grained greenstone, diabase
Relations of the two could not be
made out.

S. 22427 T. 42

R. 31

29



27

23346 1050 N 1640 W 27-42-31

At this place is a large hill of fine grained greenstone somewhat friable and soft. It is schistose with a strike of schistosity N and S.

23347 850 N 1900 W 27-42-31

Ledge of gabbro.

23348 0 N 80 W 22-42-31

Ledge of mica granite or gneiss.

(Plotted on preceding page)

23351 925 N 1100 W 28-42-31

Ledge like specimen on river bank.

23352 925 N 1450 W 28-42-31

Ledge of coarse gabbro containing veins of fine grained diabase. The relations of the two could not be made out. Sometimes one predominates and sometimes the other.

S. 23+26

T. 42

R. 31

23



26

S. 24+25-

T. 42

R. 31

24

11/11

25-


S. 357-36

T. 42

R. 31

35


36



S. 21+28

T. 42

R. 30



23357 860 N 630 W 21-42-30

Ledge of granite containing much gr. mica, on R.R. It contains patches of dark rock.

23358 Same location. Shows fragments or patches of darker fine grained granite.

23362 1650 N 655 W 28-42-30

Dyke of black hornblende rock which seems to cut the coarse granite like 23357.

S. 19420

T. 42

R. 31

