

## Tussi-Organidin advertisement.

[s.l.]: [s.n.], 1985

<https://digital.library.wisc.edu/1711.dl/AT4L7VEGFTNMQ9A>

<http://rightsstatements.org/vocab/InC/1.0/>

The libraries provide public access to a wide range of material, including online exhibits, digitized collections, archival finding aids, our catalog, online articles, and a growing range of materials in many media.

When possible, we provide rights information in catalog records, finding aids, and other metadata that accompanies collections or items. However, it is always the user's obligation to evaluate copyright and rights issues in light of their own use.

They all suppress coughs

Only One  
Moisturizes



## TUSSI-ORGANIDIN<sup>®</sup> with Codeine

LIQUID

Each teaspoonful (5 mL) contains: ORGANIDIN<sup>®</sup> (iodinated glycerol containing 15 mg organically bound iodine), 30 mg; codeine phosphate (WARNING: May be habit-forming), 10 mg.

## TUSSI-ORGANIDIN<sup>®</sup> DM

LIQUID

Each teaspoonful (5 mL) contains: ORGANIDIN<sup>®</sup> (iodinated glycerol containing 15 mg organically bound iodine), 30 mg; dextromethorphan hydrobromide, 10 mg.

Now, the only major Rx brands with an antitussive/expectorant combination

WALLACE LABORATORIES  
Div. of Carter-Wallace, Inc.  
Cranbury, N.J. 08512

Manufactured by  
Denver Chemical (Puerto Rico), Inc.  
Humacao, Puerto Rico 00661



WALLACE LABORATORIES  
Division of Carter-Wallace, Inc.  
Cranbury, New Jersey 08512

Please see the following page for a brief summary of prescribing information.

# TUSSI- ORGANIDIN<sup>®</sup> with Codeine

# TUSSI- ORGANIDIN<sup>®</sup> DM

*Before prescribing, please consult complete product information, a brief summary of which follows:*

**Indications and Usage:** For the symptomatic relief of irritating, nonproductive cough associated with respiratory tract conditions such as chronic bronchitis, bronchial asthma, tracheobronchitis, and the common cold; also for the symptomatic relief of cough accompanying other respiratory tract conditions such as laryngitis, pharyngitis, croup, pertussis and emphysema. Appropriate therapy should be provided for the primary disease.

**Contraindications:** History of marked sensitivity to inorganic iodides; hypersensitivity to any of the ingredients or related compounds; pregnancy; newborns; and nursing mothers.

**Warnings:** Discontinue use if rash or other evidence of hypersensitivity appears. Use with caution or avoid use in patients with history or evidence of thyroid disease.

**Precautions:** *General*—Iodides have been reported to cause a flare-up of adolescent acne. Children with cystic fibrosis appear to have an exaggerated susceptibility to the goitrogenic effects of iodides.

Dermatitis and other reversible manifestations of iodism have been reported with chronic use of inorganic iodides. Keep these in mind in patients receiving these preparations for prolonged periods.

*Drug Interactions*—Iodides may potentiate the hypothyroid effect of lithium and other antithyroid drugs.

*Carcinogenesis, Mutagenesis, Impairment of Fertility*—No long-term animal studies have been performed.

*Pregnancy*—Teratogenic effects: Pregnancy Category X (see CONTRAINDICATIONS).

*Nursing Mothers*—Do not administer to a nursing woman.

**Adverse Reactions:** Side effects have been rare, including those which may occur with the individual ingredients and which may be modified as a result of their combination.

*Organidin*—Gastrointestinal irritation, rash, hypersensitivity, thyroid gland enlargement, and acute parotitis.

*Codeine*—(Tussi-Organidin only): Nausea, vomiting, constipation, drowsiness, dizziness, and miosis.

*Dextromethorphan*—(Tussi-Organidin DM only): Drowsiness or gastrointestinal disturbances.

**Drug Abuse and Dependence** (Tussi-Organidin only):

*Controlled Substance*—Schedule V.

*Dependence*—Codeine may be habit-forming.

**Overdosage:** No reports of any serious problems.

**Dosage and Administration:** *Adults:* 1 to 2 teaspoonfuls every 4 hours.

*Children:* ½ to 1 teaspoonful every 4 hours.

**How Supplied:** *Tussi-Organidin Liquid*—clear red liquid, in bottles of one pint (NDC 0037-4812-10) and one gallon (NDC 0037-4812-20).

*Tussi-Organidin DM Liquid*—clear yellow liquid, in bottles of one pint (NDC 0037-4712-10) and one gallon (NDC 0037-4712-20).

**Storage:** Store at room temperature; avoid excessive heat. Keep bottle tightly closed.

Distributed by  
WALLACE LABORATORIES  
Division of  
CARTER-WALLACE, INC.  
Cranbury, New Jersey 08512

Manufactured by Denver Chemical (Puerto Rico) Inc.  
Humacao, Puerto Rico 00661

Rev. 4/84



**WALLACE LABORATORIES**  
Division of Carter-Wallace, Inc.  
Cranbury, New Jersey 08512