



LIBRARIES
UNIVERSITY OF WISCONSIN - MADISON

Dimetapp Extentabs advertisement.

[s.l.]: [s.n.], 1971

<https://digital.library.wisc.edu/1711.dl/HLXMHDLZUE5MA8M>

<http://rightsstatements.org/vocab/InC/1.0/>

The libraries provide public access to a wide range of material, including online exhibits, digitized collections, archival finding aids, our catalog, online articles, and a growing range of materials in many media.

When possible, we provide rights information in catalog records, finding aids, and other metadata that accompanies collections or items. However, it is always the user's obligation to evaluate copyright and rights issues in light of their own use.



head clear upon arising[†]

For upper respiratory allergies and infections including the common cold, Dimetapp Extentabs[®] effectively relieve the stuffiness, drip and congestion all night and all day long on just one Extentab every 12 hours. For most patients drowsiness or overstimulation is unlikely.

A-H-ROBINS

A. H. Robins Company
Richmond, Va. 23220

Dimetapp[®] Extentabs[®]

Dimetane[®] (brompheniramine maleate), 12 mg.; phenylephrine HCl, 15 mg.; phenylpropanolamine HCl, 15 mg.

prescribing information appears on next page

Dimetapp Extentabs®

INDICATIONS: Dimetapp Extentabs are indicated for symptomatic relief of allergic manifestations of upper respiratory illnesses, such as the common cold, seasonal allergies, sinusitis, rhinitis, conjunctivitis and otitis. In these cases it quickly reduces inflammatory edema, nasal congestion and excessive upper respiratory secretions, thereby affording relief from nasal stuffiness and postnasal drip.

CONTRAINDICATIONS: Hypersensitivity to antihistamines of the same chemical class. Dimetapp Extentabs are contraindicated during pregnancy and in children under 12 years of age. Because of its drying and thickening effect on the lower respiratory secretions, Dimetapp is not recommended in the treatment of bronchial asthma. Also, Dimetapp Extentabs are contraindicated in concurrent MAO inhibitor therapy.

WARNINGS: *Use in children:* In infants and children particularly, antihistamines in overdosage may produce convulsions and death.

PRECAUTIONS: Administer with care to patients with cardiac or peripheral vascular diseases or hypertension. Until the patient's response has been determined, he should be cautioned against engaging in operations requiring alertness such as driving an automobile, operating machinery, etc. Patients receiving antihistamines should be warned against possible additive effects with CNS depressants such as alcohol, hypnotics, sedatives, tranquilizers, etc.

ADVERSE REACTIONS: Adverse reactions to Dimetapp Extentabs may include hypersensitivity reactions such as rash, urticaria, leukopenia, agranulocytosis and thrombocytopenia; drowsiness, lassitude, giddiness, dryness of the mucous membranes, tightness of the chest, thickening of bronchial secretions, urinary frequency and dysuria, palpitation, hypotension/hypertension, headache, faintness, dizziness, tinnitus, incoordination, visual disturbances, mydriasis, CNS-depressant and (less often) stimulant effect, anorexia, nausea, vomiting, diarrhea, constipation, and epigastric distress.

HOW SUPPLIED: Light blue Extentabs in bottles of 100 and 500.