



LIBRARIES
UNIVERSITY OF WISCONSIN - MADISON

Lomodex 70 advertisement.

[s.l.]: [s.n.], 1974

<https://digital.library.wisc.edu/1711.dl/KASRKP3334WRF8J>

<http://rightsstatements.org/vocab/InC/1.0/>

The libraries provide public access to a wide range of material, including online exhibits, digitized collections, archival finding aids, our catalog, online articles, and a growing range of materials in many media.

When possible, we provide rights information in catalog records, finding aids, and other metadata that accompanies collections or items. However, it is always the user's obligation to evaluate copyright and rights issues in light of their own use.

THE CHOICE

"... Dextran 70 and subcutaneous heparin seem to have a comparable effect in prevention of pulmonary embolism."¹

Lomodex 70 (Dextran 70 Injection B.P.) is easily administered and is effective during the critical 48 hours following surgery. There is little risk of bleeding or haematoma formation during surgery and far less post-operative ward work.

Heparin, however, requires up to 21 injections both pre- and post-operatively, and has the added disadvantages of extra ward work, constant laboratory monitoring and appreciable patient discomfort.

"... Dextran 70 has the advantage that it is much more acceptable to patients and is less demanding of the nursing staff."²

Lomodex 70 is the practical choice in prophylaxis of thromboembolism.

1. *Brit med J*, (1975) ii, 109.
2. *J Obs & Gynae BC*, (1974) 81, 486.

Lomodex[®] 70
the practical choice

FISONS

Further information is available on request from:
Fisons Limited, Pharmaceutical Division,
Loughborough LE11 0BB, Leicestershire.

