



**LIBRARIES**  
UNIVERSITY OF WISCONSIN - MADISON

## Rondec-DM advertisement.

[s.l.]: [s.n.], 1982



<https://digital.library.wisc.edu/1711.dl/7HOHGKVYCR2OU8D>

<http://rightsstatements.org/vocab/InC/1.0/>

The libraries provide public access to a wide range of material, including online exhibits, digitized collections, archival finding aids, our catalog, online articles, and a growing range of materials in many media.

When possible, we provide rights information in catalog records, finding aids, and other metadata that accompanies collections or items. However, it is always the user's obligation to evaluate copyright and rights issues in light of their own use.

# For Big Colds in small people



## RONDEC-DM™

### Drops

**R<sub>x</sub>** (each dropperful (1 ml) contains:  
carbinoxamine maleate, 2 mg; pseudo-  
ephedrine HCl, 25 mg; dextromethorphan  
HBr, 4 mg; less than 0.6% alcohol)

For infants to 18 months

## RONDEC-DM™

### Syrup

**R<sub>x</sub>** (each teaspoonful (5 ml) contains:  
carbinoxamine maleate, 4 mg; pseudo-  
ephedrine HCl, 60 mg; dextromethorphan  
HBr, 15 mg; less than 0.6% alcohol)

For children 18 months and older

### WITH DEXTROMETHORPHAN HBr:

- Especially suited for pediatric patients
- Relieves the cough without narcotic side effects and without suppressing respiration or ciliary activity.
- Drops contain calibrated dropper for ease of administration.

for adults and children

## RONDEC-DM™ Syrup

**R<sub>x</sub>** each teaspoonful (5 ml) contains carbinoxamine maleate, 4 mg; pseudoephedrine hydrochloride, 60 mg; dextromethorphan hydrobromide, 15 mg; less than 0.6% alcohol.

for infants

## RONDEC-DM™ Drops

**R<sub>x</sub>** each dropperful (1 ml) contains carbinoxamine maleate, 2 mg; pseudoephedrine hydrochloride, 25 mg; dextromethorphan hydrobromide, 4 mg; less than 0.6% alcohol.

### INDICATIONS AND USAGE

For symptomatic relief of the common cold, nasopharyngitis with post-nasal drip, bronchitis and related respiratory conditions.

### CONTRAINDICATIONS

Patients with hypersensitivity or idiosyncrasy to any ingredients, patients taking monoamine oxidase (MAO) inhibitors, patients with narrow-angle glaucoma, urinary retention, peptic ulcer, severe hypertension or coronary artery disease, or patients undergoing an asthmatic attack.

### WARNINGS

*Use in Pregnancy:* Safety for use during pregnancy has not been established.

*Nursing Mothers:* Use with caution in nursing mothers.

*Special Risk Patients:* Use with caution in patients with hypertension or ischemic heart disease, and persons over 60 years.

### PRECAUTIONS

Before prescribing medication to suppress or modify cough, identify and provide therapy for the underlying cause of cough.

Use with caution in patients with hypertension, heart disease, asthma, hyperthyroidism, increased intraocular pressure, diabetes mellitus and prostatic hypertrophy.

*Information for Patients:* Avoid alcohol and other CNS depressants while taking these products. Patients sensitive to antihistamines may experience moderate to severe drowsiness. Patients sensitive to sympathomimetic amines may note mild CNS stimulation. While taking these products, exercise care in driving or operating appliances, machinery, etc.

*Drug Interactions:* Antihistamines may enhance the effects of tricyclic antidepressants, barbiturates, alcohol, and other CNS depressants. MAO inhibitors prolong and intensify the anticholinergic effects of antihistamines. Sympathomimetic amines may reduce the antihypertensive effects of reserpine, veratrum alkaloids, methyldopa and mecamlamine. Effects of sympathomimetics are increased with MAO inhibitors and beta-adrenergic blockers. The cough suppressant action of dextromethorphan and narcotic antitussives are additive.

*Pregnancy Category C:* Animal reproduction studies have not been conducted with Rondec-DM. It is also not known whether these products can cause fetal harm when administered to a pregnant woman or affect reproduction capacity. Give to pregnant women only if clearly needed.

### ADVERSE REACTIONS

**Antihistamines:** Sedation, dizziness, diplopia, vomiting, dry mouth, headache, nervousness, nausea, anorexia, heartburn, weakness, polyuria and dysuria and, rarely, excitability in children.

**Sympathomimetic Amines:** Convulsions, CNS depression, cardiac arrhythmias, respiratory difficulty, increased heart rate or blood pressure, hallucinations, tremors, nervousness, insomnia, weakness, palpitation and dysuria.

Dextromethorphan: Drowsiness and GI disturbance.

### DOSAGE AND ADMINISTRATION

#### DOSAGE:

AGE	DOSE*	FREQUENCY*
Rondec-DM Syrup		
18 months-6 years	½ teaspoonful (2.5 ml)	q.i.d.
adults and children		
6 years and over	1 teaspoonful (5 ml)	q.i.d.
Rondec-DM Drops		
for oral use only		
1-3 months	¼ dropperful (¼ ml)	q.i.d.
3-6 months	½ dropperful (½ ml)	q.i.d.
6-9 months	¾ dropperful (¾ ml)	q.i.d.
9-18 months	1 dropperful (1 ml)	q.i.d.

\*In mild cases or in particularly sensitive patients, less frequent or reduced doses may be adequate.

### HOW SUPPLIED

Rondec-DM Syrup, grape-flavored, in 16-fl-oz (1-pint) bottles, NDC 0074-5640-16; and 4-fl-oz bottles, NDC 0074-5640-04. Dispense in USP tight, light-resistant, glass container. Avoid exposure to excessive heat.

Rondec-DM Drops, grape-flavored, in 30-ml bottles for dropper dosage. Calibrated, shatterproof dropper enclosed in each carton. Container meets safety closure requirements. NDC 0074-5639-30. Avoid exposure to excessive heat.