



LIBRARIES
UNIVERSITY OF WISCONSIN - MADISON

Deca-Durabolin advertisement.

[s.l.]: [s.n.], 1963

<https://digital.library.wisc.edu/1711.dl/XBW5WLQBL2ARW8U>

<http://rightsstatements.org/vocab/InC/1.0/>

The libraries provide public access to a wide range of material, including online exhibits, digitized collections, archival finding aids, our catalog, online articles, and a growing range of materials in many media.

When possible, we provide rights information in catalog records, finding aids, and other metadata that accompanies collections or items. However, it is always the user's obligation to evaluate copyright and rights issues in light of their own use.

to help return
the chronically
ill patient to
a life of more
useful activity

with the economy and convenience of monthly injections.

A course of anabolic therapy with DECA-DURABOLIN (nandrolone decanoate) not only helps hasten recovery, but also helps insure the continuous integrity of vital protein tissues. Moreover, it exerts an effective sustained anti-catabolic action with a series of four monthly injections or, where prolonged therapy is required, with as few as eight injections a year.

The debilitated, the chronically ill and the geriatric patient undergoing excessive tissue breakdown often can be returned to positive nitrogen balance with resulting tissue growth and repair.


DECA-DURABOLIN (nandrolone decanoate) is preferred among anabolics because it does not interfere with normal liver function and because virilizing side effects are notably infrequent and mild.

Average Adult Dose: 50 to 100 mg. I.M. every 3 to 4 weeks for 12 weeks. If necessary, therapy may be repeated following a rest period of 4 to 8 weeks.

Precautions and Contraindications: The primary conditions should be diagnosed and treated, and improper diet or eating habits corrected to insure adequate nutrients. Women and children are more sensitive to androgenic stimulation and should be watched for possible signs of virilization (usually mild and reversible when dosage is decreased or therapy discontinued). Large doses or intensive therapy may temporarily inhibit menses. In patients with pre-existing liver disturbances therapy should be discontinued if they do not react favorably. Patients (especially malnourished or anemic) with impaired cardiac and renal function should be closely observed because of the possibility of increased retention of sodium and water. The drug is contraindicated in prostatic cancer or serious cardio-renal failure. It is not recommended for use during pregnancy.

Supplied: 50 mg./cc. in sterile sesame oil with 10% benzyl alcohol (preservative)—1 cc. ampuls, box of 4, or 2 cc. multi-dose vials.



DECA-DURABOLIN[®]
(Nandrolone Decanoate) 

Organon Inc., West Orange, N. J.