

Brondecon Elixir advertisement.

[s.l.]: [s.n.], 1973

<https://digital.library.wisc.edu/1711.dl/2B6QRM07IBSBG8T>

<http://rightsstatements.org/vocab/InC/1.0/>

The libraries provide public access to a wide range of material, including online exhibits, digitized collections, archival finding aids, our catalog, online articles, and a growing range of materials in many media.

When possible, we provide rights information in catalog records, finding aids, and other metadata that accompanies collections or items. However, it is always the user's obligation to evaluate copyright and rights issues in light of their own use.



Sometimes,
breathing isn't
easy.

Brondecon Elixir can control her bronchospasm and help make her cough more productive... breathing becomes less difficult.

Oxtriphylline, the bronchodilator component, is usually well-tolerated by children and does not cause the gastric upset associated with other forms of theophylline.

Glyceryl guaiacolate, a time-tested expectorant, helps liquefy mucoid secretions and facilitates freer breathing.

Brondecon Elixir is a valuable adjunct in the treatment of pediatric bronchitis, bronchial asthma and asthmatic bronchitis.

And children like the cherry flavor — there's seldom a problem getting them to take it.

Breathing becomes easier with

BRONDECON® ELIXIR

EACH 5 ML TEASPOONFUL CONTAINS 100 MG OXTRIPHYLLINE AND 50 MG GLYCERYL GUAIACOLATE; ALCOHOL 20%.

BRONCHODILATOR/EXPECTORANT

Caution: Federal law prohibits dispensing without prescription.

Each tablet contains 200 mg oxtriphylline and 100 mg glyceryl guaiacolate. Each 5 ml teaspoonful of elixir contains 100 mg oxtriphylline and 50 mg glyceryl guaiacolate; alcohol 20%.

Indications: Brondecon (oxtriphylline and glyceryl guaiacolate) is an adjunct in the management of bronchitis, bronchial asthma, asthmatic bronchitis, pulmonary emphysema, and similar chronic obstructive lung disease. It is indicated when both relaxation of bronchospasm and expectorant action are desirable.

Precautions: Concurrent use of other xanthine preparations may lead to adverse reactions, particularly CNS stimulation in children.

Adverse Reactions: Gastric distress and, occasionally, palpitation and CNS stimulation have been reported.

Dosage: Tablets — over 12 years of age: one tablet, 4 times a day.

Elixir — over 12 years of age: two teaspoonfuls, 4 times a day; from 2 to 12 years: one teaspoonful per 60 lb body weight, 4 times a day.

Supplied: 200 mg salmon pink tablets in bottles of 100. Elixir, dark red, cherry-flavored in 237 ml (8 fl oz) and 474 ml (16 fl oz) bottles.

Full information is available on request.



WARNER/CHILCOTT

Div. Warner-Lambert Company
Morris Plains, N.J. 07950