



LIBRARIES
UNIVERSITY OF WISCONSIN - MADISON

Otrivin Pediatric advertisement.

[s.l.]: [s.n.], 1973

<https://digital.library.wisc.edu/1711.dl/L7GAIXMUORMZU8V>

<http://rightsstatements.org/vocab/InC/1.0/>

The libraries provide public access to a wide range of material, including online exhibits, digitized collections, archival finding aids, our catalog, online articles, and a growing range of materials in many media.

When possible, we provide rights information in catalog records, finding aids, and other metadata that accompanies collections or items. However, it is always the user's obligation to evaluate copyright and rights issues in light of their own use.





It even treats



kids under six to prolonged decongestion.

Nose drops gentle enough to treat kids under six may not be long acting enough. And other nose drops that may be long acting are limited to use in older children.

Otrivin Pediatric can provide long-lasting relief for kids of all ages, including infants.

Just one application can keep noses clear for more than six hours — two or three applications can keep them clear all day.

And Otrivin works gently. It seldom causes stinging, burning, or the drowsiness that can affect school or play. Also available for older patients: Otrivin Spray.

Otrivin® Pediatric
xylometazoline HCl NF
0.05% solution

Long on action,
short on doses.

Geigy

Indications: For decongestion of the nasal mucosa.
Contraindications: Narrow-angle glaucoma. Concurrent MAO inhibitor therapy. Tricyclic antidepressant therapy. Hypersensitivity to any component. Sensitivity to even small doses of adrenergic substances as manifested by sleeplessness, dizziness, light-headedness, weakness, restlessness, or cardiac arrhythmias. **Warnings:** Systemic effects may occur if given in excess. They are most likely to occur in infants and the elderly. Use very cautiously in patients with hypertension, heart disease, angina, hyperthyroidism and advanced arteriosclerotic conditions. Overdosage may produce profound CNS depression in children, possibly requiring intensive supportive treatment. Usage in pregnancy: Safe use of Otrivin brand of xylometazoline hydrochloride has not been established with respect to fetal development. It should not be used in women of childbearing potential or during early pregnancy unless the potential benefits outweigh the possible hazards. **Precautions:** Use sparingly in all patients, particularly infants, children and the elderly. **Adverse Reactions:** Burning, stinging, dryness of the nasal mucosa, sneezing, palpitations, tachycardia, cardiac arrhythmias, increase in blood pressure, headache, blurred vision, drowsiness, nervousness, insomnia, prolonged vision, drowsiness, CNS depression. Prolonged or excessive use may cause rebound congestion. (B) 6802 (6/74)
For complete details, including dosage, please see full prescribing information.

GEIGY Pharmaceuticals
Division of CIBA-GEIGY Corporation
Ardsley, New York 10502