



**LIBRARIES**  
UNIVERSITY OF WISCONSIN - MADISON

## Novahistine advertisement.

[s.l.]: [s.n.], 1956

<https://digital.library.wisc.edu/1711.dl/XUZPMPVC2QLOV8Y>

<http://rightsstatements.org/vocab/InC/1.0/>

The libraries provide public access to a wide range of material, including online exhibits, digitized collections, archival finding aids, our catalog, online articles, and a growing range of materials in many media.

When possible, we provide rights information in catalog records, finding aids, and other metadata that accompanies collections or items. However, it is always the user's obligation to evaluate copyright and rights issues in light of their own use.

Fastest gun...



when his nose  
don't run!



Little cold sufferers  
will get **GREATER RELIEF**  
with

## Novahistine<sup>®</sup> Elixir

Novahistine Elixir combines the action of a quick-acting sympathomimetic with an antihistaminic drug for a greater decongestive effect.

Each 5 cc. teaspoonful contains:  
Phenylephrine hydrochloride . . . . . 5.0 mg.  
Prophepridine maleate . . . . . 12.5 mg.  
Chloroform (approx.) . . . . . 13.5 mg.  
l-Menthol . . . . . 1.0 mg.  
(Alcohol, 5%)

**Dosage:** Children—1 teaspoonful, three or four times daily. Infants under 1 year— $\frac{1}{4}$  to  $\frac{1}{2}$  teaspoonful, three or four times a day.

Supplied in 4 oz., pint and gallon bottles.

**PITMAN-MOORE COMPANY**

DIVISION OF ALLIED LABORATORIES, INC.  
INDIANAPOLIS 6, INDIANA