



LIBRARIES
UNIVERSITY OF WISCONSIN - MADISON

Eskatrol advertisement.

[s.l.]: [s.n.], 1960

<https://digital.library.wisc.edu/1711.dl/A6QHBNKPOTH3K9D>

<http://rightsstatements.org/vocab/InC/1.0/>

The libraries provide public access to a wide range of material, including online exhibits, digitized collections, archival finding aids, our catalog, online articles, and a growing range of materials in many media.

When possible, we provide rights information in catalog records, finding aids, and other metadata that accompanies collections or items. However, it is always the user's obligation to evaluate copyright and rights issues in light of their own use.

an escape
from
stress?



Many people overeat as an escape from stress. Whenever they are particularly upset, tense or frustrated, they devour food. The result is usually overweight.

'Eskatrol' *Spansule* capsules both control appetite and relieve the emotional stress that causes overeating. That's why so many patients are losing more pounds with 'Eskatrol'.

Eskatrol*

Each capsule contains Dexedrine* (brand of dextro amphetamine sulfate), 15 mg., and Compazine* (brand of prochlorperazine), 7.5 mg., as the maleate.

Spansule*

brand of sustained release capsules

because emotions play an
important role in overweight

RECOMMENDED DOSAGE: One 'Eskatrol' *Spansule* capsule daily, taken in the morning.

BRIEF SUMMARY OF PRINCIPAL SIDE EFFECTS AND CAUTIONS

SIDE EFFECTS: Side effects (chiefly nervousness and insomnia) are infrequent, and usually mild and transitory.

CAUTIONS: Clinical experience has demonstrated that 'Eskatrol' (containing the phenothiazine derivative, prochlorperazine) has a wide margin of safety and that there is little likelihood of blood or liver toxicity or neuromuscular reactions (extrapyramidal symptoms). The physician should be aware, however, of their possible occurrence.

'Eskatrol' *Spansule* capsules should be used with caution in the presence of severe hypertension, advanced cardiovascular disease, or extreme excitability. Excessive use of the amphetamines by unstable individuals may result in a psychological dependence; in these rare instances withdrawal of medication is recommended. It is generally recognized that in pregnant patients all medications should be used cautiously, especially in the first trimester.

SUPPLIED: Bottles of 50 capsules.

*Registered trademark

Smith Kline & French Laboratories

