

Milpath advertisement.

[s.l.]: [s.n.], 1962

https://digital.library.wisc.edu/1711.dl/WG6WPPETJXO3C8D

http://rightsstatements.org/vocab/InC/1.0/

The libraries provide public access to a wide range of material, including online exhibits, digitized collections, archival finding aids, our catalog, online articles, and a growing range of materials in many media.

When possible, we provide rights information in catalog records, finding aids, and other metadata that accompanies collections or items. However, it is always the user's obligation to evaluate copyright and rights issues in light of their own use.

Like a vacation in the sun, 'Milpath' brightens the horizon of the worry-nerve ulcer patient

promptly relieves pain, spasm, hypermotility, hypersecretion...

Milpath

meprobamate + tridihexethyl chloride Product Information: 'Milpath' (meprobamate + tridihexethyl chloride) is indicated in both organic and functional gastrointestinal disorders, especially when accompanied by anxiety and tension.

Side effects from tridihexethyl chloride are rare at recommended dosage; after high dosage, dry mouth and blurred vision, and occasional constipation or "bloating" may occur. Slight drowsiness and, rarely, allergic reactions may occur with meprobamate. Meprobamate may increase effects of excessive alcohol. Consider possibility of dependence, particularly in patients with history of drug or alcohol addiction. Withdraw gradually after prolonged use at high dosage. Tridihexethyl chloride is contraindicated in urinary bladder neck or pyloric obstruction, and glaucoma; may also be contraindicated in peptic ulcers with significant gastric retention due to stenosis.

Usual Dosage: One 'Milpath'-400 or 1-2 'Milpath'-200 tablets q.i.d. at meals and bedtime.

Composition:

'Milpath'-400:	Each tablet	contai	ns:	
meprobamate			400	mg.
tridihexethyl o	chloride		25	mg.
'Milpath'-200:	Each tablet	contai	ns:	
meprobamate			200	mg.
tridihexethyl o	chloride		25	mg.

