



Monodral advertisement.

[s.l.]: [s.n.], 1955

<https://digital.library.wisc.edu/1711.dl/CZEF54V2N3W4K9C>

<http://rightsstatements.org/vocab/InC/1.0/>

The libraries provide public access to a wide range of material, including online exhibits, digitized collections, archival finding aids, our catalog, online articles, and a growing range of materials in many media.

When possible, we provide rights information in catalog records, finding aids, and other metadata that accompanies collections or items. However, it is always the user's obligation to evaluate copyright and rights issues in light of their own use.



Functional and Organic Control

of PEPTIC ULCER

Gastro-intestinal
Irritability and Tension

MONODRAL® with MEBARAL® TABLETS

Potent ANTISECRETORY • ANTICHOLINERGIC • SEDATIVE

Each tablet contains:

Monodral bromide 5 mg.
Mebaral 32 mg.

PROVIDES:

Dependable control of hyperacidity and hypermotility. Spasmolysis. Prompt and prolonged pain relief and tranquillity.

DOSE:

Peptic ulcer, 1 or 2 tablets three or four times daily. Other gastro-intestinal disorders, 1 tablet three or four times daily.

SUPPLIED: Bottles of 100 tablets.

For unsurpassed results in PEPTIC ULCER
prescribe Monodral with Mebaral in conjunction
with

Creamalin®

PIONEER
ALUMINUM
HYDROXIDE GEL

- Fast Acting Reactive Gel
- Protective Coating

Creamalin liquid — 8 and 16 fl. oz.
Creamalin tablets — bottles of 50 and 200.

Winthrop
LABORATORIES
NEW YORK 18, N. Y.

Monodral (brand of penthenate),
Mebaral (brand of mephobarbital), and
Creamalin, trademarks reg. U. S. Pat. Off.