



LIBRARIES

UNIVERSITY OF WISCONSIN-MADISON

Central Missouri: [specimens] 15001-15259. No. 79 1888

Haworth, Erasmus

[s.l.]: [s.n.], 1888

<https://digital.library.wisc.edu/1711.dl/QHUABQ7OB4YDU8A>

<http://rightsstatements.org/vocab/InC/1.0/>

For information on re-use see:

<http://digital.library.wisc.edu/1711.dl/Copyright>

The libraries provide public access to a wide range of material, including online exhibits, digitized collections, archival finding aids, our catalog, online articles, and a growing range of materials in many media.

When possible, we provide rights information in catalog records, finding aids, and other metadata that accompanies collections or items. However, it is always the user's obligation to evaluate copyright and rights issues in light of their own use.

U. S. GEOLOGICAL SURVEY
FIELD SECTION BOOK

No. 79

1888

Central Missouri

Erasmus Haworth

15001 - 15258

SURVEY OF THE PRE-CAMBRIAN ROCKS OF THE N. W. STATES.

INSTRUCTIONS.

1. Ordinarily at least two pages of this book will be devoted to one section. On the left hand page place a map of as much of the section as has *actually been seen*. Denote rivers, lakes, marshes, etc., by the usual topographical signs. Denote the ledges of rock, when no structure is made out, by cross-hatching, making the cross-hatching cover as nearly as possible the areas occupied by the exposures. If the rock is a massive one, but still more or less plainly bedded, use the same sign with a dip arrow and figure attached, showing the amount and inclination of the dip. Denote slaty or other very plainly bedded rocks by lines running in the direction of the strike, with figures and a dip arrow attached as before. In all cases where there is the least doubt about the true bedding directions, indicate it by a query. To each exposure on the face of the map attach the number of the specimen representing it. In mapping the section count each of the spaces between the blue lines as 100 paces, and twenty of these spaces as one mile, or 2,000 paces. Usually the southeast corner will be placed at the bottom of the page, or at the first black line above the bottom of the page, and at the right hand side. If, however, for any reason, it is desirable to show portions of an adjoining section, the southeast corner may be shifted up, or the map may be turned around and the north placed at the left hand side of the page.

2. On the right hand page place the notes descriptive of the exposures. Begin in each case with the number of the specimen, placing the number on the left hand side of the red line, after which give in order on the right of the same red line the position of the ledges as reckoned in paces from the southeast corner of the section, and the dip and strike when observable, for instance: 4025 250 N., 300 W., *Strike, N. 6° E., Dip, 50° E.* Then follow with as full a description of the ledge as possible.

3. The ruling of the left hand page is also arranged so that a smaller scale can be used. Each one of the black lines may represent a section line and the red lines quarter sections and "forties." The scale of the maps may thus be reduced, if desirable, to two inches to the mile (the ordinary town plat scale.)

4. Collect a specimen from each separate ledge of rock, or wherever there is a change of rock on any one ledge. In case of trips made on foot or in canoes, for long distances, neighboring ledges, unquestionably of one kind of rock, need not be sampled, the position and extent of the ledge being marked on the map, with a note that it is of a rock identical with specimen so-and-so. Under the same conditions small sized samples, trimmed to a uniform size of $2 \times 2\frac{1}{2} \times \frac{3}{4}$ inches will be allowed, but in all other cases *large sized specimens*, trimmed to a size of $3 \times 4 \times 1$ inches, must be selected, in accordance with § 3, chapter IV, p. 44, Regulations of the U. S. Geological Survey. In all cases collect chips for slicing. All specimens are to have numbers painted on them, in white on a black background, in the field.

5. On the last twenty-five pages of the book give, as may seem desirable, a general account of the examination of the region mapped in the previous pages, correlation of observations, sketches, cross sections, etc., etc.

#79

For convenience and accuracy the colors shown on the opposite page will be used in this note book. Contact lines between any two where traced accurately will be represented by an unbroken black line thus ———, but where there is any uncertainty regarding the exact location of boundaries the division line will be represented by a broken line thus - - - - -.



= Granite.




= Porphyry.



= Limestone.



= Sandstone.



= Dyke-rock.


F.

= Flint-rock, or chert.

$\frac{1}{2}$

T. 22, A.

R. 5, E. Section 5



1

~~15022~~ This specimen was taken from the porphyry bluff at the foot of a high hill, and is a fair representative of the neighboring porphyry.
See page 9

15023 The specimens were taken from the line of contact between the granite and porphyry. The contact line seems to be pretty plainly marked if one makes a superficial observation; but to find the exact division between the two is not easy. Only a few feet on either side of the line we find characteristic rocks.

15025 Granite, taken from the right bank of the little branch. Large exposures of granite occur here.


15026-27 This is a very interesting exposure of coarse sand-stone or conglomerate. Porphyry boulders as big as a man's head, and considerably water worn are very abundant in the rock, showing that this conglomerate is younger than the neighboring
See P. 56. & K. 104. 1/2. 1/2.

1 1/2

T. 33ch

R. 5. E. Section 6.

North half of Section 6.



15038 These specimens were taken from the con-
 tact line between the granite and porphyry,
 to on the north slope of "Bunker's Hill," the
 15041 location of which is shown on the map.
 To the east the granite is very abundant, and
 many large boulders from 4 to 8 ft. in diameter
 are scattered over the surface, evidently
 never having been very far removed. The di-
 vision line, at first sight, seems to be quite
 plain, but careful examination in some
 places showed that there was no seam,
 or fissure, between the two kinds of rock.
 15038 was taken so as to show the two
 phases. 39 is porphyry 6 in. away, 40
 is porphyry 15 in. away, and 41 is porphy-
 ry 40 ft. away.

15138 This locality seemed to be so interesting
 & that I visited it a number of times, and
 15139 finally gathered a new set of specimens.
 38 is granite and 39 is porphyry from
 just across the creek to the north. They
 were about 27 yds. apart. The bank on the
 north side is quite low and the rock
 exposures are not abundant. On the south
 side there is a high bluff reaching fully
 300 ft. at a point about 300 yds. up the
 stream from the contact line.


See TP 52

2 1/2

T. 33. N.

R. 5. E. Section 6.

South half of Section 6.



15053

Along the ravine at a place from $\frac{1}{4}$ to $\frac{1}{2}$ mi.

✓

15054

south of the north line of the map there is a very peculiar looking rock. It is of almost ashy whiteness and seems to have been originally a coarse conglomerate composed of porphyry boulders and porphyry gravel. As these boulders decayed shells of decayed matter would be formed. In many instances the central part of the boulders are comparatively sound. 534, 54 are samples of this rock.

15055

This is a porphyry which fairly well represents the porphyry on the hills at the head of the ravine. There is nothing peculiar about it or its mode of occurrence which would distinguish it from other porphyries.

15056

Porphyry taken from the summit of a high hill known as "Compton Mt." The whole of the country to the east, south and west is a mass of porphyry hills, with but little difference in the character of the rocks, so far as their macroscopic appearances are concerned.

3 1/2

T. 33. N.

R. 5. E. Section 5.

North half of Section 5.



15037

This specimen of granite came from the north bank of Spout's creek about $\frac{1}{2}$ mi. below the old Tension mill. There are heavy exposures of granite for a considerable distance up and down the creek. This would be a splendid place to open up a quarry. In the summer of '86 a little work was done here, but railroad facilities were so poor the work was stopped.

15050

This granite specimen is from the roadside by the school-house on the "Gravel Road." The granite extends farther north here than in some other places.

15051

This is about the same as '50.


The area on this map north of Spout's creek represented as a granite area is generally covered with soil. But in a great many places the rocks are exposed, and in others granite boulders cover the surface, so there can be no doubt whatever with reference to the nature of the underlying rock.

4 1/2

T. 33. N.

R. 5. E. Section 4.

North half of Section 4.



15052 This specimen is a dyke rock. It was found in a small ravine which runs through Mr. McAllen's field on the north side of Shout's creek. The dyke itself could not be found, but the boulders were very numerous, and were so large that the dyke must have been fully 2 ft. thick, and probably much more.

15053a This specimen is granite. It was taken from the left bank of Shout's creek just at the point where the creek leaves the bluff and bears east into the St. Francois. The granite here were unusually fissured into angular blocks like the porphyries, but otherwise seem to be good granite.

The portion of this section which lies to the east of St. Francois was not examined in detail, and therefore is not colored on the map.

642

T. 33. N.

R. 5. E. Section 4.

South half of Section 4.
→

15066

15067

Rock Creek

15066

7

This specimen is a granite taken from the east side of Mr. McAllens field. I could find no ledge and consequently took a specimen from a big boulder in the ravine.

The whole of this section, excepting the small area in the E. N. corner is in the granite area, and although it is generally covered by soil the granite boulders and occasional out-croppings of the solid granite leave no reasonable doubt that the whole of the are is underlaid with granite.


15067

This is dyke rock taken from the ravine just south of the house of John Bank. The dyke is in the solid granite, is at least 8 or 4 ft. wide, and may be more. It trends N. E. and S. W. Just west of it about 2 rods, another dyke crosses the ravine with the same trend. The whole country through here seems to be full of small dykes, especially to the S. E. of this place. Usually nothing can be seen excepting the boulders which lie on the surface. In every case yet observed in which the trend of the dyke could be determined it was N. E. and S. W.

7 1/2

T. 33. N.

R. 5. E. Section 9



15108

E

15128

Just on the west bank of Rock creek 30 yds. ⁸
 above the contact line between the granite and
 porphyry, as shown on the map, a shaft
 has been sunk 30 ft. deep. It is now filled
 up to a depth of $18\frac{1}{2}$ ft. below the surface.
 I had the water dipped out and collected
 specimens from different places along the
 walls. This is a most interesting place. We
 have what seems to be a change from the
 non-crystalline to the crystalline condition
 and ~~the~~ by gradual transitions, and the
 whole can readily be seen by examining
 the walls of the shaft. There is an absence
 of any appearance of intrusive rock.
 The changes are much more rapid than
 one would have suspected, usually being
 quite complete within 2 ft. or less. At the
 N.E. corner near the surface is one of the
 coarse spots. Immediately below it the rock
 is not macroscopically crystalline, but a few
 feet lower the coarseness of grain again be-
 comes apparent, and at the depth of 18 ft.
 the rocks on the north, west, and south sides
 are quite coarse. Specimens were gathered as
 follows

15108, Klein shaft, middle of N. wall 3 ft. 8 in.


15109, Adjoining 108 on the E.

See P. 54

8 1/2

T. 33. N.

R. 5. E. Section 8




15022

This section is wholly occupied by Rock⁹
creek Int. The rocks throughout the whole section
are porphyry, and there is nothing to contrast
with the portions of the hill which extend
into other sections.

9 1/2

T. 33. N.

R. 5. E. Section 7.




15021

This specimen is porphyry taken about ^{10.}
75 yds. up the hillside from the ravine.
The whole country around is solid porphyry.
I traveled through the eastern portion of
the section, along the northern line and
also the western side but did not pass
directly over the summit of the hill.
The hill is locally called the Blankenship
Mk.

10 1/2

T. 33. N.

R. 5. E. Section 18.




15020 This specimen is a poor variety of limestone. It was found in the little field N. of the Dallas road. The field is on rather low ground between Rock creek mountain on the east and the Blankenship mountain on the west. The exact extent of the limestone area could not be determined, but it is evidently quite small, and represents a larger mass which has been carried away by erosion. Rock Creek runs through the section from east to west dividing it into almost equal parts. On the south of the creek near a little branch there is a very small exposure of a fine grained sandstone, the location of which is shown on the map.

11 1/2

T. 33. N.

R. R. 5. E. Section 17




- 15004 This specimen is a coarse sandstone, or a fine conglomerate occurring along a little ravine which leads into Rock creek. It has many fragments of porphyry in it which shows that the porphyry is older than the sandstone. The bedding of the rock is approximately horizontal.
- 15016 This specimen came from a ledge of porphyry by the side of the branch as shown on the map. Macroscopically it shows the flow structure remarkably well.
- 15019 Porphyry, from the bluff on the north bank of Rock creek on Mr. Payne's farm, nearly $\frac{1}{4}$ mi. W. of E. line of Section. It has been studied and described in my Dissertation under No. 287 and is a remarkably good illustration of what I have called the "veicilitic" structure.
- 15060 This rock was taken west and a little south of Mr. Payne's house. It is a remarkably light colored and flinty porphyry.

12 1/2

T. 33. N.

R. 5. E., Section 16.



15002 This section is almost wholly occupied
by "Blue Mt." The hill is a mass of porphyry,
7
15003 ledges of which can be found almost all
around it. In some places it is precipitous,
bluffs 20 ft. or more high, with almost verti-
cal walls extending for considerable distances.
In other places the detrital material has
rolled down the hillside until it is of even
descent, and one can well ride up or down
it. Specimen 2 is from the summit of
the hill; number 3 is from the western
slope about half way down the hillside.

It is on the south-eastern portion of this hill
that the so-called "asbestos mines" are located.


Extending southward from the Blue
Mt., for a number of miles, is a row of por-
phyry hills which are very steep and high.

13 1/2

T. 33, N.

R. 5, E. Section 3.

South half of Section 3.



15005 This specimen of granite is very interesting because it is from the same place from which No. 264 in my dissertation was obtained. This No. 264 was noted on account of its "feld-spar enlargement" crystals. The specimen is from a ravine at a place where there are falls of solid granite about 15 ft. high.

15006 These specimens are interesting because they show gradual transitions from the holocrystalline structure to one strongly approaching the porphyritic. There is a high bluff here about 200 ft. above the water level in the river.


15007 No. 6 is from the side of the hill and is a fairly good granite. No. 7 is from the extreme summit and in its every appearance is closely allied to the porphyries, and in fact must be called a porphyry. There is a vertical wall fully 20 ft. high at the point from which it was taken, and the fissures in the rock are exactly the same as those so common in the porphyries, dividing the rock masses into angular blocks of varying size.

15008 This specimen in appearance is about halfway between granite and porphyry. It was taken from a boulder. The surface at this point is covered with loose rock, some of which are
See P. 59.

14 1/2

T. 33. N.

R. 5. E. Section 15



15090

This number is from the porphyry of the S.E. part of Blue Mt., just above the so called asbestos mine. The porphyry on this side of the hill all shows the flow structure quite well, especially on the weathered surfaces.

15099

This porphyry is from the top of a steep hill called "Bald Mt." The hill stands out alone and is very like the frustum of a cone in shape, the upper diameter being about fifteen rods, and the height 300 or 350 ft. Macroscopically the porphyry seems to be very glassy.


15100

This is a very peculiar dyke rock which came from an old shaft to the S.W. of Bald Mt. a few rods, at a place where prospecting was done a few years ago. The size of the dyke could not be determined as the surface was entirely covered with soil. The exact location is about 100 yds. east of John Inman's house.

15 1/2

T. 33 N.

R. 5. E. Section 22.



15088 Porphyry, in ravine south of J. Mc Dowell's house $\frac{1}{4}$ mi. The whole of the section is covered with porphyry.

15089 This specimen of porphyry was picked up in the field, and was preserved because it showed the flow structure the best of any rock seen up to that time. Of course it had rolled down from ^{the hillside} above.

15091 These two specimens are dyke rocks from the S.E. corner of Blue Mt. near the "asbestos


15092 mines." Each is from a boulder taken out in mining years ago.

The detrital material at this place is quite deep and in a few places, spread out in thin sheets, an inferior grade of asbestos is found. Some little mining for it was done years ago but nothing discovered further than that above described. In a few other localities from 2 to 5 mi. away indications of similar occurrences were noted. It is quite evident that in order for asbestos mining to be successful the original home of the asbestos must be found which probably is some of the numerous dykes, indications of which are so abundant.

16 1/2

T. 33 N.

R. 5 E. Section 21




17

15086 These two specimens of porphyry were taken
from points about $\frac{1}{2}$ mi. apart. The former
15087 is from a ravine which passes the south side
of the south hill of Blue Mt. The latter
from the ^{E.} hillside near the top of the hill.
The whole section is covered by a porphyry
hill with no valley land of note. The
north part is occupied by the southern
extremity of Blue Mt., or rather a little
hill which lies just south of Blue Mt.

17 1/2

T. 33, N.

R. 5 E, Sec. 20.



15017 This specimen is a peculiar dyke rock which occurs in a ravine near the north side of section 17. Quite a number of angular blocks were seen in the ravine, and finally the dyke rock itself was found cutting through the porphyry on the bank of the ravine in such a way that the right contact line was removed so that the width of the dyke could not be determined. The rock is a very fine grained one, and macroscopically seems to be almost amorphous.


15018 This is a specimen of ordinary porphyry taken from the place indicated on the map.

15084 This also is ordinary porphyry taken from a ravine near the south side of the section. The whole of this section is occupied by porphyry hills excepting a narrow strip of sandstone in a ravine in the N. E. corner, as shown on the map.

18 1/2

T. 33. N.

R. 5. E. Section 19




15080. This section is nothing but hills of 19
+ porphyry with ~~an~~ occasional small
15081. dyke. The ravines are lined with bold
ledges of porphyry and the hillsides
are strewn with boulders, some of which
are very large. The location of the
two specimens given can be better seen
on the map than described with words.

15082. This is a dyke rock. The dyke itself could
not be found, but on the S.W. slope of
the hill the dyke boulders were numerous.
This specimen is from one of these.

1912

T. 33. n.

R. 5. & Section 30




15083. This section is simply a mass of porphyry hills. But one specimen was taken. The general appearance of the country and the rocks is very like that of all the sections near it to the north and east. Occasionally a ravine leads from the hillside into the valley either to the east or to the west. Along these ravines are numerous ledges of porphyry exposed. Also there are numerous "glades," or places on comparatively level ground over which the rock is exposed.

20 1/2

T. 33. N.

R. 46, Section 28



15094-95

15093

15093

This is a porphyry from the E. bank of the S. Rock creek, very near the S.E. corner of Section 29, and on the S.W. corner of Proctor Mt. 21

15094

This is also a porphyry which has nothing of a special interest about it so far as could be seen either in its field relations or in the character of the rock itself. Its location was near the half mile corner on the south side of the section.


15095

This is a dyke rock which came from a dyke fully 50 ft. wide and which had a N.E. and S.W. trend. Both the dyke rock and adjacent porphyry were badly fractured. The fissures in which were generally filled with quartz. The rock was a very fine grained one.

21 1/2

T. 33. n.

R. 5. 8, Section 28




15085 This specimen is porphyry from the W. side of the S part of Elide Mt. The section is occupied by this hill. No important difference was noted either in the character of the rock or in the mode of occurrence here, and a number of other porphyry hills already described.

22 1/2

T. 33. N.

R. 5. E. Section 23,



23


15106 These specimens of porphyry came from
the face of the hill and its spurs. This hill is
one of medium height, is about twice as
long N. and S. as it is broad, and is
quite steep nearly all the way around.
On the east side a spur extends out
so far that it can hardly be called a
part of the same hill. A ravine rises on
the E. side and flows N.E. to the river.
Along this ravine the banks gradually
become steeper until the river bottom
is reached.

Specimen 106 came from the S.E.
corner of the hill about half way up
the side. No. 107 is from the bed of
the ravine above mentioned.

23 1/2

T. 33. N.

R. 5. E. Section 24.



15015 This specimen of porphyry came from ^{24.}
a steep and high bluff on the E. bank
of the Little St. Francois river near the
S. E. corner of the section. The stream
runs nearly north. The bluff is from 100
to 125 ft. high and is composed of porphyry.
There are many vertical fissures which
can be traced from top to bottom of
the bluff. They are all very small and
are of the same nature as those found
every where so abundantly in the
porphyry. On top of the bluff two
high finger like projections of rock are
to be seen. One of these is fully 25 ft.
high and not more than 6 ft or 8 ft.
in diameter at the base. The opposite
bank of the creek is very low and af-
fords a good view of the bluff.


15257 This is a specimen of porphyry from
the N. W. part of the section, and
from the N. W. side of a tolerably
high hill.

The section was not examined in
sufficient detail to warrant me in
coloring the map.

24 1/2

T. 33, N,

R. 5, E, Section 14,



15242 This specimen came from immediately ²⁵ on the contact line, between the granite and porphyry. It was taken from a boulder, but as it showed in itself a transition from the coarsely crystalline to that which is much finer it is a valuable specimen. It probably had not been far removed from its original home. The granitic part is lighter in color than the other.


15243 This is a porphyry from the dump pile at the Martin shaft, and probably came from about 75 ft. under the surface, altho' there is a little uncertainty about it. A shaft was sunk with an angle of about 45° with the vertical on what was supposed to be the same vein which was worked at the Silver Mines.

This section was not examined in sufficient detail to warrant me in coloring the map, altho' I almost know that the whole of it is porphyry South of the contact line marked.

26 1/2

T. 33. N.

R. 5, E. Section 13



15241 This is a very coarse sandstone or a conglomerate. A comparatively small area, with the surface exposure as shown on the map is covered with this kind of a rock. The bedding slopes gently with the surface. The rock itself is almost entirely composed of small boulders and coarse gravel of porphyry cemented largely with iron-oxide. 26

15244 This is a porphyry thrown out in digging a cellar immediately on the contact line between the granite and porphyry. At the south end of the cellar the rock is porphyry; at north end it is granite. It is caved in and partially filled with water, so that the contact could not be seen.


15249 This is granite. The surface for all the granite in this section is level, and is mostly covered with soil.

15250-52 These were taken from the contact line. A rock was found which seemed to show the transition. I blasted out a portion of it and obtained 250, which is granite, 252, which is porphyry, and 251, which is intermediate. 52, 59.

26 1/2

T. 33. N.

R. 5. E. Section 12.



15245- This is a specimen of quartz from the ²⁷
"Apex" shaft, and was taken because it
so well represents the corrosive action
of water upon quartz under certain
conditions.

15246 Quartz-diorite, from a dyke at "Silver
mines." It was in small boulders, the
exact location of the dyke not being
known. It is very similar to No's. 15001
and 15045.

15247 Dyke rock from a big dyke just
N. of a path leading down to the river
at "Silver mines."


This place called the "Silver mines"
is very interesting. The country rock is
granite. Quite a number of dykes cross
it at different places bearing about
N. 75° E. In one or more of these
places it seems the fissure was not
filled with lava, but was left to
be filled subsequently with vein
stocks. We find evidence of se-
merole action. There are fluorite,
lepidolite, topaz and wolframite, which
indicate it.

See p. 60.

27 1/2

T. 33 N.

R. 5 E, Section 11,




15248- Granite from the bed of a ravine called "Mud creek." This whole section is granite excepting a portion in the S.W. corner, here the contact line bears to the N., but the exact boundary could not be determined on account of the soil. The rocks are exposed in different places to a sufficient extent to admit of a quite positive determination of their character.

15258- This is a dyke rock taken from a boulder in a branch of the roadside $\frac{1}{4}$ mi. E. of Mr. May's. It represents a number of large boulders which seem to be piled up winnow like trending N.E. and S.W. It is very probable that they are representatives of a large dyke concealed from view by the soil covering. Exactly the same kind of boulders occur in a ravine in Mr. Lioander Acklock's field just W. of his house in Section 10. They are peculiar in having a very rough surface where they have been weathered.

28 1/2

T. 33, N.

R. 5, E. Section 26,



15007-12 The principal interest in this section is 29
in connection with the so-called "Tin Mines".
These are located on the N. side of "Tin Mt." a
steep porphyry hill about 400 ft. high. There is
a very large dyke exposed on the northern side
of the hill and which was the cause of a big
excitement about 20 yrs. ago. The dyke is near
50 yds. wide and is filled with a very black,
compact glassy rock, which is much more
perfectly crystalline at the base than near the
top. No. 9 is from a boulder near the top of
the hill at a point the highest up of which
any dyke boulder was seen.

No. 10 from the tunnel highest up on the
hill, which the miners call tunnel No. 4

No. 11 from tunnel No. 3.

No. 12 " " " 2.

No. 13 " " " 1.


These tunnels were driven horizontally into the
hill 100 ft. or more, and the specimens were taken
from the dump piles at the several locations.
The vertical distance between tunnels No. 1 and 4
is about 175 ft. Rocks from this dyke were
described quite fully in my Dissertation.

15014 A porphyry rock from the summit of "Tin Mt."

29 1/2

T. 33. N.

R. 58. Section 10,




15001. This specimen is very interesting because ³⁰ it is a quartz bearing diabase-porphyrite dyke rock, and is the one described so fully in my Dissertation. At the time I gathered the specimen I was unable to locate the dyke but found a place at which the boulders extended at least some distance under the soil and probably were immediately over the dyke itself. Only a few feet away the granite is exposed, and during a subsequent visit to the place I found a dyke 50 in. wide, cutting through the granite and trending N.E. and S.W. There can be no doubt but that there are at least two dykes in this immediate vicinity. Porphyritic quartz crystals and feldspar crystals are very common, scattered all through the rock mass. Some of the feldspar crystals are nearly an inch in diameter.

15062 This is another dyke rock from a small ravine running N.E. It seems that the ravine is immediately over the dyke itself, which is a big one being at least 20 ft. wide. To the S.E. about 20 yds., across the section line, there is another ravine which also seems to follow a dyke. The two rocks seem to be identical and therefore a specimen of the latter was not taken.

36 1/2

T. 33, N.

R. 4. E., Sections 1, & 2,



27

These two sections, are nearly all composed of porphyry hills. On the extreme N.E. along a small ravine, there are very narrow belts of limestone and sandstone and also the western extremity of the sandstone area described under Section 6. R. 5. The hills are very high and abrupt. The porphyry is exposed in nearly all places.


15030 These specimens are coarse grained porphyry, & the former of which came from Mr. Bartch's
15031 "gold mine," that is from the place where Mr. Bartch had dug a hole about 4 ft. deep in the solid porphyry. This is a good place to note the different degrees of crystallization in the porphyry. Along the walls of this one little hole several different degrees of crystallization can be seen, and they present such an appearance that it is positive no intrusive rocks are present.

15032 These are sandstone from locations as shown & on the map. 35 is a fine grained rock and
15035 the layers were quite thin. 32 is more coarse grained. At this place there is a large boulder of
15034 porphyry imbedded in the sandstone from which specimen No. 34 was taken. The boulder is fully 3 ft. in diameter, and does not seem to be
See R. 61

37 1/2

T. 33, N.

R. 4, E. Sect. 11, 12, 13, & 14



15068. These four sections are ^{almost} entirely covered with ^{38.} porphyry hills. The southern portion of 13 and
 4 15071 14 slope off towards the large valley called
 4 the upper ^{the} "Flat woods". The whole mass of hills
 15130 are known by the name of the "Grassy hill".
 4 There is nothing of special interest about any of
 15136 these specimens and macroscopically could not
 4 well be distinguished from many others.
 15137

15180 These four numbers 15180-83 are splendid
 15181 illustrations of the transition stages of
 15182 crystallization. The first came from the
 15183 foot of the hill S. of Mrs. Henson's field.
 The distance along the foot of this spur at
 which such rocks are exposed is only a
 few hundred feet. '81 came from 150 yds.
 up the hill S. from '80. '82 came
 from 150 yds. to the S.E. from '81.
 '83, which is a good porphyry, came
 from 34 the way far to the summit
 of the big hill to the S.E. from '82.
 There seems to be a perfect gradation
 from the coarsest crystalline one
 to the one which is the most porphyritic.

38 1/2

T. 33, N.

R. 4, E. Section 3,



39

15750 This specimen of porphyry is from the little hill just across the Township line. The hill is small and almost circular and is entirely surrounded by limestone.

15757-8. We have here a very interesting change from the ordinary porphyry to a crystalline rock which seems more like granite than porphyry. These specimens are from the head of a ravine N. of the section line. The first is the crystalline, the second the porphyritic variety. They are not more than 15 ft. apart. The ravine seems to be the dividing line between the two different kinds of rock farther down.

15752a These specimens are the crystalline variety
+ mentioned above and are located as
15754 shown on map.


15753 This is a porphyry and lies between 752a and 754. It was noticed that as we ascend the hill near 754 the change from the crystalline to the non-crystalline condition occurs, so that at the top of the hill we have the porphyry alone. Following down the creek we find the same coarse rock

39 1/2

T. 33. N.

R. 4, E, Section 3,

South half of Section 3.



reaching almost all the way. A change occurs here which is easily seen and traced. At the creek almost immediately opposite the highest point in the road as it runs over the foot hill at the "Shut in" is the line between these two kinds of rock.


At one place a ledge was found with a vertical face of 6 ft. or more, and in it the change occurs. At the middle of the bottom ledge specimen No. 162 was taken. It is wholly crystalline as can be seen quite plainly with the naked eye. Either to the right or above the change occurs. Specimen No. 161 was taken 5 ft. immediately above 162 with absolutely no indication of a contact line between the two. On the right there is a fissure which might be taken for a contact line, but as there is none above it becomes probable that the one on the right is only a common fissure.

15151 This is a limestone with many small fragments of porphyry in it. The limestone is in the bed of the creek and extends up the hillside a short distance. The sandstone overlies the limestone and extends
see P. 62.

40 1/2

T. 33, N.

R. 4. E. Section 4.



41
#7

This section is nearly all limestone. In the extreme N.E. corner there is a circular porphyry hill about $\frac{1}{4}$ mi. in diameter. Near the N.W. corner there is another elevation, but it is a limestone hill. Judge Russell's house stands here. He has quite a stone quarry opened. The rock is of the same general character as the limestone throughout the valley. In some places there is so much calcine in the rock that it is practically useless for building purposes. Along the west line of the section, near the middle, is another steep circular hill covered with flint rock but which has limestone at its base.


15759. This specimen of porphyry came from the porphyry hill in the N.E. corner of the section.

41 1/2

T. 33, N.

R. 4. E. Section 5.

North half of section 5



15146 These are specimens of porphyry breccia
 & obtained from the bank of the creek

15149 at the iron wagon bridge between
 Fronton and Arcadia. It seems to be
 fragments of compact porphyry ce-
 mented together with lava which
 itself produced a porphyry upon

15147 solidification. No. 147 came from
 the summit of "Church hill" only a
 short distance away and is a very
 compact rock of conchoidal fracture
 and also shows the flow structure
 very plainly. The probable explanation
 is that as the lava flowed down the
 hillside the first crust that was
 formed was broken into fragments by
 a second flow of lava thus pro-
 ducing the porphyry breccia.

15148. This is a limestone taken from the
 creek bank just north of Mr. Dixon's
 living barn. 1


The whole of this valley is covered
 with limestone excepting "Church
 hill" and its spurs.

42 1/2

T. 33, N.

R. 4. E. Section 5,

South half of Section 5.



#3
43


15-167. Flint rock. In quite a number places in this township larger or smaller flint hills occur. They invariably have limestone at their bases, and in some undetermined way have an intimate connection with the limestone. The maps illustrate the relative positions of the limestone, flint and porphyry much better than can be given by short verbal description.

15-171. This is porphyry from the N.E. side of Lewis Mt. just S. of Arcadia. All the top land of this map is covered with limestone.

43 1/2

T. 33, N.

R. 4, E. Section 6,




15163. This is porphyry from the S.W. spur^{2/4}
of Shepherd Mt. It is nothing more than
ordinary porphyry.

15164. Limestone - Lewis Mt. does not extend
quite to the W. side of this T. but a
neck of limestone reaches S. from
Shoals creek to the larger body
around Logan.

44 1/2

T. 33 N.

R. 4 E. Sect. 7, 8, 9, & 10,



15766- This is flint rock taken from a ⁴⁵big
boulder 4 ft. in diameter taken from
the N. hillside N. of the switch of
the railroad near "Tip-Top."

15765- Limestone from the R. R. cut, just
N. of switch near "Tip-Top." There is
limestone at the greatest elevation
yet seen. The R. R. cut shows quite
heavy beds of well stratified
limestone, some layers of which are
nearly 2 ft. in diameter, and which
dips to the N. quite gently, same as
slope of the hill.

15768- Porphyry on N. side of hill as shown
on map.

15769- Dyke rock on S. side of same hill, taken
from a boulder. The dyke itself was not
seen.

15773- Limestone just N. of Mr. Guffy's house.
It is very near the line between lime-
stone and porphyry.

15774- Porphyry, from a point just E. of Mr.
See P. 62.

45 1/2

T. 33. N.

R. 4. & Sect. 15, 16, 17
20, 21, & 22,



This map represents 6 sections.

15178

&

All these are porphyries from locations as shown on the map and

15179

&

have no special properties which would distinguish them from others.

15218

15219

A very compact limestone from the base of a hill which is covered with flint.

15220

These are porphyries from the same hill.

&

15175

One from the north and the other from the southern extremity. The hill is of moderate height and is entirely surrounded by limestone.

15212

Flintrock, which seems to be a sort of flint conglomerate. It is on the top of the hill and represents a kind often seen in this locality.

15213

Limonite. These flint hills sometime have considerable deposits of limonite which seems to be of a fairly good grade. Years ago prospecting was done in a number of different places and the ore is still lying in heaps on the surface.

46 1/2

T. 33. N.

R. 4. E., Section 18,

S. L. T. M. & S. R. R.

15172

Greenville Road.

15184

15203, 15202.

15172

+

15184

47

These are porphyries from the N. and S. ends of the long porphyry hill marked on the map. On the N. the rocks have the flow structure well marked, but it could not be so well seen on the S.

15202

+


15203

These are from an old shaft on the N.W. part of the "Cuthbertson Mts." The "Manganese Mts." of former days. The first shows the flow structure very well developed and was taken on that account. The second shows the manganese ore in little stalactitic formations which is very good evidence that it was deposited by water.

47 1/2

T. 33, N.

R. 4, E. Section 18,



48

This section is a most interesting place. It is the location of the old "manganese mines" on the Leuthbertson Mt. described by Pompey, in his Rep. of Missouri Geol. Surv., 1872. It is the place where limestone blends into porphyry according to his Rep. P. 26. Leuthbertson Mt. is irregular in shape sending a sort of spur to the N.W. and to the S.W. It bears positive evidence of being an old lava flow. There are large masses of rock all twisted and contorted just like lava frequently is.


The manganese ore is often in a fine black powder, sometimes dry but often wet and "mucky." It is only occasionally in solid lumps of any much size. At other times it is in small masses or in a powder scattered through the numerous small openings in the lava. No other place in the whole area visited is half so changeable as this. There are many different looking rocks which at first sight seem to be entirely distinct, but which, I think, are closely related genetically. Thus, there are many places

See P. 63

48 1/2

T. 33, N.

R. 4, E, Section 24,



Section 23 is almost entirely in ⁴⁹ a level valley called the "Bepper Flat Woods". It is entirely under cultivation excepting the western part which is occupied by flint hills. No specimen was taken from it and it is not represented in this Note-Book by a map.

- 15069 This is a porphyry taken from the creek bank just N. of Mr. Blayton's field, N. of the "Flat Wood" school house.


All of this section is level ground and it is only here and there that the rocks are exposed. This specimen seems to be ordinary porphyry.

- 15070 This is lime stone from a well 20 ft. deep in the yard of Mr. Roebbel. No surface rocks are exposed in that vicinity.

- 15072 This is porphyry from a ledge by the roadside S. of the "Flat Wood" school house. There is an area here of only a few square rods over which the porphyry is exposed.

49 1/2

T. 33. N.

R. 4. E. Sections 25, 26,
35 & 36,

This map represents four sections.
 The interesting feature of this area is
 the alternate appearance of porphyry,
 lime stone and flint.

15076 This is a porphyry from comparatively low ground.
 The hill rises to the S.

15237 Is, also a porphyry from the high part of the
 hill. It is interesting to note how the flint
 and porphyry meet here on the hill without
 there being any apparent break in the config-
 uration. To the N. and S. valleys soon set in
 which divide the porphyry hills from the flint
 hills.

15238 This is also porphyry.

15239 Limestone, taken from a point fully 100 ft.
 above the valley, making it the highest
 limestone seen in this corner of the town ship.


15234 Porphyry, taken from hillsides as
 15240 represented on the map.

15233 Limestone from the creek bed. It has a
 great many large like porphyry gravels in it.
 15236 Are limestone, taken from well strati-
 15273 fied layers which have a gentle slope in
 See P. 64

50 1/2

T. 33. N.

R. 14. E. Sect. 27, 28, 33, & 34,



4, This map represents 4 sections. It is a ⁵¹ map of porphyry, flint and limestone alternating.

- 15295 There is nothing of special interest in the hand specimens or in their mode of occurrence. The limestone is in the valley and is always well stratified and approximately horizontal. But generally has a gentle slope in the direction of the hillsides. The map shows the location of the several specimens so that it is unnecessary to mention them in detail.


To the S.W. of this map, across the township line, two other specimens were taken.

- 15210 The former is a sandstone which came from layers along a branch.
15211 The latter is a dyke rock obtained from boulders on the surface, the dyke itself not having been found.

57 1/2

T.

R.



This map represents 4 sections 52
and is principally occupied by the
"Biser Mt." This hill is large and high
and very irregular in shape.

15205-

On the extreme N. of section 30 is a flint
hill which reaches across the section
line. The base of this hill is occupied
by a very compact limestone, called
marble by the citizens of the country.
Passing immediately S. from this we
cross a second hill which is composed
entirely of porphyry. On the northern
slope of this prospecting was done years
ago in a vain search for iron ore. The
rocks thrown out are of a dark red to
black color, waxy lustre, and some of
them seem to be a porphyry breccia.
This rock will probably prove of great
interest. Passing still farther S. the
S. we ascend the northern spur of
"Biser Mt." The porphyries of this hill
have porphyritic feldspars which
show much more plainly than the
average. The northern part of sec-
tion 32 and southern part of sec-
tion 29 are occupied by a flint
hill from which specimen 281 came.
See page 65.

15028 These are from the contact line,
the former granite, the latter
15029 porphyry, and are but 6 in.
apart.

6

Continued from P. 1.

The exposure consists of a narrow strip extending along a little branch, and covers but a few square rods. To the N.E. in a number of places small boulders and gravel of porphyry are scattered over the surface. I was first puzzled because I could not account for them, but it now seems probable they are the remnants of this same conglomerate rock, which may have at one time extended over those areas and which is now entirely disintegrated and largely removed. The rock when now exposed has a dip of fully 25° to the S.E. by S. E. giving it a strike N.W. by N.

Continued from P. 2.

15042. 42 is from the highest point of this high bluff, which is called "Bumpers Hill." 42 is
15043 from the bed of the creek immediately below 43. On the north side of the creek
15046 some distance above the contact line there is also a moderate hill but without steep bluffs.
15047 At the foot of this hill the rock is so coarsely crystalline that it will almost do to call it
15048 granite. 48 was taken from this point. Near the summit of the hill the rock is not so well crystallized. 46 & 47 are from near the top of the hill.

On my last visit to the place I followed up the creek and gathered specimens from the creek bed all the way along, as is shown on the map. Some of these are quite coarse but in general the coarseness of texture decreases as the distance from the contact line increases so that a mile or so away crystallization is little if any more nearly perfect.

than in the ordinary porphyries in other places. The individual specimens will show for themselves the way this change occurs much better than can be described in a field Note-Book.

15045 Quartz-diorite - porphyrite. This rock was found in boulders on the north ridge of the hill, as shown on the map. The dyke itself could not be found, on account of the heavy covering of soil on that part of the hill.

15044 Sand stone. The northern part of this section is covered with sand stone. The portion shown on the map is the southern extremity of a large circular area nearly 3 mi. in diameter, almost all of which is probably covered with sand stone. Only a narrow strip on the south side of this valley lies within T. 33. N. and consequently a detailed examination of the remainder was not made. Nearly all of the valley is covered with soil, and is extensively cultivated. Along the branches and occasionally at other places outcroppings can be seen. Along the little branch in the extreme N.W. part of the map very heavy beds of sand stone are laid bare. At every place noticed the bedding seemed to be horizontal, and thus far there appears to be a total absence of any indications of orographic movements of any extent since the formation of the sandstone. 44 is from the surface at a little ravine; and 49 is from Esq. Bowman's well but I could not determine

from what depth it came. The well was sunk about 25 ft. deep and did not pass through the sandstone.

Continued from P.E.

- 15110. W. side of N.E. corner 2 ft. 8 in.
- 15111. E. of and adjoining 109.
- 15112. N.E. corner 3 ft. below surface.
- 15113. N.E. corner 4 ft. 6 in.
- 15114. N.E. " 5 ft. 6 in.
- 15115. W. side of N.E. corner 7 ft. 6 in.
- 15116. S. " " " " 8 ".
- 15117. " " " " " 11 ".
- 15118. W. " " " " 14 ".
- 15119. S. " " N.W. " 15 ".
- 15120. Middle of W. side 14 ft.
- 15121. " " " " 15 ".
- 15122. N.W. corner 18 ft.
- 15123. S. side N.E. corner 17 ft.
- 15124. W. " S.E. " 17 1/2 "
- 15125. Porphyry, surface rock S.W. corner of shaft.
- 15126. Coarsest rock seen on dump pile.
- 15127. Granite from surface at contact line
30 yds N. of the shaft.
- 15128. Porphyry, surface rock 5 ft. S.W. of 127.

porphyry, some granite, and others half way between the two. The ledges of the rock are where exposed are granite. It could not be determined exactly from whence the boulders came, but it is evident they have not been moved far and it is quite probable they now exist on the same ground where they were formed.

15064. This is a coarse dyke rock from the spring near the house of Peter Weiss Jr. There is a large dyke here exposed over an area of 150 ft. by 75 ft. or more. The surface covering made it impossible to determine the direction of the trend of the dyke.

15065. This is a granite specimen taken from the hillside near the river, as shown on the map. The bluff is about 150 ft. high with granite ledges very abundant.

15061. This is a dyke rock at the foot of the high hill mentioned under no's. 6 & 7. The rock varies in color from dark to light. The latter being composed almost wholly of feldspar.

15068. This is a dyke rock found by the road side leading from Peter Weiss's to young Peter Weiss's. The dyke itself could not be located, but there was a very large amount of boulders scattered over the hillside.

Continued from P. 26.

The two extremes were not more than 20 in. apart.

15253. A coarse grained porphyry from near the

contact line, as shown on map.

15254. This came from an old shaft and was called slate by the miners, but it looks very like hornblende which it probably is. This mineral is so scarce in this part of the country that it makes it quite interesting should it prove to be hornblende. The wall rock of the shaft is porphyry.

15255. This is a half way granite, which of course came from near the contact line.

15256. This is porphyry from just across the contact line.

In tracing this contact line great care was taken. The country is sometimes level and covered with a thin coating of soil, at other times hilly with abundant exposures of rock. It is very noticeable that the line is not sharply drawn. Often a coarsely crystallized area will be rather to the S.W. than some of the porphyry. Just N. of No. 250, for example, the porphyry extends 20 to 30 ft. Such examples are often found.

Continued from P. 27.

The mining was principally done on the largest of these veins which carries argentiferous galena. The vein is not vertical but dips to the S.E. about 45° , according to Mr. Wiley who at one time was Superintendent of the mines.

57. Dike rock from E side of river from Silver Mines.
 The same as No. 290 in my classification, which see.
 58. Dike-like from Silver Mine.
 59. The so-called "serpentine" from Silver Mine.

Continued from P. 37.

water worn, and therefore must have rolled down the hillside while the sandstone was forming. This is positive evidence that the porphyry is older than the sandstone.

15133- This is limestone from along the same ravine and is so interbedded with the sandstone that there cannot be any material difference in the ages of the two rocks. Its location is shown on the map.

15131- This is porphyry from the summit of the hill above the Barker ravine, as shown on the map.

15132- These are all porphyries, the last two of which are from the bank of Shook's creek. The other is from a

15144- high point overlooking the creek.

15145-

15133 & 134- These are sandstones from the bank of the branch at Mrs. Benson's spring. The bluff here is 15 or more feet high. Well stratified sandstone overlies limestone beds and is conformable with them.

15135- This is limestone from immediately under the two sandstones just described.

The whole of these stratified rock dips gently to the N. W., and are good illustrations of what is so often seen in this township, namely, a stratified rock fringing the bases of hills and bearing gentle inclinations in the directions of the hill-sides themselves.

It may here be said once for all that in no case thus far observed is there any evidence of the bedding of the stratified rock ever having been materially changed. Usually these rocks are gently

inclined, but almost invariably in the direction of the slope of the hillside. This is most likely due to the rocks having been deposited on a sloping bottom. We look in vain for any evidence of material orographic movement since the stratified rocks were deposited.

Continued from P. 36. [p. 40]

up the Dallas road perhaps all around the hill.

15152. Porphyry from same place as 151.

15176. This is limestone from the N.W. corner of the circular hill as shown on map.

15177. Porphyry from the summit of the hill.

This circular hill of porphyry is about 250 to 300 ft high and presents a unique appearance rising as it does from a limestone valley. It is almost circular and a half mile diameter at base. There is no indication of its having been lifted up through the limestone and probably was formed long before the limestone was deposited.

Continued from P. 45.

Guff's house. It was taken from only a few feet E of the limestone limit.

15170. Limestone from N. hillside in bank of ravine. The hill is apparently composed of flint, but the limestone at its base is almost horizontal and seems to pass in under the hill.


15/55 Dyke rock. This is a most interesting occurrence of a dyke rock. It is surrounded by limestone, and on account of its resisting erosion more than the limestone has produced a narrow ridge throughout its entire length, a little more than 18 mi. The ground in most places is so covered with soil that I could not determine whether the dyke itself passed through the limestone, or whether it was formed before the deposition of the latter. It seems probable that the dyke is quite wide, 50 ft. or more, but this can not be stated definitely.

Continued from P. 48.

which have surface rocks that seem to be stratified. These overlie the massive rocks of the hill, and are at places apparently different from them, but we always find that on the hillside the pseudo-stratified rocks extend downward to about the same depth below the surface, no matter at what point on the hill the examination is made. At one place a trench about 2 rods long, has been dug into the hillside and this shows that any given layer merges into the massive rock of the hill, thus

A shaft sunk would show the layers down to a certain depth with the massive rocks below, but the cut on the hillside looks quite different.

Again, in the ravine N. of the hill at one



place the massive rocks by weathering are actually changing into shells and lusters.

All these points make it seem probable that the apparent stratification is due, (a) to lava flows, and (b) to weathering.

This is an important question and is explained entirely differently by Pampelly in his report above mentioned.

The manganese ores were probably deposited by water and fills nearly every little opening in the rocks.

15785. This seems to be a dyke rock, but it may not be, as some of the country rock looks very much like it.

15786 These three are from the same place, at one of the old mines and were taken to show

15787 that the rock mass certainly is a lava.

16788

15789. This is a specimen of the manganese ore,

All the remainder on this map are taken to show the similarity of the rock in the different parts of the hill. The map shows their location much better than can be given in words.

Continued from P. 50.

the direction of the hillside.


15074. Is a sandstone taken from a boulder found on the top of a hill. Search was made in vain for this rock in situ.

15075. This is flint taken from near 74. The hill from which the two came is flint throughout with limestone at its base.

Continued from P. 52.

All of the specimens not named from these sections are simply porphyry and need not be described in detail.

To the S.W. of section 31, across the township line, is a little knoll of very coarse granite porphyry. Some of the porphyritic feldspar crystals are nearly 2 in. long. This little knoll, which is not exceeding 300 yds. in diameter is almost entirely surrounded by ravines. To the N.W. a narrow neck reaches across on to the porphyry hills. This locality furnished an excellent opportunity for studying the crystallization question. Specimen 24 is the granite porphyry from the extreme S.E. corner of the exposure.



Specimen 25 is porphyry from 100 yds. up the ravine to the west of the northern exposure of the granite porphyry. No. 26 is up a little ravine 50 yds. to the north. No. 27 is the granite porphyry on the hillside about 50 ft. steep height from a point in the ravine at which No. 25 was taken. This latter is porphyry. No. 29 came from the same ravine about 50 ft. below 28.

So far as could be seen in the field there is a gradual transition from the coarse granite porphyry through No. 27 to which itself is comparatively fine grained, to the porphyries represented by the other numbers.

No.	P.	No.	P.	No.	P.	No.	P.
15001	30	15039	2	15077		15115	8
2	13	40	2	78		116	8
3	13	41	2	79		117	8
4	12	42	2	80	18	118	8
5	14	43	2	81	18	119	8
6	14	44	2	82	18	120	8
7	14	45	2	83	20	121	8
8	14	46	2	84	18	122	8
9	29	47	2	85	20	123	8
10	29	48	2	86	17	124	8
11	29	49	2	87	17	125	8
12	29	50	4	88	16	126	8
13	29	51	4	89	16	127	8
14	29	52	6	90	15	128	8
15	24	53	3	91	16	129	1
16	12	54	3	92	16	130	38
17	18	55	3	93	21	131	37
18	18	56	3	94	21	132	37
19	12	57	27	95	21	133	37
20	11	58	27	96		134	37
21	10	59	27	97		135	37
22	9	60	12	98		136	38
23	1	61	14	99	15	137	38
24	1	62	30	100	15	138	2
25	1	63	14	101		139	2
26	1	64	14	102		140	2
27	1	65	14	103		141	2
28	1	66	7	104		142	2
29	1	67	7	105	16	143	2
30	37	68	38	106	23	144	37
31	37	69	49	107	23	145	37
32	37	70	49	108	8	146	42
33	37	71	38	109	8	147	42
34	37	72	49	110	8	148	42
35	37	73	50	111	8	149	42
36	1	74	50	112	8	150	39
37	4	75	50	113	8	151	40
38	2	76	50	114	8	152	40

No.	P.	No.	P.	No.	P.
74	15152a39				
72	15153 39	15191 48	15229 52		
	154 39	192 48	230 52		
	155 45	193 48	231 52		
	156	194 48	232 52		
	157 39	195 48	233 50		
	158 39	196 48	234 50		
	159 41	197 48	235 51		
	160 37	198 48	236 50		
	161 40	199 48	237 50		
	162 40	200 48	238 50		
	163 44	201 48	239 50		
	164 44	202 47	240 50		
	165 45	203 47	241 26		
	166 45	204 48	242 25		
	167 43	205 52	243 25		
	168 45	206 52	244 26		
	169 45	207 52	245 27		
	170 45	208 52	246 27		
	171 43	209 52	247 27		
	172 47	210 51	248 26		
	173 45	211 51	249 26		
	174 45	212 46	250 26		
	175 46	213 46	251 26		
	176 40	214 52	252 26		
	177 40	215 51	253 26		
	178 46	216 51	254 26		
	179 46	217 51	255 26		
	180 37	218 46	256 26		
	181 37	219 46	257 24		
	182 38	220 46	258 28		
	183 38	221 48	259		
	184 47	222 52	260		
	185 48	223 52			
	186 48	224 52			
	187 48	225 52			
	188 48	226 52			
	189 48	227 52			
	190 48	228 52			

