

Anaquest advertisement.

[s.l.]: [s.n.], 1984

<https://digital.library.wisc.edu/1711.dl/TPAJRJS5BCQWT8D>

<http://rightsstatements.org/vocab/InC/1.0/>

The libraries provide public access to a wide range of material, including online exhibits, digitized collections, archival finding aids, our catalog, online articles, and a growing range of materials in many media.

When possible, we provide rights information in catalog records, finding aids, and other metadata that accompanies collections or items. However, it is always the user's obligation to evaluate copyright and rights issues in light of their own use.



AErrane™ (isoflurane)

Alyrane™ (enflurane)

Products of Original Anaquest Research

For Safety

Low biodegradation, absence of toxic metabolites

For Rapid, Precise Control

Low blood solubility, rapid induction, smooth recovery

For Cardiovascular Stability

No increase in arrhythmias due to adrenaline
used for haemostasis

For Simplified Management

Inherent muscle relaxant effect sufficient for most procedures,
including abdominal and thoracic surgeries

AErrane™ (isoflurane)

Prescribing Information

Indication: for general anaesthesia.

Dose: Inspired concentrations of 1.5% to 3.0% produce surgical anaesthesia in 7 to 10 minutes. Surgical anaesthesia can be maintained with 1.0% to 2.5% AErrane when used concomitantly with nitrous oxide.

Contraindications:

Known sensitivity to AErrane or other halogenated anaesthetics, or a known or suspected genetic susceptibility to malignant hyperthermia.

Side Effects:

Hypotension, respiratory depression, arrhythmias, elevated white blood count, shivering, nausea, vomiting and ileus have been reported.

Price: 100mL £33.50

Product Licence

Number: 5247/0002

Alyrane™ (enflurane)

Prescribing Information

Indications: For general anaesthesia.

Dose: Inspired concentrations of 2.0% to 4.5% Alyrane produce surgical anaesthesia in 7 to 10 minutes. Surgical anaesthesia can be maintained with 0.5% to 3.0% Alyrane. Analgesia equivalent to 30% to 60% nitrous oxide for vaginal delivery can be provided by 0.25% to 1.0% Alyrane. For caesarean section, Alyrane should be administered at concentrations of 0.5% to 1.0% as a supplement to other general anaesthetics.

Contraindications:

Seizure disorders; known sensitivity to Alyrane or other halogenated anaesthetics; known or suspected genetic susceptibility to malignant hyperthermia; use of product in conjunction with isoniazid or other hydrazine-containing compounds.

Side Effects:

Hypotension, respiratory depression, arrhythmias, myocardial depression, elevated white blood count, shivering, nausea and vomiting have been reported.

Price: 250 mL £33.30.

Product Licence

Number: 5247/0001.