

Ciproxin advertisement.

[s.l.]: [s.n.], 1972

<https://digital.library.wisc.edu/1711.dl/I6GOLZEGSQSBN80>

<http://rightsstatements.org/vocab/InC/1.0/>


The libraries provide public access to a wide range of material, including online exhibits, digitized collections, archival finding aids, our catalog, online articles, and a growing range of materials in many media.

When possible, we provide rights information in catalog records, finding aids, and other metadata that accompanies collections or items. However, it is always the user's obligation to evaluate copyright and rights issues in light of their own use.



Ciproxin[®]

ciprofloxacin



Parenteral power in a tablet

▽ CIPROXIN INFUSION/TABLETS ABRIDGED PRESCRIBING INFORMATION (Please refer to full data sheet before prescribing.) **Presentation** Clear, almost colourless solution containing 200mg ciprofloxacin in 100ml solution. White film-coated tablets, marked with the Bayer cross on one side, each containing 250mg ciprofloxacin. **Uses** Ciprofloxacin is indicated for the treatment of single or mixed infections caused by susceptible organisms. **Dosage and Administration** Adults — *Ciproxin Infusion* 100-200mg administered intravenously over 30 to 60 minutes twice daily. *Ciproxin Tablets* should be swallowed whole with liquid; the dosage range for adults is 250-750mg twice daily. For acute infections the usual treatment period is 5 to 7 days (intravenous) or 5 to 10 days (oral). Initial intravenous administration may be followed by treatment with oral ciprofloxacin. **Contra-indications** Hypersensitivity to ciprofloxacin or other quinolones; also in children and growing adolescents except where the benefits of treatment exceed the risks. **Warnings and Precautions** Ciprofloxacin should be used with caution in patients with a history of convulsive disorders. Increased plasma levels of theophylline have been observed following concurrent administration with ciprofloxacin. The dose of theophylline should be reduced and plasma levels of theophylline monitored. Where monitoring of plasma levels is not possible, avoid the use of ciprofloxacin in patients receiving theophylline. The tablets should not be administered within 4 hours of antacids containing magnesium/aluminium hydroxide. **Use in Pregnancy and Lactation** Administration during pregnancy or to nursing mothers is not recommended. **Side-effects** Local irritation at the site of injection (infusion only), gastro-intestinal disturbances, skin rashes, disturbances of the central nervous system. **Pharmaceutical Precautions** Unless compatibility is proven, the infusion should always be administered separately. **Legal Category** POM. **Package Quantities** Ciproxin Infusion, bottles of 50 or 100ml. Ciproxin Tablets, foil strips of 10 in packs of 10, 20, and 100. **Product Licence Numbers** PL 0010/0150 Infusion. PL 0010/0146 Tablets. **Basic NHS Cost** 10 tablets £750, 100mg Infusion £12.00. **Date of Preparation** April 1988. **Further information on request from:** Bayer UK Limited, Pharmaceutical Business Group, Bayer House, Strawberry Hill, Newbury, Berkshire RG13 1JA. Tel: (0635) 39000. © Bayer UK Limited, April 1988. © Registered trademark Bayer Germany. Bayer, Baypharm, and  are trademarks of Bayer Germany.

 Baypharm 