



**LIBRARIES**  
UNIVERSITY OF WISCONSIN - MADISON

## Dimacol advertisement.

[s.l.]: [s.n.], 1971

<https://digital.library.wisc.edu/1711.dl/RGFFKCCDINDH38U>

<http://rightsstatements.org/vocab/InC/1.0/>

The libraries provide public access to a wide range of material, including online exhibits, digitized collections, archival finding aids, our catalog, online articles, and a growing range of materials in many media.

When possible, we provide rights information in catalog records, finding aids, and other metadata that accompanies collections or items. However, it is always the user's obligation to evaluate copyright and rights issues in light of their own use.

**for coughs / for colds**



**for convenience...**

# The formula based on cough and cold logic:

For your coughing patient who also has a cold, Dimacol is just what the doctor ordered for comprehensive, symptomatic relief. Each of the four agents in the rational formula is patient proven, time tested... a leader in its class.

**Pseudoephedrine** (*Decongestant/Bronchodilator*)—Exerts a gentle vasoconstrictive effect that promptly reduces edema and congestion of mucous membranes in nasal and sinus passages. Also provides a bronchodilator effect.

**Pheniramine Maleate** (*Antihistamine*)—A potent, clinically proven agent for symptomatic relief of allergic manifestations of the common cold. Helps control mucosal edema, congestion, and hypersecretion in nose and sinuses.

**Dextromethorphan** (*Non-Narcotic Antitussive*)—Acts directly on the cough center to suppress the coughing reflex for 6 to 8 hours. Same order of effectiveness as codeine as a cough suppressant, but not habit forming or constipating.

**Glyceryl Guaiacolate** (*Expectorant*)—Increases the output of lower respiratory tract fluid (RTF). Increased RTF volume exerts a demulcent effect on tracheobronchial mucosa, promotes ciliary action, and aids the removal of inspissated mucus.

**Contraindications:** Hypersensitivity to any of the ingredients. Dimacol should not be administered to patients receiving MAO inhibitors. Not recommended for use during pregnancy. **Precautions:** Administer with caution in the presence of hypertension, heart disease, peripheral vascular disease, diabetes, or hyperthyroidism. Patients should be warned that this product may cause drowsiness in some cases. If this occurs, patients should be advised against driving or

engaging in operations which require alertness. As with all products containing sympathomimetic amines, use with caution in patients with prostatic hypertrophy or glaucoma. **Adverse Reactions:** The following adverse reactions may possibly occur: nausea, vomiting, dry mouth, nervousness, insomnia, drowsiness. **Dosage:** Adults, and children 12 years of age or older: 1 or 2 capsules or teaspoonfuls three or four times a day.

**Dimacol** CAPSULES/LIQUID™

Each capsule or 5 cc. (1 teaspoonful—alcohol 4.75%) contains:  
Pseudoephedrine hydrochloride . . . . . 30 mg.  
Pheniramine maleate . . . . . 7.5 mg.  
Dextromethorphan hydrobromide . . . . . 15 mg.  
Glyceryl guaiacolate . . . . . 100 mg.

**AH-ROBINS**

A. H. Robins Company, Richmond, Va.