



LIBRARIES
UNIVERSITY OF WISCONSIN - MADISON

Zantac advertisement.

[s.l.]: [s.n.], 1988

<https://digital.library.wisc.edu/1711.dl/UP4KN66RLFVCC8L>

<http://rightsstatements.org/vocab/InC/1.0/>

The libraries provide public access to a wide range of material, including online exhibits, digitized collections, archival finding aids, our catalog, online articles, and a growing range of materials in many media.

When possible, we provide rights information in catalog records, finding aids, and other metadata that accompanies collections or items. However, it is always the user's obligation to evaluate copyright and rights issues in light of their own use.

ZANTAC® 300 Tablets
(ranitidine hydrochloride)

The following is a brief summary only. Before prescribing, see complete prescribing information in ZANTAC® product labeling.

INDICATIONS AND USAGE: ZANTAC® is indicated in:

1. Short-term treatment of **active duodenal ulcer**. Most patients heal within four weeks.
2. **Maintenance therapy** for duodenal ulcer patients at reduced dosage after healing of acute ulcers.
3. The treatment of **pathological hypersecretory conditions** (eg, Zollinger-Ellison syndrome and systemic mastocytosis).
4. Short-term treatment of **active, benign gastric ulcer**. Most patients heal within six weeks and the usefulness of further treatment has not been demonstrated.
5. Treatment of **gastroesophageal reflux disease (GERD)**. Symptomatic relief commonly occurs within one or two weeks after starting therapy. Therapy for longer than six weeks has not been studied.

In active duodenal ulcer; active, benign gastric ulcer; hypersecretory states; and GERD, concomitant antacids should be given as needed for relief of pain.

CONTRAINDICATIONS: ZANTAC® is contraindicated for patients known to have hypersensitivity to the drug.

PRECAUTIONS: General: 1. Symptomatic response to ZANTAC® therapy does not preclude the presence of gastric malignancy.

2. Since ZANTAC is excreted primarily by the kidney, dosage should be adjusted in patients with impaired renal function (see DOSAGE AND ADMINISTRATION). Caution should be observed in patients with hepatic dysfunction since ZANTAC is metabolized in the liver.

Laboratory Tests: False-positive tests for urine protein with Multistix® may occur during ZANTAC therapy, and therefore testing with sulfosalicylic acid is recommended.

Drug Interactions: Although ZANTAC has been reported to bind weakly to cytochrome P-450 in vitro, recommended doses of the drug do not inhibit the action of the cytochrome P-450-linked oxygenase enzymes in the liver. However, there have been isolated reports of drug interactions that suggest that ZANTAC may affect the bioavailability of certain drugs by some mechanism as yet unidentified (eg, a pH-dependent effect on absorption or a change in volume of distribution).

Carcinogenesis, Mutagenesis, Impairment of Fertility: There was no indication of tumorigenic or carcinogenic effects in lifespan studies in mice and rats at doses up to 2,000 mg/kg/day.

Ranitidine was not mutagenic in standard bacterial tests (*Salmonella*, *Escherichia coli*) for mutagenicity at concentrations up to the maximum recommended for these assays.

In a dominant lethal assay, a single oral dose of 1,000 mg/kg to male rats was without effect on the outcome of two matings per week for the next nine weeks.

Pregnancy: Teratogenic Effects: Pregnancy Category B: Reproduction studies have been performed in rats and rabbits at doses up to 160 times the human dose and have revealed no evidence of impaired fertility or harm to the fetus due to ZANTAC. There are, however, no adequate and well-controlled studies in pregnant women. Because animal reproduction studies are not always predictive of human response, this drug should be used during pregnancy only if clearly needed.

Nursing Mothers: ZANTAC is secreted in human milk. Caution should be exercised when ZANTAC is administered to a nursing mother.

Pediatric Use: Safety and effectiveness in children have not been established.

Use in Elderly Patients: Ulcer healing rates in elderly patients (65 to 82 years of age) were no different from those in younger age groups. The incidence rates for adverse events and laboratory abnormalities were also not different from those seen in other age groups.

ADVERSE REACTIONS: The following have been reported as events in clinical trials or in the routine management of patients treated with ZANTAC®. The relationship to ZANTAC therapy has been unclear in many cases. Headache, sometimes severe, seems to be related to ZANTAC administration.

Central Nervous System: Rarely, malaise, dizziness, somnolence, insomnia, and vertigo. Rare cases of reversible mental confusion, agitation, depression, and hallucinations have been reported, predominantly in severely ill elderly patients. Rare cases of reversible blurred vision suggestive of a change in accommodation have been reported.

Cardiovascular: Rare reports of tachycardia, bradycardia, and premature ventricular beats.

Gastrointestinal: Constipation, diarrhea, nausea/vomiting, and abdominal discomfort/pain.

Hepatic: In normal volunteers, SGPT values were increased to at least twice the pretreatment levels in 6 of 12 subjects receiving 100 mg qid IV for seven days, and in 4 of 24 subjects receiving 50 mg qid IV for five days. With oral administration there have been occasional reports of reversible hepatitis, hepatocellular or hepatocanalicular or mixed, with or without jaundice.

Musculoskeletal: Rare reports of arthralgias.

Hematologic: Reversible blood count changes (leukopenia, granulocytopenia, thrombocytopenia) have occurred in a few patients. Rare cases of agranulocytosis or of pancytopenia, sometimes with marrow hypoplasia, have been reported.

Endocrine: Controlled studies in animals and man have shown no stimulation of any pituitary hormone by ZANTAC and no antiandrogenic activity, and cimetidine-induced gynecomastia and impotence in hypersecretory patients have resolved when ZANTAC has been substituted. However, occasional cases of gynecomastia, impotence, and loss of libido have been reported in male patients receiving ZANTAC, but the incidence did not differ from that in the general population.

Integumentary: Rash, including rare cases suggestive of mild erythema multiforme, and, rarely, alopecia.

Other: Rare cases of hypersensitivity reactions (eg, bronchospasm, fever, rash, eosinophilia) and small increases in serum creatinine.

OVERDOSAGE: Information concerning possible overdosage and its treatment appears in the full prescribing information.

DOSAGE AND ADMINISTRATION: Active Duodenal Ulcer: The current recommended adult oral dosage is 150 mg twice daily. An alternate dosage of 300 mg once daily at bedtime can be used for patients in whom dosing convenience is important. The advantages of one treatment regimen compared to the other in a particular patient population have yet to be demonstrated.

Maintenance Therapy: The current recommended adult oral dosage is 150 mg at bedtime.

Pathological Hypersecretory Conditions (such as Zollinger-Ellison syndrome): The current recommended adult oral dosage is 150 mg twice a day. In some patients it may be necessary to administer ZANTAC® 150-mg doses more frequently. Doses should be adjusted to individual patient needs, and should continue as long as clinically indicated. Doses up to 6 g/day have been employed in patients with severe disease.

Benign Gastric Ulcer: The current recommended adult oral dosage is 150 mg twice a day.

GERD: The current recommended adult oral dosage is 150 mg twice a day.

Dosage Adjustment for Patients with Impaired Renal Function: On the basis of experience with a group of subjects with severely impaired renal function treated with ZANTAC, the recommended dosage in patients with a creatinine clearance less than 50 ml/min is 150 mg every 24 hours. Should the patient's condition require, the frequency of dosing may be increased to every 12 hours or even further with caution. Hemodialysis reduces the level of circulating ranitidine. Ideally, the dosage schedule should be adjusted so that the timing of a scheduled dose coincides with the end of hemodialysis.

HOW SUPPLIED: ZANTAC® 300 Tablets (ranitidine hydrochloride equivalent to 300 mg of ranitidine) are yellow, capsule-shaped tablets embossed with "ZANTAC 300" on one side and "Glaxo" on the other. They are available in bottles of 30 tablets (NDC 0173-0393-40) and unit dose packs of 100 tablets (NDC 0173-0393-47).

ZANTAC® 150 Tablets (ranitidine hydrochloride equivalent to 150 mg of ranitidine) are white tablets embossed with "ZANTAC 150" on one side and "Glaxo" on the other. They are available in bottles of 60 tablets (NDC 0173-0344-42) and unit dose packs of 100 tablets (NDC 0173-0344-47).

Store between 15° and 30°C (59° and 86°F) in a dry place. Protect from light. Replace cap securely after each opening.

Glaxo

May 1988

Glaxo Inc.
Research Triangle Park, NC 27709

One Of A Kind



Zantac[®]
ranitidine HCl/Glaxo 150 mg and
300 mg tablets

Please see Brief Summary of Prescribing Information on adjacent page.

Glaxo/ROCHE

Zantac® 150 Tablets
(ranitidine hydrochloride)

CONDENSED BRIEF SUMMARY

Zantac® 300 Tablets
(ranitidine hydrochloride)

Zantac® Syrup
(ranitidine hydrochloride)

The following is a brief summary only. Before prescribing, see complete prescribing information in Zantac® product labeling.

INDICATIONS AND USAGE: Zantac® is indicated in:

1. Short-term treatment of **active duodenal ulcer**. Most patients heal within four weeks.
2. **Maintenance therapy** for duodenal ulcer patients at reduced dosage after healing of acute ulcers.
3. The treatment of **pathological hypersecretory conditions** (eg, Zollinger-Ellison syndrome and systemic mastocytosis).
4. Short-term treatment of **active, benign gastric ulcer**. Most patients heal within six weeks and the usefulness of further treatment has not been demonstrated.
5. Treatment of **gastroesophageal reflux disease (GERD)**.

Symptomatic relief commonly occurs within one or two weeks after starting therapy and is maintained throughout a six-week course of therapy.

In active duodenal ulcer; active, benign gastric ulcer; hypersecretory states; and GERD, concomitant antacids should be given as needed for relief of pain.

CONTRAINDICATIONS: Zantac® is contraindicated for patients known to have hypersensitivity to the drug.

PRECAUTIONS: General: 1. Symptomatic response to Zantac® therapy does not preclude the presence of gastric malignancy. 2. Since Zantac is excreted primarily by the kidney, dosage should be adjusted in patients with impaired renal function (see DOSAGE AND ADMINISTRATION). Caution should be observed in patients with hepatic dysfunction since Zantac is metabolized in the liver.

Laboratory Tests: False-positive tests for urine protein with Multistix® may occur during Zantac therapy, and therefore testing with sulfosalicylic acid is recommended.

Drug Interactions: Although recommended doses of Zantac do not inhibit the action of cytochrome P-450 enzymes in the liver, there have been isolated reports of drug interactions that suggest that Zantac may affect the bioavailability of certain drugs by some mechanism as yet unidentified (eg, a pH-dependent effect on absorption or a change in volume of distribution).

Pregnancy: Teratogenic Effects: Pregnancy Category B: Reproduction studies have been performed in rats and rabbits at doses up to 160 times the human dose and have revealed no evidence of impaired fertility or harm to the fetus due to Zantac. There are, however, no adequate and well-controlled studies in pregnant women. Because animal reproduction studies are not always predictive of human response, this drug should be used during pregnancy only if clearly needed.

Nursing Mothers: Zantac is secreted in human milk. Caution should be exercised when Zantac is administered to a nursing mother.

Pediatric Use: Safety and effectiveness in children have not been established.

ADVERSE REACTIONS: Headache, sometimes severe, seems to be related to Zantac® administration. Constipation, diarrhea, nausea/vomiting, abdominal discomfort/pain, and, rarely, pancreatitis have been reported. There have been rare reports of malaise, dizziness, somnolence, insomnia, vertigo, tachycardia, bradycardia, atrioventricular block, premature ventricular beats, and arthralgias. Rare cases of reversible mental confusion, agitation, depression, and hallucinations have been reported, predominantly in severely ill elderly patients. Rare cases of reversible blurred vision suggestive of a change in accommodation have been reported.

In normal volunteers, SGPT values were increased to at least twice the pretreatment levels in 6 of 12 subjects receiving 100 mg qid intravenously for seven days, and in 4 of 24 subjects receiving 50 mg qid intravenously for five days. There have been occasional reports of hepatitis, hepatocellular or hepatocanalicular or mixed, with or without jaundice. In such circumstances, ranitidine should be immediately discontinued. These events are usually reversible, but in exceedingly rare circumstances death has occurred.

Blood count changes (leukopenia, granulocytopenia, thrombocytopenia) have occurred in a few patients. These were usually reversible. Rare cases of agranulocytosis, pancytopenia, sometimes with marrow hypoplasia, and aplastic anemia have been reported.

Although controlled studies have shown no antiandrogenic activity, occasional cases of gynecomastia, impotence, and loss of libido have been reported in male patients receiving Zantac, but the incidence did not differ from that in the general population.

Incidents of rash, including rare cases suggestive of mild erythema multiforme, and, rarely, alopecia, have been reported, as well as rare cases of hypersensitivity reactions (eg, bronchospasm, fever, rash, eosinophilia), anaphylaxis, angioneurotic edema, and small increases in serum creatinine.

OVERDOSAGE: Information concerning possible overdose and its treatment appears in the full prescribing information.

DOSAGE AND ADMINISTRATION: (See complete prescribing information in Zantac® product labeling.)

Dosage Adjustment for Patients with Impaired Renal Function: On the basis of experience with a group of subjects with severely impaired renal function treated with Zantac, the recommended dosage in patients with a creatinine clearance less than 50 ml/min is 150 mg or 10 ml (2 teaspoonfuls equivalent to 150 mg of ranitidine) every 24 hours. Should the patient's condition require, the frequency of dosing may be increased to every 12 hours or even further with caution. Hemodialysis reduces the level of circulating ranitidine. Ideally, the dosage schedule should be adjusted so that the timing of a scheduled dose coincides with the end of hemodialysis.

HOW SUPPLIED: Zantac® 300 Tablets (ranitidine hydrochloride equivalent to 300 mg of ranitidine) are yellow, capsule-shaped tablets embossed with "ZANTAC 300" on one side and "Glaxo" on the other. They are available in bottles of 30 (NDC 0173-0393-40) tablets and unit dose packs of 100 (NDC 0173-0393-47) tablets.

Zantac® 150 Tablets (ranitidine hydrochloride equivalent to 150 mg of ranitidine) are white tablets embossed with "ZANTAC 150" on one side and "Glaxo" on the other. They are available in bottles of 60 (NDC 0173-0344-42) and 100 (NDC 0173-0344-09) tablets and unit dose packs of 100 (NDC 0173-0344-47) tablets.

Store between 15° and 30° C (59° and 86° F) in a dry place.

Protect from light. Replace cap securely after each opening.
Zantac® Syrup, a clear, peppermint-flavored liquid, contains 16.8 mg of ranitidine hydrochloride equivalent to 15 mg of ranitidine per 1 ml in bottles of 16 fluid ounces (one pint) (NDC 0173-0383-54).

Store between 4° and 25° C (39° and 77° F). Dispense in tight, light-resistant containers as defined in the USP/NF.

September 1990

 **Glaxo Pharmaceuticals™**
DIVISION OF GLAXO INC.
Research Triangle Park, NC 27709