



LIBRARIES
UNIVERSITY OF WISCONSIN - MADISON

Merck advertisement.

[s.l.]: [s.n.], 1927

<https://digital.library.wisc.edu/1711.dl/O3KIJXMGCKQP48Z>


<http://rightsstatements.org/vocab/InC/1.0/>

The libraries provide public access to a wide range of material, including online exhibits, digitized collections, archival finding aids, our catalog, online articles, and a growing range of materials in many media.

When possible, we provide rights information in catalog records, finding aids, and other metadata that accompanies collections or items. However, it is always the user's obligation to evaluate copyright and rights issues in light of their own use.

Erythrol Tetranitrate

MERCK [Erythrityl Tetranitrate]



For reduction of high blood pressure
For the relief of, and as a prophylactic for,
attacks of angina pectoris
For vascular diseases

- Vasodilator action slower and more lasting than other commonly used nitrites.
- Vasodilatation usually begins within about fifteen minutes after administration and persists for 3 to 4 hours.
- Individualized dosage may be continued with sustained effect.

Erythrol Tetranitrate Merck (Erythrityl Tetranitrate) is supplied in bottles of 50 $\frac{1}{4}$ -grain tablets; tubes of 24, and bottles of 100 $\frac{1}{2}$ -grain tablets—readily obtainable from your pharmacist.



MERCK & CO. Inc.

Manufacturing Chemists

RAHWAY, N. J.