

## Declomycin advertisement.


[s.l.]: [s.n.], 1960

https://digital.library.wisc.edu/1711.dl/BRVEOAK4NLNQY8U

http://rightsstatements.org/vocab/InC/1.0/

The libraries provide public access to a wide range of material, including online exhibits, digitized collections, archival finding aids, our catalog, online articles, and a growing range of materials in many media.

When possible, we provide rights information in catalog records, finding aids, and other metadata that accompanies collections or items. However, it is always the user's obligation to evaluate copyright and rights issues in light of their own use.



CAPSULES, 150 mg., 75 mg. *Dosage:* Average infections—150 mg. four times daily. Severe infections—Initial dose of 300 mg., then 150 mg. every six hours.

PEDIATRIC DROPS, 60 mg./cc. in 10 cc. bottle with calibrated, plastic dropper. Dosage: 1 to 2 drops (3 to 6 mg.) per pound body weight per day—divided into four doses.

SYRUP, 75 mg./5 cc. teaspoonful (cherry-flavored). Dosage: 3 to 6 mg. per pound body weight per day — divided into four doses.

PRECAUTIONS - As with other antibiotics, DECLO MYCIN may occasionally give rise to glossitis, stomatitis proctitis, nausea, diarrhea, vaginitis or dermatitis, photodynamic reaction to sunlight has been observed a few patients on DECLOMYCIN. Although reversibly discontinuing therapy, patients should avoid exposure to intense sunlight. If adverse reaction or idiosynamic and intense sunlight of the sun of the sun

crasy occurs, discontinue medication.

Overgrowth of nonsusceptible organisms is a possibilit with DECLOMYCIN, as with other antibiotics, and dentand that the patient be kept under constant observation

added measure of protection

DEMETHYLCHLORTETRACYCLINE LEDERLE

against relapse— up to 6 days' activity on 4 days' dosage

against secondary infection— sustained high activity levels

against "problem" pathogens - positive broad-spectrum antibiosis