



LIBRARIES
UNIVERSITY OF WISCONSIN - MADISON

Brondecon advertisement.

[s.l.]: [s.n.], 1962

<https://digital.library.wisc.edu/1711.dl/5PMEWONZVKMBQ8Q>

<http://rightsstatements.org/vocab/InC/1.0/>

The libraries provide public access to a wide range of material, including online exhibits, digitized collections, archival finding aids, our catalog, online articles, and a growing range of materials in many media.

When possible, we provide rights information in catalog records, finding aids, and other metadata that accompanies collections or items. However, it is always the user's obligation to evaluate copyright and rights issues in light of their own use.

new expectorant-bronchodilator

BRONDECON^{T.M.}

tablets/elixir

Tablet (each): 200 mg. oxtriphylline and 100 mg. glyceryl guaiacolate. Elixir (5 ml. tsp.): 100 mg. oxtriphylline and 50 mg. glyceryl guaiacolate.

thorough symptomatic control of

BRONCHITIS

Concentrated new Brondecon thins tenacious mucus and makes breathing easier for your bronchitis patient with *lower, more economical dosages*. Brondecon attains peak theophylline blood levels faster and facilitates a more productive cough. Its two components — glyceryl guaiacolate, a proven expectorant, and oxtriphylline, the well-tolerated bronchodilator—remain fully effective, even over prolonged periods.

During acute attacks... an increased dose may be used to initiate therapy. Patients over 12 years—3 tablets or 6 teaspoonfuls; 2 to 12 years—1 teaspoonful/20 lb. body weight. Usual doses should be employed thereafter.

Supplied: Tablets in bottles of 100. Cherry-flavored elixir in 8-ounce bottles.

Dosage: Tablets—Patients over 12 years: 1 tablet q.i.d. Elixir—Patients over 12 years: 2 teaspoonfuls q.i.d.; 6 to 12 years: 1 teaspoonful/60 lb. body weight, q.i.d. Brondecon should be taken after meals.

Side effects: Minimal, including gastric distress and, occasionally, palpitation and CNS stimulation.


Precautions: Not to be administered more often than every 4 hours or within 12 hours following oral, parenteral or rectal use of other xanthine-containing preparations. Avoid simultaneous use of other xanthines in all patients, especially children.

WARNER - CHILCOTT




Warner-Chilcott, Morris Plains, N.J.
Makers of Coly-Mycin Gelusil Peritrate Proloid Tedral


GP33



HELPS FLUSH OUT MUCUS



KEEPS BRONCHI OPEN



LETS AIR THROUGH