



LIBRARIES
UNIVERSITY OF WISCONSIN - MADISON

Nephenalin advertisement.

[s.l.]: [s.n.], 1952

<https://digital.library.wisc.edu/1711.dl/YC3UAMUSYHGHJ8Z>

<http://rightsstatements.org/vocab/InC/1.0/>

The libraries provide public access to a wide range of material, including online exhibits, digitized collections, archival finding aids, our catalog, online articles, and a growing range of materials in many media.

When possible, we provide rights information in catalog records, finding aids, and other metadata that accompanies collections or items. However, it is always the user's obligation to evaluate copyright and rights issues in light of their own use.

**In childhood asthma . . . there's an excellent chance
your little patients will prefer the fast-acting, long-lasting,
bright-red NEPHENALIN PEDIATRIC tablet.**

Two treatments in one tablet: first, hold under the tongue for five minutes; then swallow. This provides rapid bronchodilation from sublingual aludrine, then

4-hour follow-through protection from theophylline-ephedrine-phenobarbital.


Dosage: One tablet as needed (up to 5 tablets a day). In bottles of 20 and 100.



ALUDRINE (Isopropyl Arterenol) HCl → 5 mg.
(Sublingually: for rapid relief)

THEOPHYLLINE 1½ gr.
EPHEDRINE SULFATE ⅜ gr.
PHENOBARBITAL ⅛ gr.
(for 4-hour protection)

In red coating



Nephenalin[®]
PEDIATRIC (RED)

Thos. Leeming & Co. Inc. 155 East 44th Street, New York 17, N.Y.