



LIBRARIES
UNIVERSITY OF WISCONSIN - MADISON

Co Pyronil advertisement.

[s.l.]: [s.n.], 1966

<https://digital.library.wisc.edu/1711.dl/YIBADLS5TYBS58R>

<http://rightsstatements.org/vocab/InC/1.0/>

The libraries provide public access to a wide range of material, including online exhibits, digitized collections, archival finding aids, our catalog, online articles, and a growing range of materials in many media.

When possible, we provide rights information in catalog records, finding aids, and other metadata that accompanies collections or items. However, it is always the user's obligation to evaluate copyright and rights issues in light of their own use.



For more complete relief of allergic symptoms

Consider Co-Pyronil® when allergic patients need more complete relief than single antihistamines alone can give.

Each Pulvule® provides:

Two antihistamines—to give both rapid relief and prolonged activity (up to twelve hours).

A vasoconstrictor—to provide the synergistic action of a decongestant.

These ingredients act quickly to relieve the congestion and discomfort of pollinosis, the burning and weeping of ocular allergies, the itch and swelling of urticaria and minor drug reactions. For secondary symptoms concurrent with these allergies, Co-Pyronil also helps to give more complete relief.

Contraindications and Precautions: Hypersensitivity to antihistamines, though rare, is a contraindication to their use. Co-Pyronil should be

used with caution in the presence of hypertension, cardiovascular-renal disease, and hyperthyroidism. As with any preparation containing antihistamines and sympathomimetics, overdosage may produce excessive central-nervous-system depression or stimulation.

Side-Effects: Drowsiness is sometimes reported at the beginning of treatment but is usually transient. In rare instances, symptoms of sympathetic overstimulation may be noted from the vasoconstrictor ingredient in Co-Pyronil.

Dosage Range: One Pulvule every four, eight, or twelve hours, depending on the severity of the symptoms.

Co-Pyronil®

Each Pulvule contains 15 mg. Pyronil® (pyrrobutamine, Lilly), 25 mg. Histadyl® (methapyrilene hydrochloride, Lilly), and 12.5 mg. Clopane® Hydrochloride (cyclopentamine hydrochloride, Lilly).

600387

Additional information available to physicians upon request. Eli Lilly and Company, Indianapolis, Indiana.