



LIBRARIES
UNIVERSITY OF WISCONSIN - MADISON

Ornade Spansule advertisement.

[s.l.]: [s.n.], 1967

<https://digital.library.wisc.edu/1711.dl/YDPBQHK22F6RG8G>

<http://rightsstatements.org/vocab/InC/1.0/>

The libraries provide public access to a wide range of material, including online exhibits, digitized collections, archival finding aids, our catalog, online articles, and a growing range of materials in many media.

When possible, we provide rights information in catalog records, finding aids, and other metadata that accompanies collections or items. However, it is always the user's obligation to evaluate copyright and rights issues in light of their own use.

One of the best things you can do
for the cold sufferer



Ornade[®] Spansule[®]

brand of
sustained release capsules

Trademark

Each capsule contains 8 mg. of Teldrin[®] (brand of chlorpheniramine maleate); 50 mg. of phenylpropanolamine hydrochloride; 2.5 mg. of isopropamide, as the iodide.

Prompt relief from nasal congestion and hypersecretion due to colds.

Before prescribing, see complete prescribing information in SK&F literature or *PDR*.

Contraindications: Glaucoma, prostatic hypertrophy, stenosing peptic ulcer, pyloroduodenal or bladder neck obstruction.

Precautions: Use cautiously in the presence of hypertension, hyperthyroidism, coronary artery disease; warn vehicle or machine operators of possible drowsiness.

Usage in Pregnancy: Use in pregnancy, nursing mothers and women who might bear children only when potential benefits have been weighed against possible hazards.

Note: The iodine in isopropamide iodide may alter PBI test results and will suppress I¹³¹ uptake; discontinue 'Ornade' one week before these tests.

Adverse Reactions: Drowsiness; excessive dryness of nose, throat or mouth; nervousness; insomnia.

Other known possible adverse reactions of the individual ingredients: nausea, vomiting, diarrhea, rash, dizziness, fatigue, tightness of chest, abdominal pain, irritability, tachycardia, headache, incoordination, tremor, difficulty in urination. Thrombocytopenia, leukopenia and convulsions have been reported.

Supplied: Bottles of 50 capsules.

One capsule q12h for round-the-clock relief

**SK
&F** Smith Kline & French Laboratories