



LIBRARIES
UNIVERSITY OF WISCONSIN - MADISON

Mebaroin advertisement.

[s.l.]: [s.n.], 1950

<https://digital.library.wisc.edu/1711.dl/VGRFHFSBOQ67M8E>

<http://rightsstatements.org/vocab/InC/1.0/>

The libraries provide public access to a wide range of material, including online exhibits, digitized collections, archival finding aids, our catalog, online articles, and a growing range of materials in many media.

When possible, we provide rights information in catalog records, finding aids, and other metadata that accompanies collections or items. However, it is always the user's obligation to evaluate copyright and rights issues in light of their own use.

FOR *epileptic* CONTROL-

grand mal and petit mal

New

Mebaroin TRADEMARK

Highly Effective • Well Tolerated • Relatively Tasteless

In a comparative study of 108 patients^{1,2} with idiopathic epilepsy, by far the best results (85.6 per cent controlled or improved) were obtained with a combination of Mebaral and diphenylhydantoin. This combination achieved the closest approach to the ideal of maximum control of seizures with minimal toxicity. Each tablet of Mebaroin contains 90 mg. Mebaral® and 60 mg. diphenylhydantoin.

Dosage: Average adult dose: 1 or 2 tablets three times daily. For children over 6 years, 2 or 3 tablets daily; younger children ½ tablet once or twice daily. Treatment is best begun with a small dose which is increased gradually until the individual optimum is established. The usual caution should be observed when changing from another antiepileptic drug to Mebaroin.

Supplied: Bottles of 100 and 1000 scored tablets.

1. Meller, R. L., and Resch, J. A.: *Bull. Univ. Minnesota Hosp.*, 20:78, Oct. 8, 1948.

2. Meller, R. L., and Resch, J. A.: *Postgrad. Med.*, 6:452, Dec., 1949.

Wintrop Stearns INC.
NEW YORK 18, N. Y. WINDSOR, ONT.

Mebaral, trademark reg. U.S. & Canada, brand of mephobarbital