



**LIBRARIES**  
UNIVERSITY OF WISCONSIN - MADISON

## Visicol advertisement.

[s.l.]: [s.n.], 2003

<https://digital.library.wisc.edu/1711.dl/MYVWAJR2LKGFF8U>

<http://rightsstatements.org/vocab/InC/1.0/>

The libraries provide public access to a wide range of material, including online exhibits, digitized collections, archival finding aids, our catalog, online articles, and a growing range of materials in many media.

When possible, we provide rights information in catalog records, finding aids, and other metadata that accompanies collections or items. However, it is always the user's obligation to evaluate copyright and rights issues in light of their own use.

For colonoscopy bowel preparation...

“You mean there’s a tablet prep!”

- **Significantly less nausea, vomiting and bloating vs. large-volume liquid prep<sup>1</sup>**
- **Excellent colon cleansing<sup>1</sup>**
- **High patient compliance<sup>1</sup> & preference<sup>1,2</sup>**


Visit [www.Visicol.com](http://www.Visicol.com) for more information about patient-preferred VISICOL® Tablets



**VISICOL®**

(sodium phosphate monobasic monohydrate, USP,  
sodium phosphate dibasic anhydrous, USP)

*The Preferred Prep*



**Reference: 1.** Kastenberg D, et al. Efficacy and safety of sodium phosphate tablets compared with PEG solution in colon cleansing, two identically designed, randomized, controlled, parallel group, multicenter phase III trials. *Gastrointest Endoscopy*.2001;54:705-713.  
**2.** Data on file

#### **WARNINGS**

**Visicol® Tablets are not to be used in patients with congestive heart failure, ascites, unstable angina pectoris, gastric retention, ileus or acute obstruction or pseudo-obstruction, severe chronic constipation, bowel perforation, acute colitis, toxic megacolon or hypomotility syndrome.**

**Use with caution in patients with impaired renal function, pre-existing electrolyte disturbances, or people taking drugs that affect electrolyte levels.**