



## Isordil Chewable advertisement.

[s.l.]: [s.n.], 1976

<https://digital.library.wisc.edu/1711.dl/M5KIAEQDE6ZGG8R>

<http://rightsstatements.org/vocab/InC/1.0/>

The libraries provide public access to a wide range of material, including online exhibits, digitized collections, archival finding aids, our catalog, online articles, and a growing range of materials in many media.

When possible, we provide rights information in catalog records, finding aids, and other metadata that accompanies collections or items. However, it is always the user's obligation to evaluate copyright and rights issues in light of their own use.

chewable!  
chewable!  
chewable!  
chewable!



announcing

**ISORDIL<sup>®</sup> CHEWABLE TABLETS 10 mg**  
**(ISOSORBIDE DINITRATE)**

for those angina\* patients for whom  
you prefer a chewable isosorbide dinitrate.  
This higher strength chewable tablet makes  
increased dosage practical, and the tablet is scored  
for a lower starting dose and titration of dosage.

**\*Indications:** Based on a review of this drug by the National Academy of Sciences — National Research Council and/or other information, FDA has classified the indications as follows:  
"Probably" effective: When taken by the chewable route, Isordil Chewable Tablets are indicated for the treatment of acute anginal attacks and for prophylaxis in situations likely to provoke such attacks.  
Final classification of the less-than-effective indications requires further investigation.

**Contraindication:** Idiosyncrasy to this drug.

**Warnings:** Data supporting the use of nitrites during the early days of the acute phase of myocardial infarction (the period during which clinical and laboratory findings are unstable) are insufficient to establish safety.

**Precautions:** Tolerance to this drug and cross-tolerance to other nitrites and nitrates may occur.

**Adverse Reactions:** Cutaneous vasodilation with flushing. Headache is common and may be severe and persistent. Transient episodes of dizziness and

weakness as well as other signs of cerebral ischemia associated with postural hypotension may occasionally develop. This drug can act as a physiological antagonist to norepinephrine, acetylcholine, histamine, and many other agents. An occasional individual exhibits marked sensitivity to the hypotensive effects of nitrite, and severe responses (nausea, vomiting, weakness, restlessness, pallor, perspiration and collapse) can occur even with the usual therapeutic dose. Alcohol may enhance this effect. Drug rash and/or exfoliative dermatitis may occasionally occur.

*Consult direction circular before prescribing.*

May we send you reprints, detailed information and/or professional samples?

**IVES LABORATORIES INC.**

685 Third Avenue  
New York, New York 10017

DEDICATED TO IMPROVING THE QUALITY OF LIFE  
THROUGH MEDICINE<sup>®</sup>

