

Cardilate advertisement.

[s.l.]: [s.n.], 1965

<https://digital.library.wisc.edu/1711.dl/AE2OAUZFZ57N2Q85>

<http://rightsstatements.org/vocab/InC/1.0/>

The libraries provide public access to a wide range of material, including online exhibits, digitized collections, archival finding aids, our catalog, online articles, and a growing range of materials in many media.

When possible, we provide rights information in catalog records, finding aids, and other metadata that accompanies collections or items. However, it is always the user's obligation to evaluate copyright and rights issues in light of their own use.

Protection with 'CARDILATE'[®] Erythrityl Tetranitrate Tablets

With 'Cardilate' brand Erythrityl Tetranitrate, your patient can receive prompt and prolonged protection against recurrent attacks of angina pectoris. Erythrityl Tetranitrate usually acts within five minutes sublingually, and within 30 minutes orally, and provides up to four hours of sustained relief. It is particularly suitable for prophylactic administration prior to anticipated periods of stress, such as exertion or excitement. 'Cardilate' brand Erythrityl Tetranitrate decreases the need for nitroglycerin, and permits more normal activities since it increases exercise tolerance in most cases.

Announcing New Flavored Chewable Tablets 'CARDILATE'[®] Chewable Erythrityl Tetranitrate brand Tablets 10 mg.

- dual role: provides rapid onset of action (in less than 5 minutes) from buccal absorption and prolonged effect (up to 4 hours) from gastrointestinal absorption
- convenient to take—no need for water
- pleasant wintergreen flavor
- adds another dosage form to the familiar oral or sublingual tablets

Indications: For the prophylaxis and long-term treatment of patients with frequent or recurrent anginal pain and reduced exercise tolerance, rather than for the treatment of the acute attack of angina pectoris, since its onset of action is somewhat slower than that of nitroglycerin.

Precaution: As with other effective nitrites, some fall in blood pressure may occur with large doses. Caution should be observed in patients with a history of recent cerebral hemorrhage. Although therapy permits more normal activity, the patient should not be allowed to misinterpret freedom from anginal attacks as a signal to drop all restrictions.

Side effects: No serious side effects have been reported. In sublingual therapy, a tingling sensation (like that of nitroglycerin) may sometimes be noted at the point of tablet contact with the mucous membrane. If objectionable, this may be mitigated by placing the tablet in the buccal pouch. As with nitroglycerin or other effec-

tive nitrites, temporary vascular headache may occur during the first few days of therapy. This can be controlled by temporary dosage reduction in order to allow adjustment of the cerebral hemodynamics to the initial marked cerebral vasodilation. These headaches usually disappear within one week of continuous therapy but may be minimized by the administration of analgesics such as 'Empirin'® Compound.

Mild gastrointestinal disturbances occur occasionally with larger doses and may be controlled by reducing the dose temporarily.

Supplied: In 5, 10 and 15 mg. scored tablets and 10 mg. chewable tablets. Bottles of 100.

Also available for angina complicated by anxiety 'CARDILATE-P' brand Erythrityl Tetranitrate with Phenobarbital.

Erythrityl Tetranitrate 10 mg. with Phenobarbital* 15 mg. scored tablets for oral use only, bottles of 100.

*Warning: May be habit forming.



Complete literature available on request from Professional Services Dept. PML.
BURROUGHS WELLCOME & CO., (U.S.A.) INC., Tuckahoe, N. Y.