



LIBRARIES
UNIVERSITY OF WISCONSIN - MADISON

Ornade advertisement.

[s.l.]: [s.n.], 1972

<https://digital.library.wisc.edu/1711.dl/KXYRYO3A4JTYP9A>

<http://rightsstatements.org/vocab/InC/1.0/>

The libraries provide public access to a wide range of material, including online exhibits, digitized collections, archival finding aids, our catalog, online articles, and a growing range of materials in many media.

When possible, we provide rights information in catalog records, finding aids, and other metadata that accompanies collections or items. However, it is always the user's obligation to evaluate copyright and rights issues in light of their own use.

Ornade[®]

Each Spanule[®] sustained release capsule contains 8 mg. Teldrin[®] (brand of chlorpheniramine maleate); 50 mg. phenylpropanolamine hydrochloride; 2.5 mg. isopropamide, as the iodide.



to help her live with her cold

one capsule b.i.d. for prompt, prolonged relief
of sneezing, congestion, and watery eyes

Before prescribing, see complete prescribing information in SK&F literature or PDR.

Indications: Upper respiratory congestion and hypersecretion associated with: the common cold; acute and chronic sinusitis; vasomotor rhinitis; allergic rhinitis (hay fever, "rose fever," etc.).

Contraindications: Hypersensitivity to any component; concurrent MAO inhibitor therapy; severe hypertension; bronchial asthma; coronary artery disease; stenosing peptic ulcer; pyloroduodenal or bladder neck obstruction. Children under 6.

Warnings: Caution patients about activities requiring alertness (e.g., operating vehicles or machinery). Warn patients of possible additive effects with alcohol and other CNS depressants.

Usage in Pregnancy: In pregnancy, nursing mothers and women who might bear children, weigh potential benefits against hazards. Inhibition of lactation may occur.

Effect on PBI Determination and I¹³¹ Uptake: Isopropamide iodide may alter PBI test results and will suppress I¹³¹ uptake. Substitute thyroid tests unaffected by exogenous iodides.

Precautions: Use cautiously in persons with cardiovascular disease, glaucoma, prostatic hypertrophy, hyperthyroidism.

Adverse Reactions: Drowsiness, excessive dryness of nose, throat or mouth; nervousness; or insomnia. Also, nausea, vomiting, epigastric distress, diarrhea, rash, dizziness, weakness, chest tightness, angina pain, abdominal pain, irritability, palpitation, headache, incoordination, tremor, dysuria, difficulty in urination, thrombocytopenia, leukopenia, convulsions, hypertension, hypotension, anorexia, constipation, visual disturbances, iodine toxicity (acne, parotitis).

Supplied: Bottles of 50 capsules.

Smith Kline & French Laboratories, Philadelphia