

**Evolution of the family Araceae in the tropics:
Diversification rates, biogeography, and morphological evolution at different
taxonomic and geographic scales**

By

Alejandro Zuluaga Trochez

A thesis submitted in partial fulfillment of the requirements for

the degree of Doctor of Philosophy

(Botany)

at the

UNIVERSITY OF WISCONSIN-MADISON

2015

Date of final oral examination: May 29, 2015

The dissertation is approved by the following members of the final oral committee:

Kenneth M. Cameron, Professor, Botany

Cecile Ané, Associate Professor, Botany and Statistics

David Baum, Professor, Botany

Thomas Givnish, Henry Allan Gleason Professor, Botany

Kenneth J. Sytsma, Professor, Botany

ACKNOWLEDGEMENTS

During the past five years I have had the opportunity of meeting several amazing people who in some way or another have help me to complete my doctoral studies, and I would like to thank every one of them. However, first I want to express my gratitude to all the people who helped before I came to Madison, and who kept supporting me during all the last five years. First, I am grateful to my family, my parents and my brother for their infinite and unconditional support and encouragement. From the distance, they shared with me all the good and not as good moments. I also want to thank to my former undergrad advisors Gloria Galeano and Julio Betancur who introduce me to the study of plants, and who show me with incredible passion the fantastic world of botany. Also, thanks to all my Colombian botanist friends with whom I have shared memorable moments in the field and the lab. Especially, I want to thank Laura Clavijo and Julian Aguirre for all these years of friendship and for all the conversations about plants. Thanks to Laura Clavijo for the company in the field and for all the comments and help with this manuscript.

Equally, I want to express my sincere thanks to my advisor, Ken Cameron, who allowed me to come here and had always be supportive of my ideas (or discouraged the bad ones with a little of dark humor). Thank you to all members of the Cameron Lab, Rafael, Brian, Giovanni, Matthew, Alfonso and James, especially to Rafael who kindly helped me during the first years in Madison. I am also thankful to all my Committee members, Cecile Ané, David Baum, Tom Givnish and Ken Sytsma for their helpful and constructive feedback, and for always being open for discussion. I thank especially to the

Ken Sytsma and all members of the Sytsma lab whom during my time in Madison made me feel part of their lab (and were my source of coffee during this last year). I also thank Tom Givnish for the collaborative projects we developed and the interesting talks about plants (and for excellent food and movie nights). Thank you to Tom Croat who has been helpful and supportive of Colombian aroiders for many years. Thank you to my office mates Matthew and Giovanny for the good talks, and for his patience. Thank you to Ricardo Kriebel for all the constructive discussion during lunch and during beers. Also, thanks to all the graduate students in the botany department, particularly Rafael, Jeff, Daniel, Richard, Alison, Matthew and Giovanny, for all relaxing Fridays at FAC. Along the same lines, I want to express my gratitude to all the faculty and staff of the department of botany who were always willing to help, particularly to the people in the herbarium, Ted, Mark and to Mo with whom I always had a good conversation.

Finally, I want to thank to all the “non-botany” people in Madison that have shared many good moments with me for the last five years. Especially, I want to thank Isabel and Omar who introduced me to the friendly Chilean community; also to Kaitlin Sherman for her company, and the climbing and hiking (and dinners and drinks); and to all my soccer friends for the good and bad seasons.

TABLE OF CONTENT

Acknowledgments.....	i
Table of content.....	iii
Abstract.....	1
Chapter 1. Rates of diversification and patterns of morphological evolution of the tropical family Araceae at a global scale	
Title page.....	3
Abstract.....	4
Introduction.....	6
Methods.....	11
Results.....	18
Discussion.....	22
References.....	30
Tables and figures.....	41
Supplementary tables and figures.....	74
Chapter 2. Systematics, biogeography and morphological character evolution of the hemi-epiphytic subfamily Monsteroideae (Araceae)	
Title page.....	123
Abstract.....	124
Introduction.....	125
Methods.....	130
Results.....	138

Discussion.....	141
References.....	151
Tables and figures.....	164
Supplementary tables and figures.....	203

Chapter 3. Using genome skimming to resolve the phylogenetic relationships of the

Neotropical genus *Monstera* (Araceae)

Title page	209
Abstract.....	210
Introduction.....	211
Methods.....	214
Results.....	219
Discussion.....	222
References.....	228
Tables and figures.....	238

Chapter 4. Taxonomic studies in the genus *Monstera* (Araceae)

Title page.....	247
Abstract.....	248

Part 1. Two new species of *Monstera* (Araceae) with entire leaves from Panama and Costa Rica

Introduction.....	250
-------------------	-----

Methods.....	251
Taxonomic treatment.....	251
References.....	260
Tables and figures.....	262
Part 2. A preliminary interactive web tool for the identification of the species in the genus <i>Monstera</i> (Araceae)	
Introduction.....	274
Methods.....	276
Preliminary results.....	278
References.....	280
Tables and figures.....	282
Appendix 1. Zuluaga, A., K. Cameron, T. Croat & M. Medecilo. 2015. Testing the monophyly of <i>Spathiphyllum</i> , and the relationship between Asian and tropical American species. <i>Aroideana</i> 38E(1):107-115.....	
	298

ABSTRACT

This research analyzes several aspects of the systematics and evolutionary biology of the family Araceae at different taxonomic and geographic scales. First, we investigated the patterns in the diversification rates for major clades in the family in the context of the order Alismatales. We found evidence for a complex pattern including several different rate regimes within the Araceae decoupled from the background rate. Main shifts towards higher in speciation rates are located in the subfamilies Aroideae, Pothoideae, and Monsteroideae. These rates may be associated with floral characters, pollination interactions, and epiphytism. Second, we studied the subfamily Monsteroideae, and produced a robust phylogenetic hypothesis including all genera and ca. 100 taxa to test the current infrafamilial classification. We found support for the monophyly of three major clades and only seven of twelve genera within Monsteroideae. We investigated the historical biogeography of this group and estimated a Laurasian origin for the subfamily. The continental disjunction between *Monstera* and *Amydrium* + *Epipremnum* is product of a Boreotropical flora, while the other disjunctions (species of *Spathiphyllum* between South America and South East Asia; and *Rhaphidophora* South East Asia and Africa) are most probable product of long distance dispersal. Third, we studied the Neotropical genus *Monstera*. Due to the lack of genomic tools for phylogenetics in tropical aroids, we used a genome skimming approach to recover molecular data from three different DNA regions, the plastid, mitochondrial and nuclear ribosomal DNA. Despite low variation (ca. 1%) in plastome sequences, we recovered a strongly supported phylogenetic hypothesis for *Monstera*, which does not support the current infrageneric classification.

Finally, we focused in the taxonomical aspects of the genus *Monstera*. We used 108 morphological, anatomical and geographical traits from about 4200 herbarium specimens and living plants to produce a web-based interactive tool for the identification of species in the genus. In the process we discovered and described two new species of *Monstera* from Panama and Costa Rica.

Chapter 1

Rates of diversification and patterns of morphological evolution of the tropical family Araceae and their order Alismatales at a global scale

Alejandro Zuluaga¹, Ken Cameron¹

¹ Department of Botany, University of Wisconsin-Madison

ABSTRACT

The family Araceae (order Alismatales) is one of the most diverse monocot families with about 3500 described species, in contrast with the rest of the order, which includes twelve other families and ca. 500 species. The family is especially abundant in the tropics, and exhibits remarkable morphological, anatomical, ecological, and geographical variation. The highest diversity of the subfamily Aroideae (ca. 2000 species) has been hypothesized to be product of an increase in the speciation rates associated with a set of morphological characters (e.g., unisexual flowers, inaperturate and exineless pollen, chemical defense). Despite several recent phylogenetic studies, we still know rather little about the dynamics of lineage and species diversification across the family, or about the underlying drivers of diversification. We explored diversification rates of the major clades in Araceae compared to the rest of the order Alismatales using two datasets: 1) a fossil calibrated phylogeny using Bayesian relaxed molecular clock methods (two genes, ca. 230 taxa); and a supermatrix (ten genes, ca. 950 taxa). We used a Bayesian approach (BAMM), to evaluate the dynamics of the diversification rates in Alismatales. The two-gen dataset showed evidence for a different rate regime located in “True Araceae” clade. The ten-gen dataset revealed a more complex dynamics in rates of diversification with several clades presenting decoupled regimes from the background process. Finally, we analyzed the evolution of the potential characters associated with the evolutionary rates in Araceae. Although high rates in the subfamily Aroideae seems to be associated with traits previously hypothesized, other traits like epiphytism but also geography have play an important role in the diversification of the family.

Keywords

Araceae, Alismatales, BAMM, character evolution, diversification rates, speciation rates

INTRODUCTION

The angiosperm order Alismatales is a monocot clade composed of 13 families and ca. 4000 species (Stevens 2001; Iles et al., 2013) that is sister to the rest of the monocots, excluding Acoraceae (Graham et al., 2006; Iles et al., 2013; Ruhfel et al., 2014; Zeng et al., 2014). Earlier molecular phylogenetic evidence has confirmed the inclusion of Araceae and Tofieldiaceae in the traditional concept of Alismatales (Les et al., 1997; Chase 2004; Graham et al., 2006), and yet the phylogenetic relationships within Alismatales are still controversial (Iles et al., 2013; Les & Tippery 2013; Petersen et al., 2015). Three main clades are usually recovered: Araceae, Tofieldiaceae, and the rest of the Alismatales (“alismatids”), but strong support is lacking for the relationships among these clades. The most common and robust topologies among different studies show Tofieldiaceae as sister to all remaining Alismatales, including Araceae (Iles et al., 2013; Les & Tippery 2013).

Most families of Alismatales are mostly aquatic plants distributed worldwide. Tofieldiaceae with four genera and ca. 20 species is a terrestrial family distributed mainly in the northern hemisphere, though it has also reached high elevations in the Andes, and the Guyana Highlands (Campbell 2008; Azuma & Tobe 2010; Remizowa et al., 2011). The remaining eleven families of Alismatales (excluding aroids) collectively contain 56 genera and about 500 species (Les & Tippery, 2013). These have diversified in both freshwater (e.g. Hydrocharitaceae, Les et al., 1997) and marine environments (e.g. Potamogetonaceae, Zosteraceae, Posidoniaceae; Les et al., 1997) through the entire world.

One family of Alismatales, the Araceae, is significantly more diverse than all the others combined, with around 3,500 described species (or perhaps as many as 4,000; according to Boyce & Croat [2014]). Aroids contain approximately 85% of the diversity in the order, and their diversification has occurred especially in terrestrial, rather than aquatic, environments (Croat, 1988; Mayo et al., 1997). The family is currently divided into eight subfamilies. One of these, subfamily Aroideae, contains ca. 50 genera and 2000 species, more than all the other subfamilies combined. Outside the Aroideae, most of the remaining Araceae diversity is concentrated in one megadiverse genus of ca. 1,000 species, *Anthurium*, which is a member of the subfamily Pothoideae (Croat 1983; 1986; Boyce & Croat; 2014; Carlsen et al., 2014).

The family Araceae is one of the most morphological and ecologically diverse monocot families, especially diverse in the tropics (Mayo et al., 2007). Araceae includes geophytic, terrestrial, hemiepiphytic, epiphytic, rheophytic and also a few aquatic plants, including free-floating species (Croat, 1988; Grayum, 1990; Mayo et al., 2007). Despite several recent molecular phylogenetic studies (Cabrera et al., 2008; Cusimano et al., 2011; Chartier et al. 2014; Henriquez et al., 2014) there are still many problems in generic delimitation and relationships within Araceae. The most recent classification recognizes eight subfamilies, including the former family Lemnaceae, now considered to be Araceae subfamily Lemnoideae (Cusimano et al., 2011). All subfamilies have been shown convincingly to be monophyletic except Aroideae (Cusimano et al., 2011; Chartier et al., 2014; Henriquez et al., 2014). Nevertheless, Aroideae seems to be a

morphologically consistent group defined mainly by the presence of unisexual aperiogonated flowers, and inaperturate pollen with reduced exine (ektexines). The main issues concerning phylogenetic reconstruction in Aroideae relate to the backbone of the clade, and exact the position of certain problematic genera such as *Montrichardia*, *Calloopsis*, *Anubias* *Zantedeschia*, and *Calla* (the latter of which curiously has bisexual flowers), (Cusimano et al., 2011; Chartier et al., 2014; Henríquez et al., 2014). The monotypic genus *Calla* is distributed in the temperate zone of United States and has been consistently include in the subfamily Aroideae in all molecular analysis using plastid data, despite the fact that morphologically it does not resembles any members of this subfamily. Chartier et al. (2014) using the nuclear gene PHYC, recovered *Calla* as sister to the rest of the Aroideae, so the position of this genus should still be considered ambiguous.

Perhaps the most distinguishable feature of Araceae is its inflorescence organization. Unlike many other groups in Alismatales, Araceae have very reduced flowers, organized in compact inflorescences, that are specialized for a variety of pollination strategies, including food rewards, mating mutualism, ovipositing mutualism, and deception (Gibernau et al., 2010; Bröderbauer et al., 2012; Chartier, 2011). Trapping inflorescences, which usually involve deceptive pollination, have evolved multiple times in at least 27 genera, mainly in the subfamily Aroideae (Bröderbauer et al., 2012). Several patterns have been described regarding the evolution of aroid flowers, and pollination evolution (e.g. Chartier et al., 2014; Cusimano et al., 2011). Two main types of inflorescences have been described, hermaphroditic inflorescences with bisexual

flowers, present in most of the subfamilies (Orontioideae, Lemnaceae, Pothoideae, Monsteroideae and Lasiodeae); and monoecious inflorescences with unisexual flowers, which are present in the *Stylochaeton* clade and Aroideae (Mayo et al., 1997; Cusimano et al., 2011). The role of inflorescence and flower structure, together with plant-pollinator relationships, as a driver of species diversification in the family has not been explored quantitatively, but hypotheses have been proposed for how inflorescence structure and pollination might have played a role in affecting diversification rates.

Hesse (2006a; 2006b) hypothesized that the evolution of the monoecious inflorescence type, together with other traits are responsible for an increased rate of speciation in Aroideae. The other characters Hesse argued are part of the key innovation model are: 1) switch from perigonated to aperigonated flowers; 2) from physical (calcium crystals, sclereids) to chemical defense (laticiferous, biferines) against herbivory (Hay & Mabberley, 1991); 3) from aperturated to omniaperturated (inaperturated) pollen; and 4) from developed to reduced exines with a polysaccharide outer exine layer instead of sporopollenin. Pollen with reduced exine (exineless pollen) has been described from other monocot groups like Zingiberales (Kress et al., 1978), and other Alismatales like Cymodoceaceae or Hydrocharitaceae (Chanda et al. 1988; Tanaka et al., 2004; Furness & Banks, 2010). This particular feature seems to be associated to pollination in humid and wet environments, and at least for Alismatales the presence of exineless pollen is always correlated with the presence of omniaperturate pollen (Furness & Banks, 2010).

Hesse (2006a; 2006b) further argued that all these traits are “connected by causal relationships”, and may followed on from the loss of bisexual flowers in Aroideae by the early Tertiary. His hypothesis was based on the idea that that these changes are associated (as a pre-requisite or a result) with the pollination behavior observed in most of the species of the clade. Species of the Aroideae (as for most of Araceae) live in humid and wet environments (Mayo et al., 1997), and have a short time span in which to achieve successful pollination, thereby leading to: 1) pollen with short viability, 2) rapid germination of pollen tubes; and 3) brief periods of stigma receptivity (Hesse, 2006a; 2006b; Barabe et al., 2008). Other authors, in contrast, have proposed that traits like epiphytism and pollinator interactions could have contributed to some of the diversity found in Aracaceae, especially outside of subfamily Aroideae. For example, Madison (1978) suggested that the large number of species in *Anthurium* might be the result of two adaptive radiations, the first one related to the diversification of pollination interactions with bees and flies, and the second related to the epiphytism. That paper, like so many others, was written without the presence of a robust and objective phylogenetic reconstruction for the family.

The objective of this study is to investigate the patterns in the diversification rates within the entire order Alismatales, particularly in the family Araceae. We are also interested in evaluating the effect of taxon sampling in documenting rate shifts by using data sets of differing size. Additionally, we investigated the evolution of some morphological characters that may be associated with increases or decreases in rates of diversification rates.

METHODS

DNA sampling and data matrix assembly. We built two molecular datasets in order to perform analyses for this study. The first one included 229 species representing all families and 56 genera in the order Alismatales, and seven outgroups representing Petrosaviales, Acoraceae, Amborellales, Ceratophyllales, Chloranthales and Austrobailetyales. This matrix is composed of two molecular markers, *matK* and *rbcL*, mostly based on data from Cabrera et al., (2008) for Araceae, and Les et al. (1997) for Alismataceae (but see Supplementary **Table S1**, for a complete list of GENBANK accession numbers). The second dataset is a supermatrix that includes 946 species sampled across Alismatales and 10 sparsely sampled molecular markers obtained from GenBank (**Table S1**), but also from our extensive studies in the Araceae subfamily Monsteroideae (CHAPTER 2).

Sequences were downloaded and organized using Geneious 8.0 (Biomatters, <http://www.geneious.com>, Kearse et al., 2012). Each gene was aligned individually using an initial alignment was performed in MAFFT v7 (Katoh et al., 2002, Katoh, 2013) followed by two subsequent refinements using Muscle v3.8.31 (Edgar 2004). All alignments were visually inspected and manually adjusted in Geneious 8.0 using protein translation for the coding regions. For the two-gene dataset we used PartitionFinder (Lanfear et al., 2012) to select the best-fit partitioning schemes and models. However, because not all the models are available in RaxML (Statamakis, 2006) or BEAST v1.8 (Drummond & Rambaut, 2007, Drummond et al., 2012) we performed a second run

restricting PartitionFinder to only the models RaxML and BEAST could apply. For the initial data blocks in PartitionFinder we allocated nucleotides in protein-coding genes (*rbcL*, *matK*) according to whether they belonged to the first, second or third codon position, and all the other markers as independent data blocks. The PartitionFinder analysis resulted in five partitions for the two-gene dataset; the models and partition schemes are shown in **Table 1**. Since RaxML does not allow for different models, and because our 11-gene dataset has a great amount of missing data, we assigned every molecular marker as a different partition.

We performed two maximum likelihood (ML) searches for each dataset using RAxMLv8.4.2 (Stamatakis, 2014) as implemented in CIPRES (Millet et al., 2010). The first ML search was performed with all data unpartitioned, and the second one was performed using the data partitioning schemes from PartitionFinder. We inferred the maximum likelihood trees using 100 independent searches for the two-gen and 20 for the ten-gene datasets. Bootstrap support values were obtained by 1000 independent throughout searches instead of using the fast bootstrap option in RaxML. For our 10-gene supermatrix analysis we also ran a constrained analysis using clades recovered and strongly supported in previously published studies (Cusimano et al. 2011; Les & Tippery, 2103; and Henriquez et al., 2014; Ross et al., Unpublished) to reflect the most currently understood phylogenetic relationships in Alismatales. Briefly, we defined our “major” clades in Alismatales and constrained the tree so these clades are monophyletic. The clades are and constraints are: Tofieldiaceae as sister to the rest of Alismatales; the eight aroid subfamilies (Gysmonstachyoideae, Orontioideae, Lemnoideae, Pothoideae,

Monsteroideae, Lasiodieae, Zamioculcoideae and Aroideae); and two clades in alismatids (tepaloïd and petaloïd clades).

Molecular dating. To estimate ages of nodes, we used the two-gene dataset for Bayesian molecular dating using BEAST 1.8 (Drummond et al., 2012), and employed eight calibrations points following recommendations of Sytsma *et al.*, (2013) for the priors on the calibrations points. The two genes were partitioned as suggested by PartitionFinder. We performed a preliminary analysis using different prior combinations to define the best parameters for our dataset. Evaluation of the ucl.d.stdev and coefficient of variations under uncorrelated lognormal relaxed clock showed that our data do not fit a strict molecular clock. We also evaluated the differences between choosing a pure birth (Yule) prior versus a Birth-Death prior for the tree. We performed three analyses; in the first analysis the topology was not constrained (except by the fossil calibrations); secondly, we constrained the position of Tofieldiaceae as sister to the rest of Alismatales; and finally we constrained the entire tree to reflect the most updated phylogenetic relationships in Alismatales. In all analysis we used a pure birth (Yule) prior for the tree, and rate variation across branches (clock prior) as uncorrelated and log-normally distributed. For all analysis we used lognormal priors in the calibrations points with mean=1 and SD 0.75, except for the root where we used a conservative uniform prior (see fossil calibrations section). For every analysis we ran two independent chains of 75 000 000 generations, sampling every 10 000 generations in BEAST v1.8 as implemented in CIPRES, or more until they reached an effective sample sizes (ESSs) >200. Results from BEAST were checked for convergence and ESSs on Tracer, version

1.6 (<http://beast.bio.ed.ac.uk/Tracer>). We concatenated the trees using LogCombiner version 1.7.2 (<http://beast.bio.ed.ac.uk/LogCombiner>) using a 20% burn-in fraction, and we resampled the posterior distribution to obtain a final sample of 10000 trees. The final chronogram was constructed using the maximum clade credibility tree with mean node heights in TreeAnnotator version 1.7.2 (<http://beast.bio.ed.ac.uk/TreeAnnotator>).

Fossil Calibrations. We evaluated all the reliable fossils available for Alismatales based on previous studies (Nauheimer et al., 2012, Iles et al., 2015), and chose nine that covered a wide range of clades and were assignable with some certainty to a clade in Alismatales (**Table 2**). For the root we used a uniform prior (120-160 Ma) comprising some of the youngest and oldest ages estimates for Angiosperm (e.g. Bell et al., 2010; Magallón & Castillo, 2009;). For the crown node of monocots we used a constraint similar to Nauheimer et al. (2012) but a lognormal distribution so the median agrees with the age of the fossil pollen *Liliacidites* sp. A (Doyle & Hickey, 1976). Within the Alismatids we chose the oldest known *Stratiotes* fossil from the top of the Paleocene (56.0 Ma) (Sille et al., 2006) to constrain the minimum age of the crown clade of Hydrocharitaceae. The fossil of *Aponogeton harryi* from the Late Cretaceous (Lower Campanian, 82-81Ma) in United States (Hicks, 1993) was used to calibrate the stem node of Aponogetonaceae (= crown node alismatids).

For Araceae, we assigned the fossil of *Spixiarum kipea* (Coiffard et al., 2013) from the upper Cretaceous (113 Ma) in Brazil to the stem node of Orontioideae (Proto-Araceae). Friis et al., (2010) presented two fossils named “Araceae fossil sp. A” and

“Araceae fossil sp. B”, from the early Cretaceous (110 Ma) in Portugal. They argued that it represents a taxon with bisexual and unisexual flowers respectively. As a conservative approach we assigned both of these fossils to the crown node of a clade comprising Lemnoideae and the rest of Araceae (sensu Cusimano *et al.*, 2011). We assigned *Petrocardium cerrejoense* from the Mid to Late Paleocene (56-61.6 Ma) in Colombia (Herrera *et al.*, 2008) for the crown node of Pothoideae. The fossil pollen *Lasioideaacidites* from the Late Cretaceous (70-75 Ma) in Siberia (Hofmann & Zetter, 2010) represents the subfamily Lasioideae. We used this fossil to calibrate the stem node of Lasiodeae (Podolasia clade [*sensu* Cusimano *et al.*, 2011]). For the stem node of the *Heteropsis* + *Rhaphidophora* clades (in the subfamily Monsteroideae) we assigned fossil *Rhodospathodendron tomlinsonii*, from the Late Cretaceous (66 Ma) (Bonde, 2000). For the stem node of the *Typhonodorum* clade (sensu Cusimano *et al.*, [2011]) we used *Nitophyllites zaisanicus* from the Palaeocene (55.8 Ma) in Kazakhstan (Iljinskaja, 1986; Wilde *et al.*, 2005).

Molecular dating of the supermatix. Due to the size of the supermatrix a Bayesian molecular dating approach was not possible. Instead we used penalized likelihood (PL, Sanderson, 2002) as implemented in the function ‘chronos’ from the package APE (Paradis *et al.*, 2003) in R (R Development Core Team, 2008). Because our topology differences between two and ten-gene, and constrained and unconstrained analysis are not at the major clades but at the generic level -were not strong support has been found in any study-, we constrained the ten-gene dataset to our two-gene topology from BEAST in all subsequent analyses. As an input tree for PL we used the maximum likelihood tree derived from the constrained analysis described above. As calibration

points we used the age estimates from our two-gene BEAST analysis for each subfamily in Araceae, and each family in the rest of Alismatales. For lambda we tested values of 0.1, 0.25, and 0.5 to see which values will give us the closest result to our BEAST analysis.

Diversification Analysis. To investigate shifts in diversification rate through time and among lineages we performed analyses using the software BAMM v2.1 (Rabosky, 2014). When taxon sampling is incomplete and non-random BAMM requires a file in which every tip is assigned to a clade and the number of species that it represents is given. Numbers of described and currently accepted species for each genus of Araceae were obtained from Boyce & Croat (2014), and we used numbers from Les & Tippery (2013) for Alismataceae. Additional information was obtained from the *World Checklist of Selected Plant Families* (WCSP, Royal Botanic Gardens, Kew). Because of some issues concerning taxonomic uncertainty and higher-level classification within the family Araceae we assigned ambiguous genera to unambiguous clades strongly supported in our molecular analysis or in previous studies. We used two different clade schemes, the first scheme (Clades1) used recognized subfamilies within Araceae, but only considered families for the rest of Alismatales; the second scheme (Clades2) used all of the highly supported clades from Cusimano et al. (2011) for Araceae, but again only families for the rest of Alismatales. We performed two different analyses in BAMM for each of the two datasets and the two sets of clade designations. The first analysis used a MEDUSA-like approach implemented in BAMM, the second analysis used the complete BAMM model. We performed preliminary runs with poissonRatePrior of 1, 0.5 and 0.1 respectively, and

allowing shifts to occur on all branches (`minCladeSizeForShift = 1`). For each analysis we completed 50 million generations of Markov Chain Monte Carlo (MCMC) runs with `poissonRatePrior` of 1 for the two-gene dataset, and 100 million and `poissonRatePrior` of 0.5 for the ten-gene dataset, sampling parameters every 10000 generations. Results were summarized using the packages Coda (Plummer et al., 2006) and BammTools (Rabosky et al., 2014) in R. We assessed convergence of BAMM runs by computing effective sample sizes of log-likelihoods, numbers of processes, and evolutionary rate parameters using CODA. For every analysis we evaluated the 95% credibility set of shifts configurations, the best shift configuration and used Bayes Factors to evaluate rate heterogeneity and significant shifts in our dataset.

Evolution of characters potentially associated with diversification. Flower characters. We recorded data associated with flower traits proposed by others as potential key innovations and possible drivers of diversification in the subfamily Aroideae. The traits considered were: flower sexuality, presence of perianth, pollen aperture, exine presence, pollen size and shape, and pollen starch content. Data from aroid pollen was obtained mainly from Grayum (1990) for Araceae, but from many other sources for Alismatales (e.g. Chanda et al. 1988; Kubitzki, 1998; Tanaka et al., 2004; Furness & Banks, 2010). For pollen size we used the longest and shortest distances in polar view, and the diameter for spherical pollen shapes. Because there are many factors affecting pollen size (e.g. hydration, folding, angle of the photograph) we used mean value reported in the literature or calculated mean values per genera after recording as many species has possible. Finally, the flower traits associated with Aroideae were tested

for correlated evolution using the discrete module from BayesTraits (Pagel, 1994; Pagel & Maede, 2006). We tested flower sexuality against, perianth presence, pollen aperture, and exine reduction. We used the MCMC method with 50 million generations, and likelihood ratio for model testing. We also tested contingent evolution in these traits as described in (Pagel, 1994).

Ecology and Life form. Some superficial traits describing ecological and habitat preferences were also coded for each genus. Climatic region was coded as either tropical or temperate; seasonality was coded as evergreen or seasonally dormant (we included annual plants from Alismatales in this category); life form was coded in several ways based on categories defined by Croat (1988) and Mayo et al. (1997). First we coded the habitat as aquatic free-floating, aquatic rooted (including helophytes and submersed aquatics), terrestrial (including geophytes and lithophytes), or epiphytes (including hemiepiphytes). Secondly, we coded the growth form (shoot architecture) as either acaulescent, caulescent (and terrestrial), rhizomatous, tuberous, or climbing stems. Discrete characters were reconstructed using a stochastic mapping Bollback (2006) and the rerooting method as implemented in the R package phytools (Revell, 2012). Continuous characters (size) were reconstructed using ML approach as implemented in the function contMap from R package phytools (Revell, 2012).

RESULTS

Maximum likelihood analyses. The two-gene and ten-gene alignments contained 3150 bp (19% missing data) and 10343 bp (76% missing data) respectively, and the

maximum likelihood analyses from partitioned and unpartitioned datasets recovered the same topology with only slight differences in branch support (**Figures 1 & 2**).

Topologies from the two different datasets did not differ for the major clades in Alismatales (**Figures 1 & 2**), but did show differences in the generic relationships within certain clades. The main differences among our ML topologies and Cusimano et al. (2011) were the position of the aroid genus *Montrichardia*, the aroid tribe Spathiphyllaeae, and the exact intergeneric relationships within the clades Lasioideae, Spathicarpeae, Schismatoglottideae and Caladieae, although no strong support was found for these alternative relationships in the previously published trees or in our new analyses. We recovered Tofieldiaceae as sister to the rest of the Alismatales in the two-gene ML but bootstrap support was low. Within the alismatids, all families as proposed by Les & Tippery (2013) and Iles et al. 2013 were recovered as monophyletic, but family relationships did not agree completely with those of Iles et al. (2013), particularly with regards to the positions of Aponogetonaceae, Scheuzeriaceae, Posidoniaceae, Cymodoceaceae, and Ruppiaceae.

Molecular dating. We based our results on the BEAST analysis with the Tofieldiaceae constraint, but ages for major clades from the three analyses and the PL chronogram are shown in **Table 3**. Molecular clock analysis showed our data does not fit a strict molecular clock, so we used the lognormal relaxed clock. The BEAST analysis without any topology constraint (except by fossil calibrations) did not recover Tofieldiaceae sister to the rest of the Alismatales; however age estimation from the major clades did not differ greatly from the constrained analyses (**Table 3**). The age for the

crown of Alismatales is 129.6 Ma (CI 122.8-136.5) with a stem node age of 134.3 Ma (CI 127.6-141.7). Tofieldiaceae, alismatids and Araceae crown ages are 73.7, 113.6, and 121.2 Ma respectively. All stem nodes for the aroid subfamilies are dated well into the Cretaceous, whereas crown nodes for these clades are closer to the K/T boundary except for Orontioideae (52 Ma), and Lasioideae (24.1 Ma) (**Table 3**). Our penalized likelihood approach agreed with ages of the BEAST analysis for major clades and subfamilies in Araceae, but it was less accurate for clades far from the root of the tree (e.g., clades Thomsonieae and Caladieae, **Table 3**).

Diversification. *Two-gene dataset.* All the BAMM analyses using the two-gene dataset produced very similar results regarding the speciation, extinction and diversification rates, and the best shift configuration, so we will focus our discussion on the BAMM model with the clade set 1 (**Table 4 & 5, Figure 3**). The different models and clade sets produced different number of shift configurations in the 95 % credibility interval; however, one shift configuration with a single shift in diversification rates within Alismatales was recovered consistently through all the analyses (**Figure 3**). This shift in diversification rates is located along the stem of the “True Araceae” clade with probabilities between 0.45-0.8 and Bayes factors >17000. The crown Araceae showed mean speciation rates between 0.46-0.577, whereas the crowns of alismatids and Tofieldiaceae showed rates of 0.053-0.055 and 0.054-0.056 respectively (**Tables 4 & 5**).

Ten-gene dataset. As expected BAMM analyses from our ten-gene analysis produced a high number (454-677) of shift configurations in the 95 % credibility set, and

there is not a best shift configuration with significant highest probability (**Table 4**). Overall, mean speciation, extinction, and diversification rates are similar among all analyses with the ten-gene dataset so we will focus our discussion in the BAMM model with the clade set 1 (**Table 4**). The two clade sets analyses found slightly different number and location of core shifts (**Figures 5 & 6**). Different clade sets share the same shift configuration regardless of the model used (MEDUSA-like or BAMM). Taking into account the results from the two different clade sets we observed a total of ten shifts (named a, b, and 1-8, **Figures 5 & 6**). Shifts a and b are not discussed since they reflect a shift in diversification rates between outgroup (Acoraceae) and the ingroup. The other eight shifts are associated with the following clades: 1) Pothoideae + Monsteroideae (Bisexual climbers clade); 2) Aroideae (excluding *Calloopsis*); 3) *Pistia* clade + *Protarum*; 4) and 5) *Anthurium*; 6) Lasioideae; 7) and 8) *Philodendron* clade. The *Philodendron* clade showed the highest net diversification rate (0.242-0.291) followed by Pothoideae (0-0.196-0.198) and Monsteroideae (0.139) (**Table 6 & Figure 7**).

Character evolution. Analyses using the rerooting method and stochastic mapping produced very similar ancestral state reconstructions. All characters associated with the subfamily Aroideae have evolved more than once in Alismatales (**Figures 8, 9 & 12**). Unisexual flowers, absence of perianth, omniaperturate pollen, and reduced outer exine layer have evolved together in Aroideae and a clade of Alismatales comprising the families Cymodoceaceae and Posidoniaceae (**Figure 8 & 9**). The ancestral states estimated for these characters are: outer exine layer present, pollen monosulcate, and hermaphroditic flowers with perianth. Analysis of correlated evolution showed that

pollen aperture and exine reduction show patterns of correlated evolution with flower sexuality, whereas perianth evolves independently in Alismatales (**Table 7**). Pollen size did not show a lot of variation except for the long and narrow filamentous pollen from the families Zosteraceae and Cymodoceaceae (**Figure 12**). All characters associated with ecology and life form showed a complex pattern of evolution with homoplasy (**Figure 10 & 11**). As a general pattern, we found that habit, growth form, and seasonality have evolved multiple times in Aroideae, whereas in all the other subfamilies they have evolved only once (with some exceptions, such as growth form in Lasioideae). The ancestral estimated states for these characters are: tropical regions, evergreen plants, terrestrial and rhizomatous.

DISCUSSION

Phylogenetic analyses and molecular dating. The recent generation of expanded genomic tools for the Araceae has opened the possibility of developing larger molecular datasets for generating better supported hypotheses about genera and species relationships among aroids. Accurate phylogenetic reconstructions are critical for investigating historical biogeography and character evolution, and we feel confident that the current state of phylogenetic reconstruction in the order Alismatales presented here is sufficient for investigating questions about diversification in major clades in this group. The only obvious differences among previously published phylogenetic reconstructions (e.g. Cusimano et al. 2011, Iles et al. 2013) and our two and ten-gene datasets involved nodes that are only weakly resolved, mainly involving relationships of Alismatales families and among genera of Araceae. There are no examples of conflict or hard

incongruence detected with previous analyses. All genera that included more than one species were recovered monophyletic for alismatids, and again, main taxa with uncertainty in Araceae are *Montrichardia*, *Calla*, *Calloopsis*, and *Anubias*.

Our age estimates for Alismatales (129.56 Ma) agree with recent studies using different methods and fossil calibrations (Magallón & Castillo, 2009; Anderson & Janssen, 2009; Hertweck et al., 2015; Magallon et al., 2015,). Despite using older and different fossil calibrations, our estimates for subfamilies in Araceae are very similar to Nauheimer et al. (2012). The main differences for the major clades are ages estimated for Monsteroideae (56.3 Ma vs. 68.8 in our analysis), and clades within this subfamily, which are older than previously estimated. This is mainly due to differences in the relationships among clades of Monsteroideae and the inclusion of an older fossil to calibrate the stem of Monsteroideae. Recent fossil evidence has shown that at least some groups of Araceae were widespread by the early Cretaceous (Coiffard et al., 2013, Gallego et al., 2014), and that all the subfamilies were present by the late Cretaceous. However, some of the older fossils resembling Pothoideae or Monsteroideae are difficult to interpret or have not been studied in detailed (Friis et al. 2010a; Friis et al., 2010b). The main factor affecting the dating of our ten-gene phylogeny was the number of calibrations points and not the smoothing parameter (λ) used during the penalized likelihood calculation. We obtained ages similar to our BEAST tree for the major clades, but differences increased with the distance from a particular node to a calibration point.

Diversification and character evolution in Alismatales. Recent studies of diversification rates among monocots did not find any increase in the order Alismatales, but some evidence for a decrease was found in the alismatid family Butomaceae (Hertweck et al., 2015). In contrast, our findings reveal a complex pattern of ecological and morphological evolution, and changes in diversification rates throughout the order Alismatales, particularly in the family Araceae. The reason for these differences is that diversification analysis methods such as BAMM or MEDUSA will depend on the scale of the phylogenetic tree, as we showed with the different results when comparing our two vs. ten-gene dataset. In our study the Tofieldiaceae and alismatids showed both low rates of diversification and decrease in rates through time. The aquatic subfamily Lemnoideae, the Proto-Araceae, and the *Stylochaeton* clade showed the same pattern. The first two contain aquatic plants or helophytes, which suggest that adaptation to aquatic environments could have a direct negative effect on diversification rates in this order. The *Stylochaeton* clade is an interesting group, it has flower traits intermediate between the subfamily Aroideae and the rest of Araceae (having unisexual flowers with a reduced perianth), and it is the only clade restricted to Tropical Africa with only 32 species. Hesse (2006a) suggested that this group represents a “dead end” in the evolution of Araceae, because it has acquired the unisexual flowers but none of the other traits that have made Aroideae “successful”. The slowing and decreasing speciation rate in the group supports his idea.

Taxon sampling and different clade sets. As expected analyses of two vs. ten-gene datasets produce very different shift configurations due to the differences in taxon

sampling (229 vs. 946 taxa). We tested impact of allocating species numbers to different set of clades when using BAMM for incomplete sample phylogenies. We found that different clade sets produced different core shifts configurations; however in our case the shifts were located in the same major clades. Despite that clade allocation had an impact in the distribution of shifts this was not an issue for our discussion of the rates diversification in Alismatales.

Shifts in diversification. Diversification analysis of the two-gene dataset showed strong evidence for a shift in the “true Araceae” clade. We did not find evidence of an increase in rates in the subfamily Arodieae, in fact, when looking at the rates of individual clades, Aroideae and ‘Bisexual Climbers’ shared similar rates. Initially, this increase seems be associated with an adaption to terrestrial environments in the family Araceae as we see that the two sister clades to the True Araceae (Orontioideae and Lemnoideae), which have low rates of diversification, are helophytes or free-floating aquatics. However, analysis of the ten-gene dataset allowed to dissect the pattern of diversification within Araceae further, and showed that at least six clades have undergone shifts in rates of diversification.

Lasioideae (shift 6). This aroid subfamily is a small clade with only nine genera and ca. 60 species with a pantropical distribution (Mayo et al., 1997; Boyce & Croat, 2014). Members of this group are terrestrial plants or helophytes, with tuberous or rhizomatous growth form, and it is the only clade with bisexual flowers that has evolved insect trap inflorescences (Bröderbauer et al., 2012). Despite an overall increase in the net

diversification of this clade, we also detected a high rate of extinction (also reflected in the long stem branch leading to the crown of the group). The fossil record and molecular dating analyses have supported the hypothesis of a boreotropical origin for the current distribution of this clade. This may help explain the pattern on diversification rates with high extinction during the dispersal from the northern hemisphere to the tropics, and recent increase in the diversification rates. However, this seemingly contradictory pattern is intriguing and suggests that Lasioideae may be an ideal clade for future, focused investigation into the evolutionary process within Araceae.

Pothoideae + Monsteroideae (shifts 1, 4 and 5). At least two shifts showing increases in diversification rates were found in this clade of sister subfamilies, one in the stem of the clade, and the second one in the genus *Anthurium* of Pothoideae, with different locations in the clade set 1 vs. clade set 2 analysis (shifts 4 and 5 in **figures 5 & 6**). With ca. 1200 species, this clade represents most of the diversity of Araceae in the Neotropics, while only ca. 300 species from the group are present in the Paleotropics. Additionally, ca. 80% of the species are epiphytes or hemiepiphytes, whereas other non-epiphytic growth forms are restricted to the genera *Spathiphyllum*, *Stenospermation*, and *Anthurium*. Madison (1978) hypothesized that the high species diversity in *Anthurium* is a product of two adaptive radiations (related to epiphytism, and bee and fly pollination, respectively). Our analyses show that, in fact, there is an increase in speciation and net diversification in *Anthurium*, but we do not have evidence to support the idea of an adaptive radiation. Givnish (1997; 2015 In Press) has discussed the problems in the misuse of the terms adaptive radiation and explosive speciation. Based on his definition

and our observations we believe that Madison was suggesting not a true radiation following adaptation, but rather an explosive speciation event instead. Although epiphytism and pollination relationships may be factors contributing significantly to the high speciation rate in *Anthurium*, we do not consider that there is a wide range of ecological diversity that will indicate the presence of an adaptive radiation in this genus; however, we did not explore these questions directly in this study. It is worthy of further consideration.

Aroideae (shifts 2,3,7,and 8). At least three shifts were found in this subfamily, the first one at the stem node of the clade (excluding *Calloopsis*); the second one in the *Philodendron* clade (represent by shifts 7 and 8 in the different clade set analyses); and the third one in the *Pistia* clade + *Protarum*. Despite the lack of support and the uncertainty in the position of some taxa, Aroideae seems to be a clearly defined group by morphological characters (e.g., bisexual flowers lacking perianth, omniapeturate pollen, absence of sporopollenin in the exine) (Cusimano et al., 2011). At least the unisexual flowers, omniaperturate pollen and exine reduction showed a pattern of contingent evolution in Alismatales (**Table 7**). All these characters together have evolved independently at least one other time in Alismatales, but their combination, along with other vegetative characters (e.g., presence of biferines and laticifers), and specialized pollination strategy (spathe behavior and short pollination window) are unique in the order. Although it is difficult to determine the causal relationships between these characters and the pollination strategy (Hesse 2006a; 2006b), we agree with the hypothesis that these represent key innovations that led to an adaptive radiation in

Aroideae, and that in this particular case the radiation is also associated with an increase in net diversification rate. Aroideae with ca. 2000 species shows remarkable morphological and ecological diversity. It is the only subfamily with an almost cosmopolitan distribution (Boyce & Croat 2014), growing in very diverse habitats. Pollination interactions are strikingly diverse, including many types of deception and insect-trap inflorescences (Diaz & Kite, 2006; Bröderbauer et al., 2012, Chartier et al., 2014). All the growth forms and habits, including submersed and free-floating aquatic plants, are represented, and have evolved in a complex manner (i.e., independent origins for the same character, **Figure 11**). Previously, Cusimano et al. (2011a) reconstructed 81 morphological characters for Araceae and found a large amount of homoplasy, suggesting repeated adaptive shifts.

Potential drivers of diversification in Araceae. Although we did not test for any character to be directly associated with increases in speciation rates we can describe some patterns in Araceae based on our diversification and character evolution analyses. Epiphytism (especially hemiepiphytism) is the most common habit in the three most diverse clades of Araceae (*Anthurium* [ca. 80%], *Philodendron* [ca. 80%], and *Monsteroideae* [ca. 90%]), which also showed increases in diversification rates. Epiphytism has found to be related to increases in diversification in other monocot families like Bromeliaceae and Orchidaceae (Gravendeel et al., 2004; Givnish 2010; Givnish et al., 2014; Givnish et al. Unpublished). However, outside the Neotropics epiphytism is not the prevalent habit, and sister clades to these epiphytic groups present complete different patterns except by Monsteroideae. *Pothos* (plus *Pedicularum* and

Pothoidium), the sister clade to *Anthurium* has only ca. 60 species of scandent hemiepiphytes distributed across South East Asia. *Homalomena* and *Adelonema* (sister to *Philodendron*) are neither epiphytic (rheophytic is the most common habitat) or as diverse (ca. 75 species). This implies that Araceae inhabiting different tropical areas seem to be affected by different abiotic (e.g., geographical and topographical complexity) and biotic factors (e.g., pollination interactions [Chartier et al., 2014]). Further research should be focus on improving the phylogenetic hypothesis for Araceae and disentangling the factors affecting the complex pattern of speciation and diversification rates in the family.

REFERENCES

- Anderson, C.L., Janssen T., 2009. Monocots. In: *Timetree of life*, eds. Hedges S.B., Kumar S., New York: Oxford University Press, New York, pp. 203–212.
- Azuma, H., Tobe, H., 2011. Molecular phylogenetic analyses of Tofieldiaceae (Alismatales): family circumscription and intergeneric relationships. *J. Plant. Res.* 124, 349–357. doi:10.1007/s10265-010-0387-5.
- Barabé, D., Lavallée, K., Gibernau, M., 2008. Pollen viability and germination in some neotropical aroids. *Botany* 86, 98–102. doi:10.1139/B07-110.
- Bell C.D., Soltis D.E., Soltis P., 2010. The age and diversification of the angiosperms revisited. *Am. J. Bot.* 97, 1296–1303. doi: 10.3732/ajb.0900346.
- Bollback, J. P. (2006) Stochastic character mapping of discrete traits on phylogenies. *BMC Bioinformatics*, 7, 88.
- Bonde, S.D., 2000. *Rhodospathodendron tomlinsonii* gen. et sp. nov., an araceous viny axis from the Nawargaon intertrappean beds of India. *Palaeobotanist* 49, 85–92.
- Boyce, P.C., Croat, T.,B., 2014. The Überlist of Araceae: totals for published and estimated number of species in aroid genera. Available via <http://www.aroid.org/genera/140601uberlist.pdf>. Accessed 5 May 2015
- Bröderbauer, D., Diaz, A., Weber, A., 2012. Reconstructing the origin and elaboration of insect-trapping inflorescences in the Araceae. *Am. J. Bot.* 99, 1666–1679. doi:10.3732/ajb.1200274.
- Cabrera, L.I., Salazar, G.A., Chase, M.W., Mayo, S.J., Bogner, J., Dávila, P., 2008. Phylogenetic relationships of aroids and duckweeds (Araceae) inferred from coding and noncoding plastid DNA. *Am. J. Bot.* 95, 1153–1165.

- Campbell, L.M., 2008. Tofieldiaceae. In: *Nuevo Catálogo de la Flora Vascular de Venezuela*, eds., Hokche, O., Berry, P.E., Huber, O., Fundación Instituto Botánico de Venezuela & CONICIT, Caracas, pp. 823.
- Chanda, S., Nilsson, S., Blackmore, S., 1988. Phylogenetic trends in the Alismatales with reference to pollen grains. *Grana* 27, 257–272. doi:10.1080/00173138809429948.
- Chartier, M., Pélozuelo, L., Gibernau, M., 2011. Do floral odour profiles geographically vary with the degree of specificity for pollinators? -Investigation in two sapromyophilous *Arum* species (Araceae).
- Chartier, M., Gibernau, M., Renner, S.S., 2014. The evolution of pollinator-plant interaction types in the Araceae. *Evolution* 68, 1533–1543.
- Chase, M.W., 2004. Monocot relationships: An overview. *Amer. J. Bot.* 91, 1645–1655.
- Coiffard, C., Mohr, B.A.R., Bernardes-de-Oliveira, M.E.C., 2013. The Early Cretaceous Aroid, *Spixiarum kipea* gen. et sp. nov., and implications on early dispersal and ecology of basal monocots. *Taxon* 62, 997–1008. doi:10.12705/625.21.
- Croat, T., 1983. A revision of the genus *Anthurium* (Araceae) of Mexico and Central America. Part 1: Mexico and Middle America. *Ann. Missouri Bot. Gard.* 70: 211–417.
- Croat, T., 1986. A revision of the genus *Anthurium* (Araceae) of Mexico and Central America. Part 2: Panama. *Monogr. Syst. Bot. Missouri Bot. Gard.* 14: 1–204.
- Croat, T., 1988. The ecology and life forms of Araceae. *Aroideana* 11(3–4): 4–56.
- Cusimano, N., Bogner, J., Mayo, S. J., Boyce, P.C., Wong, S.Y., Hesse, M., Hetterscheid, W.L.A., Keating, R.C., French, J.C., 2011. Relationships within the Araceae:

- comparison of morphological patterns with molecular phylogenies. *Am. J. of Bot.* 98, 654–668.
- Diaz, A., Kite, G.C., 2006. Why be a rewarding trap? The evolution of floral rewards in *Arum* (Araceae), a genus characterized by saprophilous pollination systems. *Biol. J. Linn. Soc.* 88, 257–268. doi:10.1111/j.1095-8312.2006.00612.x.
- Doyle, J.A., Hickey, L.J., 1976. Pollen and leaves from the mid-Cretaceous Potomac Group and their bearing on early angiosperm evolution. In: *Origin and early evolution of angiosperms*, ed., Beck C.B., New York: Columbia University Press, New York, pp. 139–206.
- Drummond, A.J., Rambaut, A., 2007. BEAST: Bayesian evolutionary analysis by sampling trees. *BMC Evol. Biol.* 7, 214. doi:10.1186/1471-2148-7-214.
- Drummond, A.J., Suchard, M.A., Xie, D., Rambaut, A., 2012. Bayesian phylogenetics with BEAUti and the BEAST 1.7. *Mol. Biol. Evol.* 29, 1969–1973. doi:10.1093/molbev/mss075.
- Edgar, R.C., 2004. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res.* 32, 1792–1797.
- Farrell, B.D., Dussourd, D.E., Mitter, C., 1991. Escalation of plant defense: Do latex and resin canals spur plant diversification? *Am. Nat.* 138, 881–900.
- Friis, E.M., Pedersen, K.R., Crane, P.R., 2010. Diversity in obscurity: fossil flowers and the early history of angiosperms. *Philos. T. Roy. Soc. B.* 365, 369–382. doi:10.1098/rstb.2009.0227.

- Friis, E.M., Pedersen, K.R., Crane, P.R., 2010. Cretaceous diversification of angiosperms in the western part of the Iberian Peninsula. *Rev. Palaeobot. Palyno.*, 162, 341–361.
- Furness, C.A., Banks, H., 2010. Pollen evolution in the early divergent Monocot order Alismatales. *Int. J. Plant. Sci.* 171, 713–739. doi:10.1086/654848.
- Gallego, J., Gandolfo, M.A., Rubén Cúneo, N., Zamaloa, M.C., 2014. Fossil Araceae from the Upper Cretaceous of Patagonia, Argentina, with implications on the origin of free-floating aquatic aroids. *Rev. Palaeobot. Palyno.* 211, 78–86. doi:10.1016/j.revpalbo.2014.08.017.
- Gibernau, M., Chartier, M., Barabé, D., 2010. Recent advances towards an evolutionary comprehension of Araceae pollination. In: *Diversity, phylogeny and evolution in the monocotyledons*, eds., Seberg, O., Peterson, G., Barfod, A.S., Davis, J.I., Fourth international conference on the comparative biology of the monocotyledons proceedings. Aarhus Univ. Press, Denmark, pp. 101–104.
- Givnish, T.J., 1997. Adaptive radiation and molecular systematics: aims and conceptual issues. In: *Molecular Evolution and Adaptive Radiation*, eds., Givnish, T.J., Sytsma, K.J., Cambridge University Press, New York, pp. 1–54.
- Givnish, T.J., 2010. Ecology of plant speciation. *Taxon* 59, 1326–1366.
- Givnish, T.J., Barfuss, M.H.J., Ee, B.V., Riina, R., Schulte, K., Horres, R., Gonsiska, P.A., Jabaily, R.S., Crayn, D.M., Smith, J.A.C., Winter, K., Brown, G.K., Evans, T.M., Holst, B.K., Luther, H., Till, W., Zizka, G., Berry, P.E., Sytsma, K.J., 2014. Adaptive radiation, correlated and contingent evolution, and net species

- diversification in Bromeliaceae. *Mol. Phylogenet. Evol.* 71, 55–78.
doi:10.1016/j.ympev.2013.10.010.
- Graham, S.W., Zgurski, J.M., McPherson, M.A., Cherniawsky, D.M., Saarela, J.M., Horne, E.F.C., Smith, S.Y., Wong, W.A., O'Brien, H.E., Biron, V.L., Pires, J.C., Olmstead, R.G., Chase, M.W., Rai, H.S., 2006. Robust inference of monocot deep phylogeny using an expanded multigene plastid data set. In: *Monocots: Comparative Biology and Evolution Excluding Poales*, eds., Columbus, J.T., Friar, E.A., Porter, J.M., Prince, L.M., Simpson, M.G., Rancho Santa Ana Botanical Garden, Claremont, Ca., pp. 3–21. [Aliso 22, 3–21.].
- Gravendeel, B., Smithson, A., Slik, F.J.W., Schuiteman, A., 2004. Epiphytism and pollinator specialization: drivers for orchid diversity? *Philos. Trans. R. Soc. Lond., B, Biol. Sci.* 359, 1523–1535. doi:10.1098/rstb.2004.1529.
- Grayum, M.H. 1990. Evolution and phylogeny of the Araceae. *Ann. Mo. Bot. Gard.* 77, 628–697.
- Hay, A., Mabberley, D.J., 1991. Transference of function and the origin of aroids: their significance in early angiosperm evolution. *Bot. Jb*, 113, 339–428.
- Henriquez, C.L., Arias, T., Pires, J.C., Croat, T.B., Schaal, B.A., 2014. Phylogenomics of the plant family Araceae. *Mol. Phylogenet. Evol.* 75, 91–102.
doi:10.1016/j.ympev.2014.02.017.
- Herrera, F.A., Jaramillo, C.A., Dilcher, D.L., Wing, S.L., Gómez-N, C., 2008. Fossil Araceae from a Paleocene neotropical rainforest in Colombia. *Am. J. Bot.* 95, 1569–1583. doi:10.3732/ajb.0800172.

- Hertweck, K.L., Kinney, M.S., Stuart, S.A., Maurin, O., Mathews, S., Chase, M.W., Gandolfo, M.A., Pires, J.C., 2015. Phylogenetics, divergence times and diversification from three genomic partitions in monocots. *Bot. J. Linn. Soc.* doi:10.1111/boj.12260.
- Hesse, M., 2006a. Conventional and novel modes of exine patterning in members of the Araceae –the consequence of ecological paradigm shifts? *Protoplasma* 228, 145–49. doi:10.1007/s00709-006-0165-3.
- Hesse, M., 2006b. Reasons and consequences of the lack of a sporopollenin ectexine in Aroideae (Araceae). *Flora* 201, 421–428. doi:10.1016/j.flora.2005.10.002.
- Hicks, J.F., 1993. Chronostratigraphic analysis of the Foreland Basin sediments of the latest Cretaceous, Wyoming, U.S.A. PhD Thesis, Yale University.
- Hofmann, C.C., Zetter, R., 2010. Upper Cretaceous sulcate pollen from the Timerdyakh Formation, Vilui Basin(Siberia). *Grana* 49, 170–193.
- Iles, W.J., Smith, S.Y., Graham, S.W., 2013. A well-supported phylogenetic framework for the monocot order Alismatales reveals multiple losses of the plastid NADH dehydrogenase complex and a strong long-branch effect. In: *Early Events in Monocot Evolution*, eds., Wilkin, P., Mayo, S.J., Cambridge University Press, Cambridge, pp. 1–28.
- Iles, W.J., Smith, S.Y., Gandolfo, M.A., Graham, S.W., 2015. Monocot fossils suitable for molecular dating analyses. *Bot. J. Linn. Soc.* doi 0.1111/boj.12233.
- Iljinskaja, I.A., 1986. Paleogene floras of Kiin-Kerish Mountain (eastern Kazakhstan) and their new species. *Bot. Zh. (St. Petersburg)* 71, 1193–1202.

- Katoh, K., Misawa, K., Kuma, K., Miyata, T., 2002. MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. *Nucl. Acids. Res.* 30, 3059–3066. doi:10.1093/nar/gkf436.
- Katoh, K., Standley, D.M., 2013. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Mol. Biol. Evol.* 30, 772–780. doi:10.1093/molbev/mst010.
- Kearse, M., Moir, R., Wilson, A., Stones-Havas, S., Cheung, M., Sturrock, S., Buxton, S., Cooper, A., Markowitz, S., Duran, C., Thierer, T., Ashton, B., Meintjes, P., Drummond, A., 2012. Geneious Basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* 28, 1647–1649. doi:10.1093/bioinformatics/bts199.
- Kress, W.J., Stone, D.E., Sellers, S.C., 1978. Ultrastructure of Exine-Less Pollen: *Heliconia* (Heliconiaceae). *Am. J. Bot.* 65, 1064–1076. doi:10.2307/2442323.
- Kubitzki, K., ed., 1998. *The Families and Genera of Vascular Plants*. Vol. 4: Flowering Plants. Monocotyledons: Alismatanae to Commelinanae (except Gramineae). 126, fig. 511. p. 4to. Hardcover.
- Lanfear, R., Calcott, B., Ho, S.Y.W., Guindon, S., 2012. PartitionFinder: Combined selection of partitioning schemes and substitution models for phylogenetic analyses. *Mol. Biol. Evol.* 29, 1695–1701. doi:10.1093/molbev/mss020.
- Les, D.H., Cleland, M.A., Waycott, M., 1997. Phylogenetic studies in Alismatidae II: Evolution of Marine Angiosperms (Seagrasses) and Hydrophily. *Syst. Bot.* 22, 443–463. doi:10.2307/2419820.

- Les, D.H., Tippery, N.P., 2013. In time and with water...the systematics of alismatid monocotyledons. In: *Early Events in Monocot Evolution*, eds., Wilkin, P., Mayo, S.J., Cambridge University Press, New York, pp. 118–164.
- Magallón S, Castillo A., 2009. Angiosperm diversification through time. *Am. J. Bot.* 96: 349–365.
- Magallón, S., Gómez-Acevedo, S., Sánchez-Reyes, L.L., Hernández-Hernández, T., 2015. A metacalibrated time-tree documents the early rise of flowering plant phylogenetic diversity. *New Phytol.* Article first published online: 23 JAN 2015. . doi:10.1111/nph.13264.
- Mayo, S.J., 1978. A new species of *Anthurium* (Araceae) from Bahia, Brazil. *Bradea* 2, 281–286.
- Mayo, S.J., Bogner, J., Boyce, P.C., 1997. *The genera of Araceae*. Kew: Royal Botanic Gardens, Kew.
- Nauheimer, L., D. Metzler, Renner, S.S., 2012. Global history of the ancient monocot family Araceae inferred with models accounting for past continental positions and previous ranges based on fossils. *New Phytol.* 195, 938–950.
- Pagel, M. 1994. Detecting correlated evolution on phylogenies: a general method for the comparative analysis of discrete characters. *Proceedings of the Royal Society (B)* 255, 37-45.
- Pagel, M. and Meade, A. 2006. Bayesian analysis of correlated evolution of discrete characters by reversible-jump Markov chain Monte Carlo. *American Naturalist*, 167, 808-825.

- Paradis, E., Claude, J., Strimmer, K., 2004. APE: Analyses of phylogenetics and evolution in R language. *Bioinformatics* 20, 289–290.
doi:10.1093/bioinformatics/btg412.
- Petersen, G., Seberg, O., Cuenca, A., Stevenson, D.W., Thadeo, M., Davis, J.I., Graham, S., Ross, T.G., 2015. Phylogeny of the Alismatales (Monocotyledons) and the relationship of *Acorus* (Acorales?). *Cladistics*. doi:10.1111/cla.12120.
- Plummer M., Best, N., Cowles, K., Vines, K. 2006. Coda: Convergence Diagnosis and Out-put Analysis for MCMC." *RNews*, 6, 7–11. URL <http://CRAN.R-project.org/doc/Rnews/>.
- R Development Core Team, 2008. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. ISBN 3-900051-07-0, URL <http://www.R-project.org>.
- Rabosky, D.L., 2014. Automatic detection of key innovations, rate shifts, and diversity-dependence on phylogenetic trees. *PLoS One*, 9: e89543. doi: 10.1371/journal.pone.0089543.
- Rabosky, D.L., Grudler, M., Anderson, C., Title, P., Shi, J.J., Brown, J.W., Huang, H., Larson, J.G., 2014. BAMMtools: an R package for the analysis of evolutionary dynamics on phylogenetic trees. *Methods Ecol. Evol.* 5, 701–707.
doi:10.1111/2041-210X.12199.
- Remizowa, M.V., Sokoloff, D.D., Campbell, L.M., Stevenson, D.W., Rudall, P.J., 2011. *Harperocallis* is congeneric with *Isidrogalvia* (Tofieldiaceae, Alismatales): evidence from comparative floral morphology. *Taxon* 60, 1076–1094.

- Revell, L.J., 2012. phytools: an R package for phylogenetic comparative biology (and other things). *Methods Ecol. Evol.* 3, 217–223. doi:10.1111/j.2041-210X.2011.00169.x.
- Ruhfel, B.R., Gitzendanner, M.A., Soltis, P.S., Soltis, D.E., Burleigh, J.G., 2014. From algae to angiosperms –inferring the phylogeny of green plants (Viridiplantae) from 360 plastid genomes. *BMC Evol. Biol.* 14, 23. doi:10.1186/1471-2148-14-23.
- Sanderson, M.J., 2002. Estimating absolute rates of molecular evolution and divergence times: a penalized likelihood approach. *Mol. Biol. Evol.* 19, 101–109. doi: 10.1093/molbev/msm088.
- Sille, N.P., Collinson, M.E., Kucera, M., Hooker, J.J., 2006. Morphological evolution of stratiotes through the Paleogene in England: An example of microevolution in flowering plants. *Palaios* 21, 272–288. doi:10.2110/palo.2005.P05-21e.
- Stamatakis, A., 2006. RAxML-VI-HPC: maximum likelihood-based phylogenetic analyses with thousands of taxa and mixed models. *Bioinformatics* 22, 2688–2690.
- Stamatakis, A., 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* 30, 1312–1313. doi:10.1093/bioinformatics/btu033
- Stevens, P.F., 2001 onwards. Angiosperm Phylogeny Website. Version 12, July 2012 [and more or less continuously updated since] <<http://www.mobot.org/MOBOT/research/APweb/>>.

- Sytsma, K.J., Spalink, D., Berger, B., 2014. Calibrated chronograms, fossils, outgroup relationships, and root priors: re-examining the historical biogeography of Geraniales. *Biol. J. Linnean Soc.* 113, 29–49
- Tanaka, N., Uehara, K., Murata, J., 2004. Correlation between pollen morphology and pollination mechanisms in the Hydrocharitaceae. *J. Plant. Res.* 117, 265–276. doi:10.1007/s10265-004-0155-5.
- WCSP, 2015. World Checklist of Selected Plant Families. Facilitated by the Royal Botanic Gardens, Kew. Published on the Internet. <http://apps.kew.org/wcsp/> Accessed July 2014.
- Wilde, V., Kvaček, Z., Bogner, J., 2005. Fossil leaves of the Araceae from the European Eocene and notes on other aroid fossils. *Int. J. Plant Sci.* 166, 157–183.
- Zeng, L., Zhang, Q., Sun, R., Kong, H., Zhang, N., Ma, H., 2014. Resolution of deep angiosperm phylogeny using conserved nuclear genes and estimates of early divergence times. *Nat. Commun.* 5. doi:10.1038/ncomms5956.

TABLES and FIGURES

Table 1. DNA substitutions models, partitioning scheme and partition subsets resulting from analyses conducted in PartitionFinder using the BIC criterion for RAXML and BEAST analyses.

	Partition no.	Best Model	Partition subsets
RaxML	1	GTR+G	matK_pos1, matk_pos2
	2	GTR+I+G	matk_pos3
	3	GTR+I+G	rbcL_pos1
	4	GTR+I+G	rbcL_pos2
	5	GTR+I+G	rbcL_pos3
BEAST	1	GTR+G	matK_pos1, matk_pos2
	2	GTR+I+G	matk_pos3
	3	GTR+I+G	rbcL_pos1
	4	SYM+I+G	rbcL_pos2
	5	GTR+I+G	rbcL_pos3

Table 2. Fossil calibrations used in the BEAST analyses. The name of the fossil and estimated ages, node constraint by the fossil, and the source are shown. The prior distribution for the calibration is represented by U (Uniform) or L (lognormal). St.Dev represents the log(St.Dev.) value use for the lognormal distribution in Beauti. A range is shown for the root node.

Name of fossil	Age	Node	Prior	St.Dev. or range	Source
<i>Niophylites zaisanicus</i>	55.8	Stem node of <i>Typhonodorum</i> clade, sensu Cusimano et al (2011)	L	0.75	Ijinskaja, 1986; Wilde <i>et al.</i> , 2005
<i>Rhodospathodendron tomlinsonii</i>	66	Stem node of <i>Heteropsis</i> + <i>Rhaphidophora</i> clades	L	0.75	Bonde, 2000
<i>Lasiodeaacidites</i>	70-75	Stem node of subfamily Lasiodeae	L	0.75	Hofmann & Zetter, 2010
<i>Petrocardium cerrejoense</i>	56-61.6	Crown node of Pothoideae	L	0.75	Herrera et al., 2008
“Araceae fossil sp. A” and “Araceae fossil sp. B”	110	Crown node of Lemnoideae + “True Araceae”	L	0.75	Friis et al., 2010
<i>Spixium kipea</i>	113	Stem node of Oroniideae (crown of Proto-Araceae)	L	0.75	Coiffard et al., 2013
<i>Stratiotes</i> sp.	56	Crown of Hydrocharitaceae	L	0.75	Stille et al., 2006
		Stem node of			
<i>Aponogeton hurryi</i>	82-81	Aponogetonaceae (Crown of alismatids)	L	0.75	Hicks, 1993
<i>Liliacidites</i> sp	113-124	Crown node of monocots	L	0.75	Doyle & Hickey, 1976
Root			U	120-160	

Table 3. Calculated ages and confidence intervals (CI) for the main clades in Alismatales from the BEAST analyses with the Tofieldiaceae constraint, the complete constraint constrained topologies and the penalized likelihood. Clade names in Araceae follow Cusimano et al. (2011).

CLADE	Tofieldiaceae constraint		Tree constraint		Penalized likelihood
	Age	CI(5-95%)	Age	CI(5-95%)	Age
ROOT	158.35	(153.37-160)	158.23	(153.09-160)	NA
Monocots	139.94	(132.35-146.9)	138.59	(131.19-146.12)	139
Tofieldiaceae	73.67	(42.93-106.72)	73.66	(44.76-107.04)	73.7
Alismatids	113.61	(102.92-123.59)	112.69	(100.54-124.04)	113.5
Alismatids clade 2	101.08	(88.39-113.1)	100.21	(84.73-114.45)	101.8
Alismatids clade 1	103.49	(90.89-116.23)	102.44	(87.18-115.94)	103.5
Alismatales	129.56	(122.76-136.5)	128.78	(122.15-135.93)	132.1
Alsimatids+Araceae	128.52	(122.32-135.88)	127.73	(121.33-134.96)	125.2
Araceae	121.24	(116.28-127.47)	121.04	(116.13-127.08)	118.3
Proto-Araceae	115	(110.92-120.62)	114.97	(110.95-120.2)	115
Orontioideae	51.97	(22.42-86.71)	52.08	(24.91-87.89)	76.67
Lemnaceae + True Araceae	111.09	(108.65-114.81)	112.11	(109.72-115.8)	111.4
Lemnaceae	74.73	(51.73-102.07)	74.14	(52.73-97.6)	74.7
True Araceae	101.68	(88.33-111.35)	103.32	(90.16-113.14)	104.5
Pothoideae + Monsteroideae	85.44	(73.44-100.11)	85.51	(72.91-100.45)	84.32
Monsteroideae	68.14	(63.97-75.15)	68.35	(63.98-75.64)	64.15
Pothoideae	64.41	(56.47-77.24)	64.73	(56.54-77.72)	55.71
Podolasia clade	87.72	(74.82-101.68)	89.21	(76.98-102.95)	97.6
Zamioculcoideae	63.6	(32.88-87.99)	64.92	(32.74-87.46)	60.47
Lasiodeae	24.09	(12.65-39.9)	26.87	(14.27-45.46)	24.1
Unisexual Clade	83.01	(71.58-96.27)	84.49	(73.14-97.6)	90.7
Aroideae	76.82	(66.7-87.93)	78.6	(68.86-90.37)	83.8
Schismatoglottideae	19.8	(9.09-34.63)	20.63	(9.09-35.97)	38.4
Cryptocoryneae	14.89	(6.59-26.64)	15.18	(6.18-26.96)	35.3
Zantedeschia	51.83	(36.39-68.25)	65.89	(49.04-79.09)	44.6
Spathicarpeae	36.58	(23.95-52.13)	62.32	(45.6-76.04)	41.2
Nephtytideae	14.23	(4.81-27.87)	14.36	(5.02-27.98)	15.4
Aglaonemateae	14.62	(5.25-27.07)	15.44	(5.12-30.23)	17.29
Philodendron clade	19.24	(9.92-32.13)	20.86	(10.62-34.38)	19.2
Culcasieae	24.77	(7.46-44.7)	26.35	(7.72-48.57)	19.21
Calla	68.64	(61.52-78.23)	67.34	(60.56-75.66)	63.1
Dracunculus	62.96	(57.48-70.14)	62.23	(57.19-69.14)	56.2
Ambrosina	59.83	(55.33-65.73)	59.52	(55.4-65.22)	49.3
Protarum Clade	42.43	(30.22-54.04)	42.74	(31.27-54.66)	42.4
Pistia Clade	38.76	(27.88-50.33)	NA	NA	40.7
Thyphonodorum clade	56.43	(53.8-60.51)	56.44	(53.72-60.37)	39.44
Caladieae	34.33	(22.33-48.27)	36.56	(23.92-50.53)	48.42

Thomsonieae	18.52	(7.25-35.92)	19.02	(6.86-36.34)	48.9
-------------	-------	--------------	-------	--------------	------

Table 4. Information from the diversification analyses using BAMM. Mean speciation (Lambda), extinction (Mu) and net diversification are shown. For the two-gene dataset a single shift was recovered; probability and Bayes factors are given.

Model, Dataset	Clade set	Distinct shift configurations in credible set	Number of core shifts	Bayes Factor	Probability	Mean Lambda	Mean Mu	Mean net diversification
BAMM, two-gene	1	34	1	179132.5	0.449	0.249 (0.183,0.328)	0.204 (0.132,0.284)	0.046(0.037-0.055)
BAMM, two-gene	2	4	1	359657.7	0.695	0.311 (0.219,0.425)	0.265 (0.167,0.384)	0.046(0.037-0.055)
MEDUSA-like, two-gene	1	16	1	Inf	0.59	0.279 (0.219,0.345)	0.236 (0.173,0.305)	0.043(0.036-0.051)
MEDUSA-like, two-gene	2	2	1	358543.8	0.799	0.343 (0.270,0.420)	0.3 (0.225,0.381)	0.043(0.036-0.051)
BAMM, ten-gene	1	585	NA	NA	NA	0.098 (0.09,0.108)	0.02 (0.01,0.033)	0.078(0.073-0.082)
BAMM, ten-gene	2	677	NA	NA	NA	0.102 (0.093,0.111)	0.024 (0.013,0.037)	0.078(0.073-0.082)
MEDUSA-like, ten-gene	1	454	NA	NA	NA	0.098 (0.09,0.108)	0.019 (0.009,0.032)	0.079(0.074-0.083)
MEDUSA-like, ten-gene	2	636	NA	NA	NA	0.1 (0.092,0.111)	0.021 (0.011,0.033)	0.079(0.074-0.084)

Table 5. Speciation (Lambda) and extinction rates (Mu) with confidence intervals (CI)

for the major clades in Alismatales from BAMM analyses using the two-gene dataset.

Values for the MEDUSA-like model and the complete BAMM model, for both of the alternative clade sets are provided.

Analysis	Clade set	Clade	Lambda	CI	Mu	CI
BAMM, two-gene	1	Tofieldiaceae	0.055	(0.041-0.079)	0.019	(0.002-0.048)
BAMM, two-gene	2	Tofieldiaceae	0.053	(0.04-0.074)	0.017	(0.001-0.043)
MEDUSA-like, two-gene	1	Tofieldiaceae	0.055	(0.041-0.078)	0.019	(0.002-0.047)
MEDUSA-like, two-gene	2	Tofieldiaceae	0.055	(0.041-0.077)	0.019	(0.002-0.046)
BAMM, two-gene	1	Alismatids	0.055	(0.041-0.077)	0.019	(0.002-0.048)
BAMM, two-gene	2	Alismatids	0.054	(0.041-0.075)	0.017	(0.001-0.043)
MEDUSA-like, two-gene	1	Alismatids	0.055	(0.041-0.077)	0.019	(0.002-0.045)
MEDUSA-like, two-gene	2	Alismatids	0.056	(0.041-0.078)	0.019	(0.002-0.045)
BAMM, two-gene	1	Araceae	0.407	(0.287-0.544)	0.353	(0.222-0.502)
BAMM, two-gene	2	Araceae	0.519	(0.355-0.727)	0.465	(0.292-0.681)
MEDUSA-like, two-gene	1	Araceae	0.46	(0.355-0.577)	0.411	(0.297-0.535)
MEDUSA-like, two-gene	2	Araceae	0.577	(0.447-0.712)	0.527	(0.389-0.669)
BAMM, two-gene	1	True Araceae	0.472	(0.331-0.633)	0.415	(0.262-0.59)
BAMM, two-gene	2	True Araceae	0.605	(0.41-0.848)	0.548	(0.344-0.801)
MEDUSA-like, two-gene	1	True Araceae	0.535	(0.41-0.673)	0.484	(0.349-0.631)
MEDUSA-like, two-gene	2	True Araceae	0.673	(0.517-0.832)	0.621	(0.46-0.791)
BAMM, two-gene	1	Aroideae	0.472	(0.33-0.634)	0.414	(0.26-0.591)
BAMM, two-gene	2	Aroideae	0.607	(0.413-0.852)	0.549	(0.341-0.804)
MEDUSA-like, two-gene	1	Aroideae	0.533	(0.407-0.674)	0.482	(0.345-0.631)
MEDUSA-like, two-gene	2	Aroideae	0.674	(0.519-0.834)	0.622	(0.462-0.795)
BAMM, two-gene	1	Pothoideae + Monsteroideae	0.478	(0.338-0.653)	0.419	(0.267-0.603)
BAMM, two-gene	2	Pothoideae + Monsteroideae	0.607	(0.413-0.849)	0.550	(0.345-0.8)
MEDUSA-like, two-gene	1	Pothoideae + Monsteroideae	0.541	(0.413-0.687)	0.489	(0.35-0.643)
MEDUSA-like, two-gene	2	Pothoideae + Monsteroideae	0.675	(0.521-0.835)	0.623	(0.464-0.796)

Table 6. Values and confidence intervals (CI) for clades with shifts in diversification from the BAMM analyses using the ten-gene dataset. Calculations are provided for speciation (Lambda), extinction rate (Mu), and net diversification (NetDiv) within for major clades of Alismatales, and subfamilies of Araceae. All values are for the complete BAMM model analysis but with the two clade sets of different size (1 & 2) also showed.

Clade Name	Clade set	lambda	CI	mu	CI	NetDiv	CI
Tofieldiaceae	1	0.039	(0.035-0.044)	0.003	(<0.001-0.009)	0.036	(0.032-0.04)
Alismatids	1	0.038	(0.035-0.043)	0.003	(<0.001-0.009)	0.035	(0.031-0.038)
Alismatids tepaloid	1	0.039	(0.035-0.046)	0.004	(<0.001-0.01)	0.035	(0.032-0.039)
Alismatids petaloid	1	0.037	(0.034-0.042)	0.003	(<0.001-0.008)	0.034	(0.03-0.038)
Araceae	1	0.144	(0.131-0.163)	0.032	(0.016-0.056)	0.111	(0.104-0.118)
Proto-Araceae	1	0.046	(0.038-0.054)	0.004	(<0.001-0.019)	0.041	(0.03-0.048)
Lemnoideae	1	0.037	(0.033-0.041)	0.003	(<0.001-0.008)	0.034	(0.029-0.038)
Bisexual climbers clade	1	0.264	(0.21-0.345)	0.098	(0.027-0.197)	0.166	(0.138-0.191)
Pothoideae	1	0.338	(0.248-0.504)	0.14	(0.032-0.331)	0.198	(0.159-0.234)
Monsteroideae	1	0.2	(0.156-0.256)	0.061	(0.008-0.133)	0.139	(0.107-0.167)
Lasioideae	1	0.194	(0.098-0.342)	0.108	(0.004-0.271)	0.086	(0.036-0.134)
<i>Stylochaeton</i> clade	1	0.048	(0.036-0.102)	0.004	(<0.001-0.016)	0.043	(0.033-0.091)
Aroideae	1	0.128	(0.117-0.141)	0.016	(0.005-0.033)	0.112	(0.104-0.12)
<i>Philodendron</i> clade	1	0.397	(0.323-0.508)	0.106	(0.02-0.255)	0.291	(0.238-0.34)
<i>Pistia</i> clade + <i>Protarum</i>	1	0.106	(0.097-0.117)	0.008	(<0.001-0.023)	0.098	(0.09-0.105)
Tofieldiaceae	2	0.04	(0.035-0.045)	0.003	(<0.001-0.009)	0.036	(0.032-0.04)
Alismatids	2	0.039	(0.035-0.044)	0.004	(<0.001-0.009)	0.035	(0.031-0.039)
Alismatids tepaloid	2	0.039	(0.035-0.048)	0.004	(<0.001-0.011)	0.036	(0.032-0.04)
Alismatids petaloid	2	0.038	(0.034-0.043)	0.003	(<0.001-0.009)	0.035	(0.03-0.038)
Araceae	2	0.15	(0.135-0.167)	0.039	(0.02-0.062)	0.111	(0.103-0.118)
Proto-Araceae	2	0.047	(0.04-0.056)	0.006	(<0.001-0.021)	0.042	(0.03-0.049)
Lemnoideae	2	0.037	(0.033-0.041)	0.003	(<0.001-0.009)	0.034	(0.029-0.038)
Bisexual climbers clade	2	0.263	(0.21-0.337)	0.098	(0.03-0.19)	0.165	(0.139-0.188)
Pothoideae	2	0.336	(0.246-0.482)	0.141	(0.033-0.31)	0.196	(0.157-0.229)

Monsteroideae	2	0.199	(0.156-0.255)	0.06	(0.01-0.134)	0.139	(0.112-0.166)
Lasioideae	2	0.163	(0.091-0.314)	0.08	(0.008-0.244)	0.083	(0.043-0.129)
<i>Stylochaeton</i> clade	2	0.072	(0.037-0.124)	0.016	(<0.001-0.054)	0.056	(0.034-0.087)
Aroideae	2	0.138	(0.123-0.156)	0.027	(0.009-0.052)	0.111	(0.101-0.12)
<i>Philodendron</i> clade	2	0.31	(0.25-0.386)	0.068	(0.01-0.168)	0.242	(0.2-0.285)
<i>Pistia</i> clade + <i>Protarum</i>	2	0.146	(0.121-0.181)	0.026	(0.003-0.071)	0.119	(0.102-0.135)

Table 7. Tests for correlated and contingent evolution of the characters associated with subfamily Aroideae. Flower sexuality vs. perianth presence, pollen aperture, and exine reduction. P-values from likelihood ratio test comparing the likelihood of the models produced by BayesTraits are shown. Significance for correlated test is 0.05 and for contingent evolution test is 0.05/4=0.0125

Character	States	Dependent	State 1 evolves	Unisexual flowers	Bisexual flowers	State 0 evolves
		vs. Independent	more often in unisexual flowers	evolved more often in clades with state 1	evolved more often in clades with state 0	more often in bisexual flowers
Perianth	absent(0), present(1)	1	NA	NA	NA	NA
Pollen Aperture	other(0), omniaperturate (1)	5.34E-27	1.60E-20	8.21E-21	3.55E-21	2.44E-20
Exine	complete(0), reduced (1)	1.82E-25	2.89E-02	4.12E-05	7.09E-05	1

Figure 1. Phylogram from maximum likelihood analysis in RaxMLtree using the two-gene dataset and 229 taxa with no constraints. Major clades in Alismatales are labeled and bootstrap support values are shown above the branches.



Figure 2. Phylogram from Maximum likelihood analysis in RaxMLtree using the ten-gene dataset and 946 taxa with no constraints. Major clades in Alismatales are labeled and bootstrap support values are shown above the branches.







Figure 3. Chronograms depicting rates of net diversification obtained with BAMM. Rates are indicated on the branches, with warmer colors representing higher rates, and cooler colors representing lower rates; shifts are indicated by red colors. The top figures correspond to analyses using the complete BAMM model with clade set 1 (left) and clade set 2 (right). The bottom figures correspond to analyses using a MEDUSA-like model with clade set 1 (left) and clade set 2 (right). Bars next to the trees indicate the size of the clades; left bar indicates outgroups (black), Tofieldiaceae, alismatids, Araceae (grey shades); bars to the right indicate clade sets 1 and 2.




Best Configuration BAMM Model, 2 Genes-Clades set 1



Best Configuration BAMM Model, 2 Genes-Clades set 2



Best Configuration Medusa-like Model, 2 Genes-Clades set 1



Best Configuration Medusa-like Model, 2 Genes-Clades set 2

Figure 4. Rates through time plots from the two-gene diversification analyses. Figure **a** shows the rate through time plot for speciation (left), extinction (center), and net diversification (right) for Alismatales; **b** represents a comparison of the speciation (solid line), extinction (dot line), and net diversification (dashed line) for all of the four different two-gene analyses; **c** shows the speciation rate through time plot for All Alismatales (left), the True Araceae (center), and background (Alismatales without “True Araceae” [right]).







Figure 5. Chronograms depicting rates of net diversification obtained with BAMM for the ten-gene dataset with 946 taxa. Rates are indicated on the branches;h warmer colors represent higher rates, cooler color low rates, and shifts are numbered and indicated by red colors. The top figures correspond to analyses using a complete BAMM model with clade set 1(left) and clade set 2 (right). Bottom figures correspond to analyses using a MEDUSA-like model with clade set 1(left) and clade set 2 (right). Bars next to the trees indicate the size of the clades; left bar indicates outgroups (black), Tofieldiaceae, alismatids, Araceae (grey shades); bars to the right indicate clade set 1 and 2.




Best Configuration BAMM Model, 10 Genes–Clades set 1



Best Configuration BAMM Model, 10 Genes–Clades set 1




Best Configuration BAMM Model, 10 Genes–Clades set 1




Best Configuration BAMM Model, 10 Genes–Clades set 1

Figure 6. Bayes factor values associated with the different shifts from the BAMM analysis for the ten-gene dataset using the complete BAMM model and clade set 1 (top figures) and clade set 2 (bottom figures). Figures to the right show branches of the phylogram proportional to the Bayes factor values for the probability of a shift occurring on a particular branch. Figures to the left show branches color code by Bayes factors, yellow >100, orange >500, and red >1000.



Bayes Factors BAMM Model, 10 Genes-Clades set 1



Best Configuration BAMM Model, 10 Genes-Clades set 2

Figure 7. Rate through time plots for the different BAMM analyses with the ten-gene dataset (**A**), and for the different clades in Alismatales (**B**) and Araceae (**C**). **A**, rate through time plots for the BAMM analyses using the complete BAMM and the MEDUSA-like models, and the two different clade sets. Speciation is represented by the solid line and extinction by the dot line. **B**, rate through time plots from the BAMM analysis using the complete BAMM model and the clade set 1. Gray dot lines represent the rates for entire Alismatales (speciation, net diversification and extinction from top to bottom); solid lines represent the net diversification rates of the major clades in Alismatales. **C**, rate through time plots rate through time plots from the BAMM analysis using the complete BAMM model and the clade set 1. Gray dot lines represent the rates for entire Alismatales, speciation (dot line), net diversification (solid line), and extinction (dashed line); solid lines represent the the net diversification rates for the subfamilies in Araceae.




Figure 8. Ancestral state reconstruction for the characters exine reduction (top) and pollen aperture (bottom) in Alismatales. Figures to the left show reconstructions using the rerooting method (ML); figures to the right show reconstruction using 1000 stochastic mappings. States and color codes for each character are shown in the legend.





Figure 9. Ancestral state reconstruction for the characters flower sexuality (top) and perianth (bottom) in Alismatales. Figures to the left show reconstructions using the rerooting method (ML); figures to the right show reconstruction using 1000 stochastic mappings. States and color codes for each character are shown in the legend.







Figure 10. Ancestral state reconstruction for the characters distribution (top) and seasonality (bottom) in Alismatales. Figures to the left show reconstructions using the rerooting method (ML); figures to the right show reconstruction using 1000 stochastic mappings. States and color codes for each character are shown in the legend.







Figure 11. Ancestral state reconstruction for the characters habitat (top) and growth form (bottom) in Alismatales. Figures to the left show reconstructions using the rerooting method (ML); figures to the right show reconstruction using 1000 stochastic mappings. States and color codes for each character are shown in the legend.









Figure 12. Ancestral state reconstruction for the characters seed length (top) and seed width (bottom) in Alismatales. Figures to the left show reconstructions using the rerooting method (ML); figures to the right show reconstruction using 1000 stochastic mappings. States and color codes for each character are shown in the legend.



SUPPLEMENTARY TABLES AND FIGURES

TABLE S1. GenBank accession numbers for the taxa used in the BAMM analyses. NOTE: This table corresponds to all the sequences gathered for Alismatales from GENBANK and our own data. Some were excluded from the analysis because several reasons (length of the sequence, identification issues, repeat sequences). Additionally, GENBANK accessions numbers from chapter two are missing.

Species	CHS	ETS	ITS	mark	rbcl	rp120- rps12	trnC-ycf6	trnG	trnH-psbA	trnLF
<i>Aglaodorum_griffithii</i>				AM920580	AM905758					EF173563
<i>Aglaonema_crispum</i>				EF173528						
<i>Aglaonema_moderatum</i>				AM920579	AM905757					AY054700
<i>Aglaonema_nitidum</i>										GQ220953
<i>Aglaonema_rotundum</i>										AY290843
<i>Alloschemone_occidentalis</i>										AY398567
<i>Allocastia_acuminata</i>				AM920566	AM905744					
<i>Allocastia_alba</i>				JQ238813		JQ238900				
<i>Allocastia_argyrea</i>				JQ238814		JQ238901				
<i>Allocastia_argyrea</i>				JQ238815		JQ238902				
<i>Allocastia_atropurpurea</i>				JQ238816		JQ238903				
<i>Allocastia_beccarii</i>				JQ238817		JQ238904				
<i>Allocastia_boa</i>				JQ238818		JQ238905				
<i>Allocastia_boyceana</i>				JQ238819		JQ238906				
<i>Allocastia_brancheolata</i>				JQ238820		JQ238907				
<i>Allocastia_brisbanensis</i>				JQ238821	KF496352,					
<i>Allocastia_chaitii</i>				JQ238822	JN105555	JQ238908				
<i>Allocastia_clypeolata</i>				JQ238823		JQ238909				
<i>Allocastia_cucullata</i>	AB429332			EU886579	JQ237188	JQ238910				JQ237218
<i>Allocastia_cucullata</i>										
<i>Allocastia_culionensis</i>				JQ238825		JQ238911				

<i>Alocasia_cuprea</i>		IQ238826		IQ238912	
<i>Alocasia_denudata</i>		IQ238827		IQ238913	
<i>Alocasia_gageana</i>		EU886580		AY248909	
<i>Alocasia_grandis</i>		IQ238828			IQ238743
<i>Alocasia_heterophylla</i>		IQ238829		IQ238914	
<i>Alocasia_holtrungii</i>		IQ238830		IQ238915	
<i>Alocasia_infernalis</i>		IQ238832		IQ238917	IQ238745
<i>Alocasia_inornata</i>		IQ238833			
<i>Alocasia_korthalsii</i>		IQ238834		IQ238918	
<i>Alocasia_lauterbachiana</i>		IQ238835		IQ238919	
<i>Alocasia_longiloba</i>		IQ238836		IQ238922	
<i>Alocasia_lowii</i>		IQ238840		IQ238924	IQ238751
<i>Alocasia_macrorrhizos</i>		IQ238841	JN407238	IQ238925	
<i>Alocasia_maquilingensis</i>		IQ238842		IQ238926	
<i>Alocasia_melo</i>		IQ238843		IQ238927	
<i>Alocasia_micholiziana</i>		IQ238844			
<i>Alocasia_monticola</i>		IQ238845		IQ238928	
<i>Alocasia_navicularis</i>		EU886581		AY248925	
<i>Alocasia_nebula</i>		IQ238846		IQ238929	
<i>Alocasia_nyctericis</i>		IQ238847		IQ238930	
<i>Alocasia_odora</i>	AB429312	IQ238848	KC466582	IQ238931	
<i>Alocasia_odora</i>					AF469028
<i>Alocasia_pungeran</i>		IQ238849		IQ238932	
<i>Alocasia_peltata</i>		IQ238850		IQ238933	
<i>Alocasia_perakensis</i>		IQ238851			IQ238767
<i>Alocasia_plumbea</i>		IQ238852		IQ238934	
<i>Alocasia_portei</i>		IQ238853		IQ238935	
<i>Alocasia_princeps</i>		IQ238854		IQ238936	
<i>Alocasia_principiculus</i>		IQ238855		IQ238937	
<i>Alocasia_ramosii</i>		IQ238856		IQ238938	
<i>Alocasia_reginae</i>		IQ238857		IQ238939	
<i>Alocasia_reginula</i>		IQ238858		IQ238940	
<i>Alocasia_reversa</i>		IQ238859		IQ238941	
<i>Alocasia_ridleyi</i>		IQ238860		IQ238942	
<i>Alocasia_robusta</i>		IQ238862		IQ238945	
<i>Alocasia_sanderiana</i>		IQ238864	EU193192	IQ238946	

<i>Alocasia_sarawakensis</i>	JQ238865	JQ238947	
<i>Alocasia_scabrituscula</i>	JQ238866	JQ238948	
<i>Alocasia_scalprum</i>	JQ238867	JQ238949	
<i>Alocasia_simata</i>	JQ238868	JQ238950	
<i>Alocasia_sp._Bo011</i>	JQ238869	JQ238951	
<i>Alocasia_sp._Bo07</i>	JQ238870	JQ238952	
<i>Alocasia_sp._Gomantong</i>	JQ238871	JQ238953	
<i>Alocasia_sp._inopinata</i>	JQ238879	JQ238961	
<i>Alocasia_sp._kelamensis</i>	JQ238880	JQ238962	
<i>Alocasia_sp._Kerby</i>	JQ238872	JQ238954	
<i>Alocasia_sp._kulat</i>	JQ238881	JQ238963	
<i>Alocasia_sp._MPM1</i>	JQ238873	JQ238955	
<i>Alocasia_sp._MPM2</i>	JQ238874	JQ238956	
<i>Alocasia_sp._MPM3</i>	JQ238875	JQ238957	
<i>Alocasia_sp._MPM4</i>	JQ238876	JQ238958	
<i>Alocasia_sp._ridleythai</i>	JQ238882	JQ238964	
<i>Alocasia_sp._Septier</i>	JQ238877	JQ238959	
<i>Alocasia_sp._Setiam</i>	JQ238878	JQ238960	
<i>Alocasia_suhirmaniana</i>	JQ238883	JQ238965	
<i>Alocasia_warburgii</i>	JQ238884	JQ238966	
<i>Alocasia_watsoniana</i>	JQ238886	JQ238967	
<i>Alocasia_wongii</i>	JQ238887	JQ238969	
<i>Alocasia_zebrina</i>	JQ238888	JQ238970	
<i>Ambrosina_bassii</i>	AM920620	AM905798	AY555185
<i>Amorphophallusabyssinicus</i>	AF387379	AF497060	
<i>Amorphophallusabyssinicus</i>			AF387433
<i>Amorphophallus_albus</i>	EU193570	EU193179	
<i>Amorphophallus_amygdales</i>		DQ012482	
<i>Amorphophallus_angolensis</i>	AF387404	AF497061	AF387458
<i>Amorphophallus_angolensis</i>			
<i>Amorphophallus_angustispathu</i> ^s		DQ012483	
<i>Amorphophallus_ankarana</i>	AF387380	AF497062	AF387434
<i>Amorphophallus_baumannii</i>	AF387382	AF497063	AF387436
<i>Amorphophallus_beccarii</i>	AF387383	AF497064	AF387437
<i>Amorphophallus_bhandarensis</i>		KM093849	

<i>Amorphophallus_bonaccordensis</i> <i>is</i>			
<i>Amorphophallus_borneensis</i>	KM093850		
<i>Amorphophallus_brevispatus</i>	DQ012484		
<i>Amorphophallus_bulbifer</i>	AF387384	AF497065	AF387438
<i>Amorphophallus_canaliculatus</i>	EU542582	KM093839	
<i>Amorphophallus_cirrififer</i>	AF387385	AF497066	AF387439
<i>Amorphophallus_coetaneus</i>	AF387386	AF497067	AF387440
	AF387381	AF497068	AF387435
		KM093835, KM093836, KM093837, KM093838, AF497069	
<i>Amorphophallus_commutatus</i>	AF387387	AF387392	AF387441
<i>Amorphophallus_corrugatus</i>		AF497070	AF387442
<i>Amorphophallus_dactylifer</i>		DQ012485	
<i>Amorphophallus_declinatus</i>		DQ012486	
<i>Amorphophallus_decus-silvae</i>	AF387388	AF497071	AF387443
<i>Amorphophallus_discophorus</i>		DQ012487	
<i>Amorphophallus_dracontioides</i>	AF387389	AF497072	AF387444
<i>Amorphophallus_eburneus</i>	AF387390	AF497073	AF387445
<i>Amorphophallus_eichleri</i>	AF387391	AF497074	AF387446
<i>Amorphophallus_galbra</i>	AF387393	AF497075	AF387447
<i>Amorphophallus_glossophyllus</i>		DQ012489	
<i>Amorphophallus_henryi</i>	AF387394	AF497076	AF387448
<i>Amorphophallus_hewitii</i>		DQ012490	
<i>Amorphophallus_hirsutus</i>		KM093840,	
<i>Amorphophallus_hirtus</i>	AF387395	AF497077	AF387449
	AF387396	AF497078	AF387450
		KM093843, DQ012491	
<i>Amorphophallus_hohenackeri</i>		AF497079	AF387451
<i>Amorphophallus_hottiae</i>	AF387397	DQ012493	
<i>Amorphophallus_impersus</i>		DQ012492	
<i>Amorphophallus_interruptus</i>		DQ012494	
<i>Amorphophallus_johnsonii</i>		AF497080	AF387452
<i>Amorphophallus_konjac</i>	AF387398	KM093842, DQ012495	
<i>Amorphophallus_konkanensis</i>			

<i>Amorphophallus krausei</i>	AF387399	AF497081		AF387453
<i>Amorphophallus lambii</i>	AF387400	AF497082		AF387454
<i>Amorphophallus lanuginosus</i>		DQ012496		
<i>Amorphophallus laoticus</i>		DQ012497		
<i>Amorphophallus lewalliei</i>	AF387401	AF497083		AF387455
<i>Amorphophallus longiconnectivus</i>		KM093851, DQ012498		
<i>Amorphophallus longituberosus</i>	AF387402	AF497084		AF387456
<i>Amorphophallus margaritifer</i>		KM093844,		
<i>Amorphophallus maxwellii</i>	AF387403	AF497085		AF387457
<i>Amorphophallus mossambicensis</i>	AF387405	AF497086		AF387459
		DQ012499		
<i>Amorphophallus muelleri</i>		KM093841,		
<i>Amorphophallus mysorensis</i>	AF387406	AF497087		AF387460
<i>Amorphophallus nepalensis</i>		KM093852		
<i>Amorphophallus nepalensis</i>	AF387408	AF497088		AF387462
<i>Amorphophallus napiger</i>	AF387407	AF497089		AF387461
<i>Amorphophallus nicolsonianus</i>		KM093853		
<i>Amorphophallus ochroleucus</i>	AF387409	AF497090		AF387463
		KM093846, KM093847, AF497091		
<i>Amorphophallus paeoniifolius</i>	AF387410			AY054703
<i>Amorphophallus palawanensis</i>	AF387411	AF497092		AF387465
<i>Amorphophallus pendulus</i>		DQ012501		
<i>Amorphophallus pingbianensis</i>	AF387412	AF497093		AF387466
<i>Amorphophallus pusillus</i>	AF387413	AF497094		AF387467
<i>Amorphophallus pygmaeus</i>	AF387414	AF497095		AF387468
<i>Amorphophallus rhizomatosus</i>	AF387415	AF497096		AF387469
<i>Amorphophallus sagittarius</i>	AF387416	AF497097		AF387470
<i>Amorphophallus salmoneus</i>		DQ012502		
<i>Amorphophallus scutatus</i>		DQ012503		
		KM093854,		
<i>Amorphophallus smithsonianus</i>	AF387417	AF497098		AF440738
<i>Amorphophallus sp. KP-2014</i>		KM093845		
<i>Amorphophallus sumawongii</i>	AF387418	AF497099		AF387471
<i>Amorphophallus sylvaticus</i>		KM093855		

<i>Amorphophallus symonians</i>	AF387419	AF497100	AF387472
<i>Amorphophallus taurostigma</i>	AF387420	AF497101	AF387473
<i>Amorphophallus thaensis</i>		DQ012504	
<i>Amorphophallus tinekeae</i>		DQ012505	
<i>Amorphophallus titanium</i>	AF387421	AF497102	AY555180
<i>Amorphophallus variabilis</i>	AF387422	AF497103	AF387475
<i>Amorphophallus yunnanensis</i>	AF387423	AF497104	AF387476
<i>Amorphophallus zenkeri</i>	AF387424	AF497105	AF387477
<i>Amydrum hainanense</i>			AY398552
<i>Amydrum humile</i>	AM920567	AM905745	AY398550
<i>Amydrum magnificum</i>			AY054735
<i>Amydrum medium</i>			AY555161
<i>Anadendrum microstachyum</i>	AM920547	AM905740	AY398559
<i>Anadendrum_sp._Boyce_1173</i>	AM920575	AM905753	AY398560
<i>Anadendrum_sp._Chase_9985</i>			AY054726
<i>Anaphyllopsis americana</i>	HQ687766		
<i>Anaphyllum wightii</i>	AF387427	AF497108	AY555186
<i>Anchomanes difformis</i>	AM920557	AM905735	
<i>Anthurium acule</i>			
<i>Anthurium andicola</i>	JX894473	JX894684	JX894791
<i>Anthurium andraeanum</i>	JX894474	JX894685	JX894792
	JN602206	FN668824	
<i>Anthurium antioquiense</i>			JX894686
<i>Anthurium bakeri</i>	JX894475	JX894578	JX894793
<i>Anthurium balaanum</i>	JX894476	JX894687	JX894579
<i>Anthurium bellum</i>	JQ586582	JQ590265	JX894794
<i>Anthurium berriozabalense</i>	JX894477	JX894688	JX894580
<i>Anthurium besseae</i>	JX894478	JX894689	JX894581
<i>Anthurium bredeanum</i>	JX894479	JX894690	JX894796
<i>Anthurium brenesii</i>	JX894480	JX894691	JX894582
<i>Anthurium brevipedunculatum</i>	JX894482	JX894693	JX894583
<i>Anthurium bromelicola_subsp._bahense</i>	JX894483	JQ589934	JX894585
<i>Anthurium brownii</i>	JX894484	JQ594929	JX894694
<i>Anthurium carchiense</i>	JX894485		JX894586
<i>Anthurium chiapasense_subsp._tlaxiicense</i>	JX894486		JX894587
	JX894489		JX894695
			JX894696
			JX894588
			JX894697
			JX894589
			JX894804
			JX894700
			JX894592
			JX894807

<i>Anthurium citrifolium</i>	JX894490				JX894701	JX894593	JX894808	
<i>Anthurium clarinervium</i>	JX894491				JX894702	JX894594	JX894809	
<i>Anthurium clavigerum</i>	JX894492				JX894703	JX894595	JX894810	
<i>Anthurium clidmioides</i>	JX894471				JX894704	JX894596	JX894811	
<i>Anthurium consobrinum</i>			JQ586585	JQ590267				AY555155
<i>Anthurium cordatum</i>	JX894493				JX894705	JX894597	JX894812	
<i>Anthurium cordifolium</i>	JX894494				JX894706	JX894598	JX894813	
<i>Anthurium coriaceum</i>	JX894495				JX894707	JX894599	JX894814	
<i>Anthurium crenatum</i>	JX894496				JX894708	JX894600	JX894815	
<i>Anthurium croatii</i>	JX894497				JX894709	JX894601	JX894816	
<i>Anthurium crystallinum</i>	JX894498				JX894710	JX894602	JX894817	
<i>Anthurium curvispadix</i>	JX894499				JX894711	JX894603	JX894818	
<i>Anthurium cuspidatum</i>	JX894500				JX894712	JX894604	JX894819	
<i>Anthurium digitatum</i>	JX894501				JX894713	JX894605	JX894820	
<i>Anthurium dolichostachyum</i>	JX894502				JX894714	JX894606	JX894821	
<i>Anthurium dominicense</i>	JX894503				JX894715	JX894607	JX894822	
<i>Anthurium dressleri</i>	JX894504				JX894716	JX894608	JX894823	
<i>Anthurium durandi</i>	JX894481				JX894692	JX894584	JX894799	
<i>Anthurium emiens</i>	JX894505				JX894717	JX894609	JX894824	
<i>Anthurium flexile</i> subsp. <i>flexile</i>								
<i>e</i>	JX894472				JX894718	JX894610	JX894825	
<i>Anthurium formosum</i>	JX894506				JX894719	JX894611	JX894826	
<i>Anthurium formicifolium</i>	JX894507				JX894720	JX894612	JX894827	
<i>Anthurium friedrichshalii</i>	JX894508				JX894721	JX894613	JX894828	
<i>Anthurium furcatum</i>	JX894509				JX894722	JX894614	JX894829	
<i>Anthurium gracile</i>	JX894510				JX894723	JX894615	JX894830	
<i>Anthurium graciliacuminatum</i>	JX894511				JX894724	JX894616	JX894831	
<i>Anthurium grandifolium</i>	JX894512				JX894725	JX894617	JX894832	
<i>Anthurium halmorei</i>	JX894513				JX894726	JX894618	JX894833	
<i>Anthurium harrisii</i>	JX894514				JX894727	JX894619	JX894834	
<i>Anthurium hoffmannii</i>	JX894515				JX894728	JX894620	JX894835	
<i>Anthurium huxiense</i>	JX894516				JX894729	JX894621	JX894836	
<i>Anthurium interruptum</i>			JQ586589	JQ590272				
<i>Anthurium kaulthii</i>	JX894517				JX894730	JX894622	JX894837	
<i>Anthurium lacinosum</i>	JX894518				JX894731	JX894623	JX894838	
<i>Anthurium lancea</i>	JX894519				JX894732	JX894624	JX894839	

<i>Anthurium lanceolense</i>	JX894520			JX894733	JX894625	JX894840	
<i>Anthurium lancifolium</i>	JX894521			JX894734	JX894626	JX894841	
<i>Anthurium leuconeurum</i>	JX894522			JX894735	JX894627	JX894842	
<i>Anthurium longicaudatum</i>				JX894737	JX894629	JX894844	
<i>Anthurium longipes</i>	JX894524			JX894738	JX894630	JX894845	
<i>Anthurium longissimum</i>	JX894525			JX894739	JX894631	JX894846	
<i>Anthurium lucens</i>	JX894526			JX894740	JX894632	JX894847	
<i>Anthurium lucidum</i>	JX894527			JX894741	JX894633	JX894848	
<i>Anthurium maclearii</i>	JX894529			JX894743	JX894635	JX894850	
<i>Anthurium maculosum</i>	JX894530			JX894744	JX894636	JX894851	
<i>Anthurium margaricarpum</i>	JX894531			JX894745	JX894637	JX894852	
	JX894532,			JX894746,	JX894638,	JX894853,	
<i>Anthurium michelii</i>	JX894487			JX894698	JX894590	JX894805	
<i>Anthurium microspadix</i>	JX894533			JX894747	JX894639	JX894854	
<i>Anthurium mindense</i>	JX894534			JX894748	JX894640	JX894855	
<i>Anthurium morae</i>	JX894535			JX894749	JX894641	JX894856	
<i>Anthurium myunense</i>	JX894536			JX894750	JX894642	JX894857	
<i>Anthurium napaeum</i>				JX894751	JX894643	JX894858	
<i>Anthurium obtusilobum</i>	JX894537			JX894752	JX894644	JX894859	
	JX894488,			JX894753,	JX894591,	JX894806,	
<i>Anthurium obtusum</i>	JX894538			JX894699	JX894645	JX894860	
<i>Anthurium ochrantum</i>			IQ586590	IQ590274			AY290833
<i>Anthurium ovatifolium</i>	JX894540				JX894755	JX894647	JX894862
<i>Anthurium oxyphyllum</i>	JX894541				JX894756	JX894648	JX894863
<i>Anthurium palenquense</i>					JX894757	JX894649	JX894864
<i>Anthurium parasiticum</i>	JX894539				JX894754	JX894646	JX894861
<i>Anthurium parvispathum</i>	JX894542				JX894758	JX894650	JX894865
<i>Anthurium parvum</i>	JX894543				JX894759	JX894651	JX894866
<i>Anthurium pedatoradiatum</i>	JX894544				JX894760	JX894652	JX894867
<i>Anthurium pedatum</i>	JX894545						
<i>Anthurium peligerum</i>	JX894546				JX894762	JX894654	JX894869
<i>Anthurium polydactylum</i>	JX894547				JX894763	JX894655	JX894870
<i>Anthurium protensum</i>	JX894548				JX894764	JX894656	JX894871
<i>Anthurium punctatum</i>	JX894549				JX894765	JX894657	JX894872
<i>Anthurium radicans</i>	JX894550				JX894766	JX894658	JX894873
<i>Anthurium roventii</i>			IQ589978	IQ590282			AY290835

[illegible]

<i>Arisaema_album</i>				AY279138	
<i>Arisaema_amurense</i>		KC466571	DQ859161		
<i>Arisaema_amurense_var._serratum</i>	FJ980445, GQ434867 AF291914				
<i>Arisaema_angustatum</i>				AY248912	
<i>Arisaema_aridum</i>				AY376845	
<i>Arisaema_auriculatum</i>				AY279139	AY275568
<i>Arisaema_balansae</i>					
<i>Arisaema_bonatianum</i>			JQ237193		JQ237223
<i>Arisaema_candidissimum</i>				AY279140	
<i>Arisaema_ciliatum</i>				AY248913	
<i>Arisaema_clavatum</i>				AY279142	AY275595
<i>Arisaema_concinnum</i>				AY279143	AY275591
<i>Arisaema_consanguineum</i>	KF913386	KF913421			AY275587
<i>Arisaema_costatum</i>		KC466572	KC466584	AY279144	
<i>Arisaema_costatum</i>					
<i>Arisaema_elephas</i>		KC466573	KC466585	AY248914	AY248951, AY248989
			JQ237194	AY279145	
<i>Arisaema_erubescens</i>	JF975895, JF975892	GQ434310	GQ436779	AY279146	JN043993
<i>Arisaema_fargesii</i>				AY279148	AY275592
	JF975899, JF975900	JF953250	JF940894	AY248915	
<i>Arisaema_flavum</i>				AY279151	JN043994
<i>Arisaema_fornosanum</i>		AM920628	AM905806	AY279152	
<i>Arisaema_franchetianum</i>				AY279153	
<i>Arisaema_hamandii</i>				AY279154	
<i>Arisaema_heterocephalum</i>				AY248916	
<i>Arisaema_heterophyllum</i>	FJ980446	KC466574	GQ436775	AY279155	JN043996
<i>Arisaema_itanense</i>				AY279156	
<i>Arisaema_jacquemonitii</i>				AY248917	
<i>Arisaema_macrospathum</i>		KC466575	KC466587	AY279158	
<i>Arisaema_matsudae</i>				AY279159	
<i>Arisaema_microspadix</i>					AY275604
<i>Arisaema_mildbraedii</i>					
<i>Arisaema_negishii</i>				AY279161	
<i>Arisaema_nepenthoides</i>				AY279162	AY275584

<i>Arisaema polyphyllum</i>						AY248918	
<i>Arisaema rhizonatum</i>						AY248919	
<i>Arisaema ringens</i>							JN043999
<i>Arisaema schimperianum</i>		AF274295		JF953255	JF940900		
<i>Arisaema serratum</i>							
<i>Arisaema sikokianum</i>		AB605025,		KC466576	KC466588	AY279166	
<i>Arisaema siko-</i>		EF017383		EU193586	EU193195	AB494543	
<i>kianum</i> var. <i>s-</i>		AB513178					
<i>erratum</i>							
<i>Arisaema speciosum</i>		GQ434307	GQ436776	EU886502	EU193196	AY279168	EU193411
<i>Arisaema thun-</i>							
<i>bergii</i> subsp. <i>-</i>							
<i>urashima</i>	AB894250						
<i>Arisaema tortuosum</i>				AF387428	AF497109	AY248920	AF387481
<i>Arisaema tosaense</i>							
<i>Arisaema triphyllum</i>		AB513179					
<i>Arisaema und-</i>		KF977433,					
<i>ulatifolium</i> su		AF203678		KC466577	AY298817	AY248921	HQ596605
<i>bsp. uwajimen</i>							
<i>se</i>	AB611695						
<i>Arisaema watii</i>		AF469031			JQ237195		
<i>Arisarum vulgare</i>				EU886582	HM849791	EU886630	
<i>Arophyton buchetii</i>				AM920642	AM905820		AY555181
<i>Arum alpinum</i>				EU193636	EU193245		
<i>Arum apulum</i>		GU067541		GU067591	GU067566		
<i>Arum balansanum</i>		GU067542		EU886512	GU067567	EU886624	GU067617
<i>Arum bessenianum</i>							
<i>Arum byzantinum</i>		GU067543		GU067593	GU067568		
<i>Arum concinnum</i>		GU067544		EU886516	GU067569	GU255991	GU067618
<i>Arum creticum</i>		GU067545		EU886504	GU067570	EU886595	
<i>Arum cylindraceum</i>		GU067540		EU886514	EU193244	EU886626	
<i>Arum cyrenaicum</i>		GU067546		EU886515	GU067581	EU886623	
<i>Arum dioscoridis</i>		GU067547		EU886505	GU067572		
<i>Arum dioscoridis</i> var. <i>cypriu</i>							
<i>m</i>							
<i>Arum dioscoridis</i> var. <i>dioscor</i>							

<i>idlis</i>								
<i>Arum elongatum</i>	GU067549	GU067598	GU067573					
<i>Arum elongatum_subsp. elongatum</i>								
<i>Arum euxinum</i>	GU067548	GU067599	GU067574					GU067624
<i>Arum gratum</i>	GU067552	GU067600	GU067575					
<i>Arum hygrophilum</i>	GU067550	GU067601	GU067576			EU886620		AM932296
<i>Arum idaeum</i>	GU067553	GU067602	GU067577					
<i>Arum italicum</i>	GU067554	EU886517	GU067578					
<i>Arum italicum_subsp. albispatham</i>								
<i>Arum italicum_subsp. canariense</i>	EU193627	EU193236						
<i>Arum jacquemontii</i>	GU067555	GU067604	GU067579					GU067630
<i>Arum korolkowii</i>	GU067556	EU886589	GU067580		EU886598			GU067631
<i>Arum lucanum</i>								
<i>Arum maculatum</i>	GU067558	EU886506	GU067582		EU886593			
<i>Arum megobrybi</i>		EU886513			EU886625			
<i>Arum nigrum</i>	GU067559	EU886507	GU067583		EU886597			
<i>Arum orientale</i>		EU886510	EU193240		EU886621			
<i>Arum orientale_subsp. longispatum</i>								
<i>Arum orientale_subsp. orientale</i>	EU193639	EU193248						
<i>Arum palaestinum</i>	GU067551	GU067607	GU067584					GU067629
<i>Arum pictum</i>	GU067560	EU886518	GU067585		EU886596			AY555182
<i>Arum purpureospatum</i>	GU067561	EU886508	GU067586		EU886594			
<i>Arum rupicola</i>	GU067562	EU886519	GU067587		EU886592			
<i>Arum rupicola_subsp. rupicola</i>								
<i>Arum rupicola_subsp. virescens</i>								
<i>Arum sinensisii</i>	GU067563	GU067613	GU067588					
<i>Arum_sp. Lobin_26940</i>					EU886622			
<i>Asterostigma_cryptostylum</i>		EF173530						EF173565

<i>Asterostigma cubense</i>			EF173531			EF173566
<i>Asterostigma lividum</i>			EF173532			EF173567
<i>Asterostigma lombardii</i>			EF173533			EF173568
			EF173534, EF173535			EF173569
<i>Asterostigma reticulatum</i>			EF173536			EF173571
<i>Asterostigma tweedii</i>			GQ220894			GQ220962
<i>Bakoa lucens</i>			EU193608	EU193217		
<i>Biarrum arundanum</i>			EU886529			
<i>Biarrum bovei</i>			EU886521			
<i>Biarrum carduchorum</i>			EU886525			
<i>Biarrum davisii</i>						
<i>Biarrum davisii_subsp._davisii</i>				EU193220		
<i>Biarrum davisii</i>						
<i>i_subsp._mar</i>						
<i>marisense</i>		EU193221				
<i>Biarrum dispar</i>			EU886522	EU193222		
<i>Biarrum dischianum</i>			EU886526			
<i>Biarrum kotschy</i>			EU886527			
<i>Biarrum pyrami</i>			EU886523			
<i>Biarrum straussii</i>			EU886524			
<i>Biarrum tenuifolium</i>			EU886528	AM905810		
<i>Biarrum tenuif</i>						
<i>oilum_subsp._</i>						
<i>abbreviatum</i>	EU193603	EU193212				
<i>Biarrum tenuif</i>						
<i>oilum_subsp._i</i>						
<i>domenaeum</i>	EU193605	EU193214				
<i>Biarrum tenuif</i>						
<i>oilum_subsp._</i>						
<i>zeleborii</i>	EU193602	EU193211				
<i>Bognera recondita</i>			EF173537	AM905765		EF173572
<i>Bucephalandra molleyana</i>	JN544425	AM920644	AM905822		KC454008	GQ220963
<i>Caladium bicolor</i>		EU886501	EU193181	AY248943		AY054708
<i>Caladium lindenii</i>		AM920610	AM905788			AF521871
<i>Calla palustris</i>	AF168829	HQ687764	AM905819			AY555174
<i>Calloopsis volkensii</i>		AM920595	AM905773			AM932330
<i>Carlephyton glaucophyllum</i>		AM920643	AM905821			

<i>Cercesis mirabilis</i>	AM920639	AM905817		EF173564
<i>Cercesis stigmaticus</i>				AY555170
<i>Chlorospatha atropurpurea</i>	EU542584			
<i>Chlorospatha longipoda</i>				AF521872
<i>Chlorospatha sp._Chase_1191</i> <i>2</i>	AM920613, AM905791, AM920613 AM905791	AM905791, AM905791		
<i>Colletogyne perrieri</i>	AM920645	AM905823		
<i>Colocasia affinis var._jenningsii</i>	JQ238889	JN105544	JQ238971	
<i>Colocasia antiquorum</i>	AH001599, AF469032	JF828113	JQ237199	JF828132
<i>Colocasia esculenta</i>	AY081000	JQ238890	AM905800	JQ238972
<i>Colocasia esculenta var._fo</i> <i>manesii</i>				AB494616
<i>Colocasia fallax</i>	JQ238892	JQ238974		GU135448
<i>Colocasia formosana</i>	JQ238891	JN105548	JQ238973	AF469018
<i>Colocasia heterochroma</i>	JN105375	JN105547		AY555184
	JF828117	JF828099		
		JF828094, JF828098		JF828138
<i>Colocasia lilengiae</i>	JF828118	JF828098		
<i>Colocasia menglaensis</i>	JQ238894		JQ238976	
<i>Colocasia sp._Boyce s.n.</i>	JQ238895		JQ238977	
<i>Colocasia yunnanensis</i>	JF828112	JF828092		
<i>Croatitella integrifolia</i>	EF173538			JF828131
<i>Cryptocoryne auriculata</i>	JX218026	JX258151		EF173573
<i>Cryptocoryne balansae</i>				
<i>Cryptocoryne bulbosa</i>	JX258137	JX196335		AY555173
<i>Cryptocoryne ciliata</i>	L41594, JN695007			
<i>Cryptocoryne ciliata var._latifolia</i>	JX258138	JX196336		
<i>Cryptocoryne cordata var._zonata</i>	JX258139	JX196337		
<i>Cryptocoryne ferruginea</i>	JX258141	JX196339		GQ220965
<i>Cryptocoryne fusca</i>	JX258142	JX196340		
<i>Cryptocoryne keei</i>	JX258143	JX196341		
<i>Cryptocoryne lingua</i>	AM920601	AM905779		AM933329
<i>Cryptocoryne longicauda</i>	JX258145	JX196343		

<i>Cryptocoryne pallidivervia</i>		JX258146	JX196344	
<i>Cryptocoryne</i> sp. <i>FL09-16</i>		JX258150	JX196349	
<i>Cryptocoryne spiralis</i>	JQ436581			
<i>Cryptocoryne striolata</i>		JX258148	JX196347	
<i>Cryptocoryne uenoi</i>		JX258140	JX196338	
<i>Cryptocoryne yuiii</i>		GQ220897		GQ220966
<i>Cryptocoryne zaidiana</i>		JX258149	JX196348	
<i>Culcasia liberica</i>		AM920638	AM905816	
<i>Culcasia manni</i>				AY555169
<i>Culcasia saxatilis</i>				AY054713
<i>Culcasia striolata</i>				AY290844
<i>Cyrtosperma cuspidispalum</i>		AM920572	AM905750	AY290836
<i>Cyrtosperma macrodon</i>				
<i>Dieffenbachia aglaonematiifolia</i>		AM920586	AM905764	
<i>Dieffenbachia humilis</i>		EF173540		EF173574
<i>Dieffenbachia oerstedii</i>		EF173541	JQ590304	EF173575
<i>Dieffenbachia parlatorei</i>		EF173542		EF173576
<i>Dieffenbachia seguiniae</i>		FN668814		
<i>Dieffenbachia</i> sp. <i>A.Guadalupe</i>			JQ590299	
<i>Dieffenbachia</i> sp. <i>Goncalves_164</i>		EF173543		
<i>Dieffenbachia</i> sp. <i>KRI-2009</i>		GQ429082	AI005631	
<i>Dieffenbachia</i> sp. <i>Qiu_96007</i>		GQ220898		
<i>Dieffenbachia spruceana</i>		JQ586623	JQ590319	
<i>Dieffenbachia tonduzii</i>		AM920576	AM905754	AM932316
<i>Dracontoides desciscens</i>		JQ586629	JQ590325	
<i>Dracontium gigas</i>		AM920569	AM905747	AY054727
<i>Dracontium polyphyllum</i>		EU886531	EU193230	
<i>Dracunculus canariensis</i>	GU067564	EU886532	GU067589	
<i>Dracunculus vulgaris</i>		EU886520		
<i>Eminium jaegeri</i>		EU886530	AM905813	
<i>Eminium spiculatum</i>		JQ238831		JQ238916
<i>Englerarum hyposomum</i>		GQ434255	AI005625	
<i>Epipremnum aureum</i>				

<i>Epipremnum_falcifolium</i>				AY398558 LC000733, AY398557
<i>Epipremnum_papuanum</i>				
<i>Epipremnum_pinnatum</i>				
<i>Filium_manserichense</i>				
<i>Furtadoa_mixta</i>				
<i>Furtadoa_sumatrensis</i>				
<i>Gearum_brasiliense</i>				
<i>Gonatopus_angustus</i>				
<i>Gorgonidium_intermedium</i>				
<i>Gorgonidium_striatum</i>				
<i>Gorgonidium_vermicidum</i>				
<i>Gymnostachys_anceps</i>				
<i>Hapaline_benthamiana</i>				
<i>Hapaline_ellipticifolium</i>				
<i>Hapaline_sp._HAR_056</i>				
<i>Helicodicteros_muscivorus</i>				
<i>Hestia_longifolia</i>				
<i>Heteropsis_oblongifolia</i>				
<i>Heteropsis_sp._Barabe_147</i>				
<i>Holochlamys_beccarii</i>				
<i>Homalomena_asmiae</i>				
<i>Homalomena_atrox</i>				
<i>Homalomena_borneensis</i>				
<i>Homalomena_clandestina</i>				
<i>Homalomena_cochinchinensis</i>				
<i>Homalomena_crinipes</i>				
<i>Homalomena_curvata</i>				

<i>Homalomena deblicrista</i>		JQ413313			
<i>Homalomena erythropus</i>		JQ955574			
	DQ870562	DQ866879			
<i>Homalomena expedita</i>		JQ413312,	JX024965		
<i>Homalomena giamensis</i>		JX076778			
		JQ929129	JX024996		
		JQ413339,			
		JX076779,			
		JX076804,			
		JX076803,			
		JX076802,			
		JX076801,			
		JX076780,	JX024974		
<i>Homalomena hanneae</i>		JQ413318			
<i>Homalomena havilandii</i>		JX076781,	JX024985		
		JQ413332			
		JQ413316,			
		JX076806,			
<i>Homalomena humilis</i>		JX076805	JX024969		
		JX076803,			
		JX076782,			
		JQ413331,			
<i>Homalomena insignis</i>		JQ413330	JX024987		GQ220968
		JX076784,			
		JQ413325	JX024980		
			AM920596	AM905774	
<i>Homalomena josephi</i>		JQ955577			
<i>Homalomena magna</i>			GQ434258	GQ436695	
<i>Homalomena matangae</i>					GQ435386
<i>Homalomena occulta</i>					
<i>Homalomena panamensis</i>	DQ870563	DQ866880			
<i>Homalomena philippinensis</i>	DQ870564	DQ866881			
<i>Homalomena picturata</i>	DQ870565	DQ866882			
		JX076785,			
<i>Homalomena punctulata</i>		JQ413337	JX024989		
		JX076786,			
<i>Homalomena rostrata</i>		JX024998	JX024995		
<i>Homalomena rubescens</i>	DQ870566				AY054723
		JX076790,			
		JX076789,			
<i>Homalomena sengkenyang</i>		JX076788,	JX024977		

<i>Homalomena</i> <i>sp.</i> <i>Ar3065</i>					
<i>Homalomena</i> <i>sp.</i> <i>Barabe_151</i>					
			JX076787, JQ413321, JQ413319, JX076794		AY054724
<i>Homalomena</i> <i>sp.</i> <i>Batang_Ai</i>			JX076791, JQ413334	JX024991	
<i>Homalomena</i> <i>sp.</i> <i>Croat_90162</i>	DQ870559				
<i>Homalomena</i> <i>sp.</i> <i>HYC-2012a</i>		JQ955572			
<i>Homalomena</i> <i>sp.</i> <i>HYC-2012b</i>		JQ955575			
<i>Homalomena</i> <i>sp.</i> <i>HYC-2012c</i>		JQ955576			
		JX076795,			
<i>Homalomena</i> <i>sp.</i> <i>KKN-2012</i>		JQ413338	JX024988		
		JQ413315,			
<i>Homalomena</i> <i>sp.</i> <i>Melaka</i>		JX076792	JX024968		
		JX076793,			
<i>Homalomena</i> <i>sp.</i> <i>Pahang</i>		JQ413326	JX024981		
<i>Homalomena</i> <i>sp.</i> <i>SYW-2013</i>		JX076773			
<i>Homalomena</i> <i>sp.</i> <i>Speariae</i>			KC466578	KC466589	
		JQ413322,			
<i>Homalomena</i> <i>symplocarpifolia</i>		JX076808,			
		JX076807	JX024984		
		JX076809,			
<i>Homalomena</i> <i>vagans</i>		JQ413333	JX024990		
		JX076796,			
		JX076797,			
<i>Homalomena</i> <i>vivens</i>		JQ413324,			
		JQ413323	JX024978		
		JX076798,			
<i>Homalomena</i> <i>wallisii</i>		JQ413335	JX024992		
<i>Homalomena</i> <i>wendlandii</i>	DQ870567	DQ866883			
<i>Incarum</i> <i>pavonii</i>			AM920590	AM905768	EF173583
<i>Jasarrum</i> <i>steyermarkii</i>			AM920614	AM905792	
<i>Lagenandra</i> <i>ovata</i>			AM920602	AM905780	AM933330
<i>Landoltia</i> <i>punctata</i>			AY034185	HM850382	
<i>Lasia</i> <i>spinosa</i>			AM920571	AM905749	AM932312
<i>Lasiorapha</i> <i>senegalensis</i>			AM920577	AM905755	
<i>Lemna</i> <i>aequinoctialis</i>			AY034190	AY034228	AM932317
<i>Lemna</i> <i>disperma</i>			AY034198	AY034236	

<i>Lemna ecuadoriensis</i>									
<i>Lemna gibba</i>	AY034193	AY034231							
<i>Lemna japonica</i>	AY034197	AY034235							AY290848
<i>Lemna minor</i>	AY034195	AY034233							
<i>Lemna minuta</i>	AM920552	AM905730							
<i>Lemna obscura</i>	AY034186	AY034224							
<i>Lemna perpusilla</i>	AY034194	AY034232							
<i>Lemna tenera</i>	AY034191	AY034229							
<i>Lemna trisulca</i>	AY034189	AY034227							
<i>Lemna turionifera</i>	AY034199	AY034237							
<i>Lemna valdiviana</i>	AY034192	AY034230							
<i>Lemna yungensis</i>	AY034187	AY034225							
<i>Leucocasia gigantea</i>	AY034188	AY034226							
<i>Lorenzia umbrosa</i>	JQ238893	EU193194	JQ238975						JQ237227
<i>Lysichiton umbrosa</i>	JF803495								
<i>Lysichiton americanus</i>	AM920549	AM905728		KC241917	KC241908	KC241913		DQ400863	
<i>Lysichiton camtschaticensis</i>					AB206655			AY054740	
<i>Mangonia tweediana</i>	AM920588	AM905766						EF173584	
<i>Monstera adansonii</i>	AM920565								
<i>Monstera ada</i>									
<i>nsonii_var._kl</i>									
<i>orzechiana</i>									AY555162
<i>Monstera deliciosa</i>									
<i>Monstera falcifolia</i>								AY398547	
<i>Monstera obliqua</i>								AY398548	
<i>Monstera tenuis</i>									
<i>Montrichardia arborescens</i>	JQ586635								
<i>Nephtytis afzelii</i>	AM920640								
<i>Nephtytis bintuluensis</i>	AM920581								
<i>Nephtytis poissonii</i>	KC466579								
<i>Ooia grabowski</i>									AY290842
<i>Oronitium aquaticum</i>	GQ220903								
<i>Oronitium aquaticum</i>	AM920550								
<i>Pedicularum paiei</i>	AF293753								
<i>Peltandra virginica</i>	AM920555								
<i>Phlodeudron bipinnatifidum</i>	EU886583								
<i>Phlodeudron corcovadense</i>	JX024993								
	KF981850								

<i>Philodendron fragrantissimum</i>			JQ586638
<i>Philodendron lundii</i>			KF971332
<i>Philodendron lundii</i>			KF981857
<i>Philodendron pedatum</i>			KF981852
<i>Philodendron petraeum</i>			KF981853
<i>Philodendron radiatum</i>			JX024994
<i>Philodendron saxicola</i>			KF971327
<i>Philodendron saxicola</i>			KF981854
<i>Philodendron solimoesense</i>			KF981858
<i>Philodendron stenolobum</i>			KF971334
<i>Philodendron stenolobum</i>			KF981859
<i>Philodendron undulatum</i>			KF971328
<i>Philodendron undulatum</i>			KF981855
<i>Philodendron venezuelense</i>			KF971329
<i>Philodendron williamsii</i>			KF971330
<i>Philodendron acutatum</i>	DQ870570		
<i>Philodendron adamantinum</i>	KF895425		
<i>Philodendron angustisectum</i>	DQ870576		
<i>Philodendron anisotomum</i>		DQ866885	
<i>Philodendron barrosocum</i>	DQ870577		
<i>Philodendron billietiae</i>	DQ870578		
<i>Philodendron bipinnatifidum</i>	DQ870614		
<i>Philodendron brasiliense</i>	KF895413		
<i>Philodendron brevispatum</i>	DQ870579		
<i>Philodendron callosum</i>	DQ870580		
<i>Philodendron canifolium</i>	DQ870581		
<i>Philodendron corcovadense</i>	KF895417		
<i>Philodendron crassinervium</i>	DQ870582		
<i>Philodendron dardanianum</i>	KF895411		
<i>Philodendron davidsonii</i>	DQ870583		
<i>Philodendron deltoideum</i>		AM920597	
<i>Philodendron distantilobum</i>	DQ870584		
<i>Philodendron duckei</i>			
<i>Philodendron ecor datum</i>		DQ866890	
<i>Philodendron erubescens</i>	DQ870585		
<i>Philodendron fmdens</i>	DQ870586		

<i>Philodendron_fragrantissimum</i>	DQ870587	
<i>Philodendron_glaziovii</i>	DQ870588	
<i>Philodendron_gloriosum</i>	DQ870589	
<i>Philodendron_goeldii</i>	DQ870571	
<i>Philodendron_grandifolium</i>	DQ870590	
<i>Philodendron_grandipes</i>	DQ870591	
<i>Philodendron_grazielae</i>		
<i>Philodendron_hederaceum</i>	DQ870613	
<i>Philodendron_hederaceum_v</i>		
<i>ar._oxycardiu</i>		
<i>m</i>		
	DQ401355	
<i>Philodendron_heleniae</i>	DQ870592	
<i>Philodendron_hylaeae</i>	DQ870593	
<i>Philodendron_ilsemanii</i>	DQ870594	
<i>Philodendron_imbe</i>	DQ870595	
<i>Philodendron_insigne</i>	DQ870596	
<i>Philodendron_lazorii</i>	DQ870597	
<i>Philodendron_leal-costae</i>	KF895427	
<i>Philodendron_lindenii</i>	DQ870598	
<i>Philodendron_linnaei</i>	DQ870599	
<i>Philodendron_loefgrenii</i>	DQ870600	
<i>Philodendron_longistilum</i>	DQ870601	
<i>Philodendron_lundii</i>	KF895420	
<i>Philodendron_malessevichiae</i>	DQ870602	
<i>Philodendron_manei</i>		DQ866904
<i>Philodendron_megalophyllum</i>		
<i>Philodendron_melinonii</i>	DQ870603	
<i>Philodendron_mello-</i>		
<i>barretoanum</i>	KF895423	
<i>Philodendron_oblongum</i>		KF981851
<i>Philodendron_ornatum</i>	DQ870604	
<i>Philodendron_palmicola</i>	KF895412	
<i>Philodendron_panamense</i>	DQ870605	
<i>Philodendron_panduriforme</i>	DQ870606	
<i>Philodendron_pedatum</i>	DQ870607	

<i>Philodendron_petraceum</i>	KF895422		
<i>Philodendron_pinnatifidum</i>	DQ870608		
<i>Philodendron_placidum</i>	DQ870609		
<i>Philodendron_platypetalum</i>			JQ590343
<i>Philodendron_plerotum</i>		DQ866907	
<i>Philodendron_radialum</i>	DQ870610		
<i>Philodendron_rhodoxis</i>			JQ586644
<i>Philodendron_rudgeanum</i>	DQ870568		
<i>Philodendron_ruizii</i>	DQ870611		
<i>Philodendron_sagittifolium</i>	DQ870612		
<i>Philodendron_saxicola</i>	KF895426		
<i>Philodendron_schottii_subsp._latamancæ</i>	DQ870619		
<i>Philodendron_serpens</i>		DQ866911	
<i>Philodendron_simsii</i>	DQ870615		
<i>Philodendron_smithii</i>	DQ870616		
<i>Philodendron_sodiroi</i>		DQ866913	
<i>Philodendron_solimoesense</i>	DQ870572		
<i>Philodendron_sp._Barabed_Archambault_184</i>			
<i>Philodendron_sp._Croat_84914</i>			
<i>Philodendron_speciosum</i>	DQ870573		
<i>Philodendron_squamiferum</i>	KF895414		
<i>Philodendron_stenolobum</i>	DQ870617		
<i>Philodendron_stenophyllum</i>	KF895424		
<i>Philodendron_sulcatum</i>	DQ870618		
<i>Philodendron_surinamense</i>		JQ589932	
<i>Philodendron_tenuë</i>		DQ866918	
<i>Philodendron_tripartitum</i>	DQ870620		JQ586647
<i>Philodendron_tweedieanum</i>	KF895421		
<i>Philodendron_uliginosum</i>	KF895419		
<i>Philodendron_undulatum</i>	DQ870574		
<i>Philodendron_venezuelense</i>	KF895415		

<i>Philodendron verrucosum</i>	DQ870621	
<i>Philodendron wendlandii</i>	DQ870622	
<i>Philodendron williamsii</i>	KF895416	
<i>Philodendron xanadu</i>	DQ870575	
<i>Philonotion americanum</i>		GQ220908
<i>Phylonotion americanum</i>	JN544445	
<i>Phymatarum borneense</i>		AM920605
<i>Phymatarum borneense</i>	JN544433	
<i>Pichnia disticha</i>		GQ220917
<i>Pinellia cordata</i>		JF828130
<i>Pinellia pedatisecta</i>		AM920629
<i>Pinellia peltata</i>		JF828126
<i>Pinellia polyphylla</i>		JF828125
<i>Pinellia ternata</i>		EU886503
<i>Pinellia tripartita</i>		AB494681
<i>Pinellia cordata</i>	AF469039	
<i>Pinellia fujianensis</i>		JF828129
<i>Pinellia integrifolia</i>	AF469033	
<i>Pinellia pedatisecta</i>	AF469040	
<i>Pinellia peltata</i>	AF469034	
<i>Pinellia polyphylla</i>	AF469035	
<i>Pinellia ternata</i>	AF469036	
<i>Pinellia tripartita</i>	AF469037	
<i>Pinellia yaoluopingensis</i>	AF469038	
<i>Piptospatha truncata</i>		JN177493
<i>Piptospatha burbridgei</i>		GQ220901
<i>Piptospatha elongata</i>		GQ220902
<i>Piptospatha impolita</i>		GQ220905
<i>Piptospatha perakensis</i>		JN177495
<i>Piptospatha ridleyi</i>		AM920603
<i>Piptospatha truncata</i>	JN544443	
<i>Piptospatha viridistigma</i>		GQ220906
<i>Pistia stratiotes</i>		EU886585
<i>Pistia stratiotes</i>	AF168869	
<i>Podolasia stipitata</i>		AM920574
<i>Pothoidium lobbianum</i>		AM920556

<i>Pothos_chinensis</i>						JX894678
<i>Pothos_jungluhnii</i>						JX894679
<i>Pothos_longipes</i>			KF496718			
<i>Pothos_ovatifolius</i>						JX894680
<i>Pothos_scandens</i>		AM920554				JX894681
<i>Pothos_tener</i>						
<i>Protarum_sechellarum</i>						
<i>Pseudodracontium_harmandii</i>		EU886588				
<i>Pseudodracontium_lacourii</i>		AF387425				
<i>Pseudodracontium_lanceolatum</i>		AM920608				
		AF387426				
<i>Pseudohydrosme_gabunensis</i>		AM920582				
<i>Pycnospatha_arietina</i>		AM920573				
<i>Remusatia_hookeriana</i>	AF469041	JF828124	JQ237206			JF828134
<i>Remusatia_pumila</i>	AY081001	JQ238896	JQ237209	JQ238978		AF469027
				JQ238979,		AY081006
<i>Remusatia_vivipara</i>	AY081002	EU886584	KC466593	AY248934		AY081005
<i>Remusatia_yunnanensis</i>	AY081003	JF828114	JQ237214			AY081007
<i>Rhaphidophora_africana</i>					JF828135	AY054736
<i>Rhaphidophora_angulata</i>						AY398512
<i>Rhaphidophora_angustata</i>						AY398514
<i>Rhaphidophora_apiculata</i>						AY398543
<i>Rhaphidophora_australasica</i>						AY398519
<i>Rhaphidophora_beccarii</i>						AY398526
<i>Rhaphidophora_bognerii</i>		KC466581	KC466594			
<i>Rhaphidophora_bonii</i>						AY398517
<i>Rhaphidophora_crassicaulis</i>						AY398529
<i>Rhaphidophora_crassifolia</i>	AM920563	AM905741				AY398511
<i>Rhaphidophora_decursiva</i>						AY398523
<i>Rhaphidophora_elliptifolia</i>						AY398509
<i>Rhaphidophora_foraminifera</i>						AY398542
<i>Rhaphidophora_geniculata</i>						AY398546
<i>Rhaphidophora_glauca</i>						AY398524
<i>Rhaphidophora_hayi</i>						AY398537
<i>Rhaphidophora_hongkongensis</i>						AY398507
<i>Rhaphidophora_hookeri</i>						AY398531

<i>Rhaphidophora_korthalsii</i>				AY398534
<i>Rhaphidophora_latavaginata</i>				AY398532
<i>Rhaphidophora_lobbii</i>				AY398508
<i>Rhaphidophora_luchunensis</i>				AY398533
<i>Rhaphidophora_mainingyi</i>				AY398520
<i>Rhaphidophora_megaphylla</i>				AY398527
<i>Rhaphidophora_megasperma</i>				AY398513
<i>Rhaphidophora_montana</i>				AY398506
<i>Rhaphidophora_nicolsonii</i>				AY398541
<i>Rhaphidophora_novo-guineensis</i>				AY398535
<i>Rhaphidophora_pachyphylla</i>				AY398540
<i>Rhaphidophora_peepla</i>				AY398518
<i>Rhaphidophora_pertusa</i>				AY398539
<i>Rhaphidophora_pertusa</i>				AY398515
<i>Rhaphidophora_puberula</i>				AY398545
<i>Rhaphidophora_schlechteri</i>				AY398536
<i>Rhaphidophora_schottii</i>				AY398516
<i>Rhaphidophora_spathacea</i>				AY398525
<i>Rhaphidophora_spathacea</i>				AY398530
<i>Rhaphidophora_subfalcata</i>				AY398538
<i>Rhaphidophora_syvestris</i>				AY398510
<i>Rhaphidophora_tetrasperma</i>				AY398528
<i>Rhaphidophora_tetrasperma</i>				AY398544
<i>Rhaphidophora_warra</i>				AY398563
<i>Rhodospatha_falconensis</i>				AY398562
<i>Rhodospatha_guanchensis</i>				AY398564
<i>Rhodospatha_oblongata</i>				
<i>Saurumatium_gaoligongense</i>				
<i>Scaphispatha_gracilis</i>				
<i>Schismatoglottis_adoceta</i>	JN544432	EU542587	AM905739	KC460384
<i>Schismatoglottis_asperata</i>		JN570739	AM905793	
<i>Schismatoglottis_bauensis</i>		GQ220909		GQ220979
<i>Schismatoglottis_brevicuspis</i>		GQ220910		GQ220980
<i>Schismatoglottis_calyptrata</i>		GQ220911		GQ220981
<i>Schismatoglottis_ciliata</i>				AY555172
<i>Schismatoglottis_clausula</i>		GQ220913		GQ220982
<i>Schismatoglottis_clausula</i>		GQ220949		GQ221014

<i>Schismatoglottis confinis</i>			GQ220914	GQ220983
<i>Schismatoglottis corneri</i>			GQ220915	GQ220985
<i>Schismatoglottis dulosa</i>			GQ220918	
<i>Schismatoglottis elegans</i>			JN570747	
<i>Schismatoglottis erecta</i>			GQ220919	GQ220990
<i>Schismatoglottis jeponii</i>			GQ220925	GQ220993
<i>Schismatoglottis jiinae</i>			GQ220927	GQ220994
<i>Schismatoglottis josephi</i>	JN544431		GQ220947	GQ221012
<i>Schismatoglottis linae</i>			JN570743	KC454013
<i>Schismatoglottis matangensis</i>			JN570741	
<i>Schismatoglottis mayoana</i>			GQ220929	GQ220997
<i>Schismatoglottis monoplacenta</i>			GQ220931	GQ220999
<i>Schismatoglottis moleyana</i>			GQ220933	GQ221001
<i>Schismatoglottis multiflora</i>			GQ220934	
<i>Schismatoglottis multinervia</i>			GQ220940	GQ221005
<i>Schismatoglottis nervosa</i>	JX076800		JN570742	
<i>Schismatoglottis nicolsonii</i>			GQ220941	GQ221006
<i>Schismatoglottis patentinervia</i>			GQ220942	GQ221007
<i>Schismatoglottis pyrthias</i>			GQ220944	GQ221008
<i>Schismatoglottis sarikensis</i>	JN544434			KC454015
<i>Schismatoglottis simonii</i>			JN570740	JN177488
<i>Schismatoglottis sp._AR-1930</i>		GQ220946		
<i>Schismatoglottis sp._AR-2078</i>			JN570744	
<i>Schismatoglottis sp._AR-2482</i>			JN570749	
<i>Schismatoglottis sp._AR108</i>			JN570745	
<i>Schismatoglottis sp._AR134</i>			GQ220945	
<i>Schismatoglottis sp._AR134</i>			GQ220948	
<i>Schismatoglottis sp._B_SLL-2011</i>	JN544424	KC454001		
<i>Schismatoglottis sp._SLL-2011c</i>			JN544429	
<i>Schismatoglottis sp._SYW-2010f</i>			JN544430	
<i>Schismatoglottis tecturata</i>			GQ220950	GQ221015

<i>Schismatoglottis tessellata</i>	JN570748								
<i>Schismatoglottis trifasciata</i>	AM920604	AM905782							
<i>Schismatoglottis turbata</i>	JN570750								
<i>Schismatoglottis ulsarikeniensis</i> ^s	JN570746								
<i>Schismatoglottis viridissima</i>	GQ220952							GQ221017	
<i>Schottarella muricata</i>	GQ220912							GQ221009	
<i>Scindapsus hederaceus</i>	AM920564	AM905742						AY398556	
<i>Scindapsus longistipitatus</i>								AY398555	
<i>Scindapsus pictus</i>								AY398554	
<i>Spathantheum orbignyianum</i>	EF173550							EF173585	
<i>Spathicarpa gardneri</i>	EF173552								
<i>Spathicarpa hastifolia</i>	AM920594	AM905772							EF173589
<i>Spathicarpa lanceolata</i>	EF173556								
<i>Spathicarpa sagittifolia</i>	EF173558								
<i>Spathiphyllum canniifolium</i>									AY398570
<i>Spathiphyllum cochlearispathu</i> <i>m</i>	AF542575							AY398569	
<i>Spathiphyllum floribundum</i>								AY555159	
<i>Spathiphyllum patinii</i>								AY555158	
<i>Spathiphyllum phrynifolium</i>									
<i>Spathiphyllum wallisii</i>	AF207023	AM920559	JQ590355		KC241920	KC241911	KC241916	AY054738	
<i>Spirodela intermedia</i>		AY034183	U68092				FN675813		
<i>Spirodela polytricha</i>	KC573839	AM920553	AM905731						
<i>Stenospermation multiovulatu</i> <i>m</i>					KC241919,	KC241910	KC241915	AY398566	
<i>Stenospermation popayanense</i>					KC241919			AY054737	
<i>Stenospermation ullei</i>								AY398565	
<i>Stenodnera assamica</i>	AM920561	AM905738							
<i>Stenodnera colocasiifolia</i>	JQ238898				EF517224				
<i>Stenodnera discolor</i>	AM920623	AM905801							
<i>Stenodnera henryana</i>	EU886586				EF517221		JF828140	JQ237247	
<i>Stenodnera kerrii</i>	AF469042							AF469029	
<i>Stylochaeton bogneri</i>	JQ238899	KC466595			EF517223				
<i>Symplocarpus foetidus</i>	AM920598	AM905776							
<i>Symplocarpus nabekuraensis</i>	FJ874946	AM920551	L10247			AB206656	KC241912	AY054741	
						AB206658			

<i>Symplocarpus nipponicus</i>							AB206657	DQ400870
<i>Symplocarpus renifolius</i>								DQ400867
<i>Synandropadix vermitoxicus</i>								EF173594
<i>Syngonium angustatum</i>								AY555177
<i>Syngonium auritum</i>								
<i>Syngonium podophyllum</i>								
<i>Syngonium wendlandii</i>								
<i>Taccarum crassispalum</i>								
<i>Taccarum warningii</i>								
<i>Taccarum weddellianum</i>								
<i>Theriotophonum dalzielii</i>								
<i>Theriotophonum infaustum</i>								
<i>Typhonium adnatum</i>								
<i>Typhonium albidinervum</i>								
<i>Typhonium alismifolium</i>								
<i>Typhonium angustilobum</i>								
<i>Typhonium bachmaense</i>								
<i>Typhonium baoshanense</i>								
<i>Typhonium blumei</i>								
<i>Typhonium brevipes</i>								
<i>Typhonium brownii</i>								
<i>Typhonium bulbiferum</i>								
<i>Typhonium calcicola</i>								
<i>Typhonium cincinnatum</i>								
<i>Typhonium cordifolium</i>								
<i>Typhonium digitatum</i>								
<i>Typhonium divaricatum</i>								
<i>Typhonium diversifolium</i>								
<i>Typhonium echinulatum</i>								
<i>Typhonium eliosorum</i>								
<i>Typhonium filiforme</i>								
<i>Typhonium flagelliforme</i>								
<i>Typhonium gallowayi</i>								
<i>Typhonium gaoligongense</i>								

<i>Typhonium_tentaculatum</i>		EU886543	EU886612	
<i>Typhonium_trilobatum</i>	KC478077	EU886571	AY248941	
<i>Typhonium_tubispalum</i>		EU886574	KC460382	AB494591
<i>Typhonium_varians</i>		EU886575	KC460383	
		KM093848		
<i>Typhonium_venosum</i>		EU886544	AF497113	AB494602
<i>Typhonium_violifolium</i>		EU886562	EU886611	
<i>Typhonium_wilbertii</i>		GU255976	KF496356	
<i>Typhonodorum_lindleyanum</i>		EU886578	AM905814	EU886627
<i>Ulearum_sagittatum</i>		EU542590	AM905794	
<i>Urospatha_sagittifolia</i>		AM920570	AM905748	
<i>Wolffia_angusta</i>		AY034215	AY034253	
<i>Wolffia_arhiza</i>		AY034216	AY034254	
<i>Wolffia_australiana</i>		AY034213	AY034251	
<i>Wolffia_borealis</i>		AY034212	AY034250	
<i>Wolffia_brasiliensis</i>		AY034210	AY034248	
<i>Wolffia_columbiana</i>		AY034217	AY034255	
<i>Wolffia_cylindracea</i>		AY034218	AY034256	
<i>Wolffia_elongata</i>		AY034220	AY034258	
<i>Wolffia_globosa</i>		AY034219	AY034257	
<i>Wolffia_microscopica</i>		AY034211	AY034249	
<i>Wolffia_neglecta</i>		AY034214	AY034252	
<i>Wolffia_caudata</i>		AY034206	AY034244	
<i>Wolffia_denticulata</i>		AY034209	AY034247	
<i>Wolffia_gladinata</i>		AY034205	AY034243	
<i>Wolffia_hyalina</i>		AY034202	AY034240	AY290851
<i>Wolffia_lingulata</i>		AY034203	AY034241	
<i>Wolffia_neotropica</i>		AY034208	AY034246	
<i>Wolffia_oblonga</i>		AY034204	AY034242	
<i>Wolffia_repanda</i>			AY034239	
<i>Wolffia_rotunda</i>		AY034200	AY034238	
<i>Wolffia_welwitschii</i>		AY034207	AY034245	
<i>Xanthosoma_brasiliense</i>		EU542591		
<i>Xanthosoma_helleborifolium</i>		AM920612	AM905790	
<i>Xanthosoma_majaffa</i>		DQ401357	AJ007543	

<i>Xanthosoma_mexicanum</i>	JQ586654	JQ590366		
<i>Xanthosoma_sagittifolium</i>	EU886500	L10246	AY248944	AY555175
<i>Xanthosoma_violaceum</i>	EU193571	EU193180		
<i>Xanthosoma_wendlandii</i>	JQ586659	JQ590370		
<i>Zamioculcas_zamiifolia</i>	AM920600	AM905778		AY054725
<i>Zantedeschia_aethiopica</i>	JQ412304	AF065474		
<i>Zantedeschia_albomaculata</i>	AM920584	AM905762		
<i>Zomicarpa_steigeriana</i>	EU542592			
<i>Zomicarpella_amazonica</i>	EU542593	AM905796		AM932346

TABLE S2. Clade set and species numbers used in the BAMM analyses.

Clade set 1	Code	Species number	Clade set 2	Code	Species number
Alismataceae	Ali	111	Aglaonemateae	Agl	24
Aponogetonaceae	Apo	50	Alismataceae	Ali	111
Aroideae	Aro	1889	<i>Anthurium</i>	Ant	950
Butomaceae	But	1	<i>Anubias</i>	Anu	8
Cymodoceaceae	Cym	20	Aponogetonaceae	Apo	50
Gymnostachioideae	Gym	1	Butomaceae	But	1
Hydrocharitaceae	Hyd	126	Caladieae	Cal	169
Juncaginaceae	Jun	17	<i>Calla</i>	Caa	1
Lasioideae	Las	58	<i>Calloopsis</i>	Cll	1
Lemnoideae	Lem	38	Culcasieae	Cul	38
Maundiaceae	Mau	1	Cymodoceaceae	Cym	20
Monsteroideae	Mon	358	<i>Heteropsis</i> clade	Het	97
Orontioideae	Oro	8	Hydrocharitaceae	Hyd	126
Posidoniaceae	Pos	5	Juncaginaceae	Jun	17
Potamogetonaceae	Pot	115	Lasioideae	Las	58
Pothoideae	Poh	1010	Lemnoideae	Lem	38
Scheuchzeriaceae	Sch	1	Maundiaceae	Mau	1
Tofieldiaceae	Tof	28	<i>Rhaphidophora</i> clade	Mon	211
Zamioculcoideae	Zam	32	<i>Montrichardia</i>	Mot	2
Zosteraceae	Zos	14	Nephtytideae	Nep	13
			<i>Philodendron</i> clade	Phi	572
			<i>Philonotum</i> clade	Phi	272
			Posidoniaceae	Pos	5
			Potamogetonaceae	Pot	115
			<i>Pothos</i> clade	Poh	60
			Protarum clade	Pro	472
			Protoaraceae	Pra	9
			Scheuchzeriaceae	Sch	1
			Spathicarpeae	Spa	92
			Spathiphyllaeae	Spt	50
			Thomsonieae	Tho	197
			<i>Thyphonodorum</i> clade	Thy	20
			Tofieldiaceae	Tof	28
			Zamioculcoideae	Zam	32
			<i>Zantedeschia</i> clade	Zan	8
			Zosteraceae	Zos	14

TABLE S3. Clade set allocation for each taxon in the two-gene dataset, with the species fraction from the total number of species in the clade that they represent. Ten-gene dataset follows the same allocation for each genus.

Species	Clade set 1	Sp fraction	Clade set 2	Sp fraction
<i>Acorus calamus</i>	Aco	0.5	Aco	0.5
<i>Aglaodorum griffithii</i>	Aro	0.050291159	Agl	0.125
<i>Aglaonema modestum</i>	Aro	0.050291159	Agl	0.125
<i>Nephtytis bintuluensis</i>	Aro	0.050291159	Agl	0.125
<i>Luronium natans</i>	Ali	0.14159292	Ali	0.14159292
<i>Damasonium alisma</i>	Ali	0.14159292	Ali	0.14159292
<i>Baldellia ranunculoides</i>	Ali	0.14159292	Ali	0.14159292
<i>Alisma canaliculatum</i>	Ali	0.14159292	Ali	0.14159292
<i>Alisma plantagoaquatica</i>	Ali	0.14159292	Ali	0.14159292
<i>Limnophyton LYC2011</i>	Ali	0.14159292	Ali	0.14159292
<i>Hydrocleys nymphoides</i>	Ali	0.14159292	Ali	0.14159292
<i>Butomopsis latifolia</i>	Ali	0.14159292	Ali	0.14159292
<i>Limnocharis flava</i>	Ali	0.14159292	Ali	0.14159292
<i>Ranalisma rostratum</i>	Ali	0.14159292	Ali	0.14159292
<i>Caldesia parnassifolia</i>	Ali	0.14159292	Ali	0.14159292
<i>Echinodorus cordifolius</i>	Ali	0.14159292	Ali	0.14159292
<i>Helanthium parvulum</i>	Ali	0.14159292	Ali	0.14159292
<i>Sagittaria latifolia</i>	Ali	0.14159292	Ali	0.14159292
<i>Wiesneria triandra</i>	Ali	0.14159292	Ali	0.14159292
<i>Astonia australiensis</i>	Ali	0.14159292	Ali	0.14159292
<i>Anthurium acaule</i>	Poh	0.010891089	Ant	0.008421053
<i>Anthurium andreanum</i>	Poh	0.010891089	Ant	0.008421053
<i>Anthurium bakeri</i>	Poh	0.010891089	Ant	0.008421053
<i>Anthurium brenesii</i>	Poh	0.010891089	Ant	0.008421053
<i>Anthurium interruptum</i>	Poh	0.010891089	Ant	0.008421053
<i>Anthurium ravenii</i>	Poh	0.010891089	Ant	0.008421053
<i>Anthurium scandens</i>	Poh	0.010891089	Ant	0.008421053
<i>Anthurium spectabile</i>	Poh	0.010891089	Ant	0.008421053
<i>Anubias barteri</i>	Aro	0.050291159	Anu	0.25
<i>Anubias heterophylla</i>	Aro	0.050291159	Anu	0.25
<i>Aponogeton fenestralis</i>	Apo	0.02	Apo	0.02
<i>Butomus umbellatus</i>	But	1	But	1
<i>Caladium bicolor</i>	Aro	0.050291159	Cal	0.059171598
<i>Chlorospatha Chase1821912</i>	Aro	0.050291159	Cal	0.059171598
<i>Filarum manserichense</i>	Aro	0.050291159	Cal	0.059171598
<i>Hapaline benthamiana</i>	Aro	0.050291159	Cal	0.059171598

<i>Jasarum steyermarkii</i>	Aro	0.050291159	Cal	0.059171598
<i>Scaphispatha gracilis</i>	Aro	0.050291159	Cal	0.059171598
<i>Syngonium AZ128</i>	Aro	0.050291159	Cal	0.059171598
<i>Ulearum sagittatum</i>	Aro	0.050291159	Cal	0.059171598
<i>Xanthosoma sagittifolium</i>	Aro	0.050291159	Cal	0.059171598
<i>Zomicarpella amazonica</i>	Aro	0.050291159	Cal	0.059171598
<i>Calla palustris</i>	Aro	0.050291159	Caa	1
<i>Calloopsis volkensii</i>	Aro	0.050291159	Cll	1
<i>Cercestis mirabilis</i>	Aro	0.050291159	Cul	0.052631579
<i>Culcasia liberica</i>	Aro	0.050291159	Cul	0.052631579
<i>Ruppia megacarpa</i>	Cym	0.285714286	Cym	0.285714286
<i>Halodule uninervis</i>	Cym	0.285714286	Cym	0.285714286
<i>Cymodocea serrulata</i>	Cym	0.285714286	Cym	0.285714286
<i>Syringodium filiforme</i>	Cym	0.285714286	Cym	0.285714286
<i>Thalassodendron ciliatum</i>	Cym	0.285714286	Cym	0.285714286
<i>Amphibolis antarctica</i>	Cym	0.285714286	Cym	0.285714286
<i>Alloschemone occidentalis</i>	Mon	0.097765363	Het	0.092783505
<i>Heteropsis oblongifolia</i>	Mon	0.097765363	Het	0.092783505
<i>Heteropsis spruceana</i>	Mon	0.097765363	Het	0.092783505
<i>Rhodospatha AZ125</i>	Mon	0.097765363	Het	0.092783505
<i>Rhodospatha AZ82</i>	Mon	0.097765363	Het	0.092783505
<i>Rhodospatha wendlandii</i>	Mon	0.097765363	Het	0.092783505
<i>Stenospermation AZ129</i>	Mon	0.097765363	Het	0.092783505
<i>Stenospermation AZ80</i>	Mon	0.097765363	Het	0.092783505
<i>Stenospermation escobariae</i>	Mon	0.097765363	Het	0.092783505
<i>Stratiotes aloides</i>	Hyd	0.129032258	Hyd	0.129032258
<i>Lagarosiphon muscoides</i>	Hyd	0.129032258	Hyd	0.129032258
<i>Ottelia acuminata</i>	Hyd	0.129032258	Hyd	0.129032258
<i>Blyxa echinosperma</i>	Hyd	0.129032258	Hyd	0.129032258
<i>Apalanthe granatensis</i>	Hyd	0.129032258	Hyd	0.129032258
<i>Elodea nuttallii</i>	Hyd	0.129032258	Hyd	0.129032258
<i>Egeria densa</i>	Hyd	0.129032258	Hyd	0.129032258
<i>Hydrilla verticillata</i>	Hyd	0.129032258	Hyd	0.129032258
<i>Hydrocharis dubia</i>	Hyd	0.129032258	Hyd	0.129032258
<i>Limnobium laevigatum</i>	Hyd	0.129032258	Hyd	0.129032258
<i>Najas flexilis</i>	Hyd	0.129032258	Hyd	0.129032258
<i>Vallisneria asiatica</i>	Hyd	0.129032258	Hyd	0.129032258
<i>Nechamandra alternifolia</i>	Hyd	0.129032258	Hyd	0.129032258
<i>Halophila ovalis</i>	Hyd	0.129032258	Hyd	0.129032258
<i>Enhalus acoroides</i>	Hyd	0.129032258	Hyd	0.129032258
<i>Thalassia hemprichii</i>	Hyd	0.129032258	Hyd	0.129032258
<i>Tetroncium magellanicum</i>	Jun	0.235294118	Jun	0.235294118
<i>Cycnogeton rheophilum</i>	Jun	0.235294118	Jun	0.235294118

<i>Triglochin maritima</i>	Jun	0.235294118	Jun	0.235294118
<i>Anaphyllopsis americana</i>	Las	0.101694915	Las	0.101694915
<i>Cyrtosperma macrothum</i>	Las	0.101694915	Las	0.101694915
<i>Dracontioides desciscens</i>	Las	0.101694915	Las	0.101694915
<i>Dracontium polyphyllum</i>	Las	0.101694915	Las	0.101694915
<i>Lasia spinosa</i>	Las	0.101694915	Las	0.101694915
<i>Lasimorpha senegalensis</i>	Las	0.101694915	Las	0.101694915
<i>Podolasia stipitata</i>	Las	0.101694915	Las	0.101694915
<i>Pycnospatha arietina</i>	Las	0.101694915	Las	0.101694915
<i>Urospatha sagittifolia</i>	Las	0.101694915	Las	0.101694915
<i>Landoltia punctata</i>	Lem	0.131578947	Lem	0.131578947
<i>Lemna minor</i>	Lem	0.131578947	Lem	0.131578947
<i>Spirodela polyrhiza</i>	Lem	0.131578947	Lem	0.131578947
<i>Wolffia columbiana</i>	Lem	0.131578947	Lem	0.131578947
<i>Wolffiella gladiata</i>	Lem	0.131578947	Lem	0.131578947
<i>Maundia triglochinoides</i>	Mau	1	Mau	1
<i>Amydrium hainanense</i>	Mon	0.097765363	Mon	0.104265403
<i>Amydrium humile</i>	Mon	0.097765363	Mon	0.104265403
<i>Amydrium medium</i>	Mon	0.097765363	Mon	0.104265403
<i>Anadendrum microstachyum</i>	Mon	0.097765363	Mon	0.104265403
<i>Epipremnum aureum</i>	Mon	0.097765363	Mon	0.104265403
<i>Epipremnum falcifolium</i>	Mon	0.097765363	Mon	0.104265403
<i>Epipremnum papuanum</i>	Mon	0.097765363	Mon	0.104265403
<i>Epipremnum pinnatum</i>	Mon	0.097765363	Mon	0.104265403
<i>Monstera adansonii</i>	Mon	0.097765363	Mon	0.104265403
<i>Monstera deliciosa</i>	Mon	0.097765363	Mon	0.104265403
<i>Monstera dubia</i>	Mon	0.097765363	Mon	0.104265403
<i>Monstera tuberculata</i>	Mon	0.097765363	Mon	0.104265403
<i>Monstera xanthospatha</i>	Mon	0.097765363	Mon	0.104265403
<i>Rhaphidophora bognerii</i>	Mon	0.097765363	Mon	0.104265403
<i>Rhaphidophora crassifolia</i>	Mon	0.097765363	Mon	0.104265403
<i>Rhaphidophora decursiva</i>	Mon	0.097765363	Mon	0.104265403
<i>Rhaphidophora hongkongensis</i>	Mon	0.097765363	Mon	0.104265403
<i>Rhaphidophora lobbii</i>	Mon	0.097765363	Mon	0.104265403
<i>Rhaphidophora megaphylla</i>	Mon	0.097765363	Mon	0.104265403
<i>Rhaphidophora spuria</i>	Mon	0.097765363	Mon	0.104265403
<i>Rhaphidophora sylvestris</i>	Mon	0.097765363	Mon	0.104265403
<i>Scindapsus hederaceus</i>	Mon	0.097765363	Mon	0.104265403
<i>Montrichardia arborescens</i>	Aro	0.050291159	Mot	0.5
<i>Anchomanes difformis</i>	Aro	0.050291159	Nep	0.230769231
<i>Nephtytis afzelii</i>	Aro	0.050291159	Nep	0.230769231
<i>Pseudohydrosme gabunensis</i>	Aro	0.050291159	Nep	0.230769231
<i>Japonolirion osense</i>	Pet	0.5	Pet	0.5

<i>Petrosavia sakuraii</i>	Pet	0.5	Pet	0.5
<i>Furtadoa mixta</i>	Aro	0.050291159	Phi	0.01048951
<i>Homalomena magna</i>	Aro	0.050291159	Phi	0.01048951
<i>Homalomena speariae</i>	Aro	0.050291159	Phi	0.01048951
<i>Philodendron bipinnatifidum</i>	Aro	0.050291159	Phi	0.01048951
<i>Philodendron deltoideum</i>	Aro	0.050291159	Phi	0.01048951
<i>Philodendron fragrantissimum</i>	Aro	0.050291159	Phi	0.01048951
<i>Apoballis acuminatissima</i>	Aro	0.050291159	Phl	0.055147059
<i>Aridarum nicolsonii</i>	Aro	0.050291159	Phl	0.055147059
<i>Bakoa lucens</i>	Aro	0.050291159	Phl	0.055147059
<i>Bucephalandra motleyana</i>	Aro	0.050291159	Phl	0.055147059
<i>Cryptocoryne ferruginea</i>	Aro	0.050291159	Phl	0.055147059
<i>Cryptocoryne uenoi</i>	Aro	0.050291159	Phl	0.055147059
<i>Hestia longifolia</i>	Aro	0.050291159	Phl	0.055147059
<i>Lagenandra ovata</i>	Aro	0.050291159	Phl	0.055147059
<i>Ooia grabowskii</i>	Aro	0.050291159	Phl	0.055147059
<i>Philonotion americanum</i>	Aro	0.050291159	Phl	0.055147059
<i>Phymatarum borneense</i>	Aro	0.050291159	Phl	0.055147059
<i>Pichinia disticha</i>	Aro	0.050291159	Phl	0.055147059
<i>Piptospatha ridleyi</i>	Aro	0.050291159	Phl	0.055147059
<i>Schismatoglottis trifasciata</i>	Aro	0.050291159	Phl	0.055147059
<i>Schottariella mirifica</i>	Aro	0.050291159	Phl	0.055147059
<i>Posidonia oceanica</i>	Pos	0.2	Pos	0.2
<i>Stuckenia pectinata</i>	Pot	0.051282051	Pot	0.051282051
<i>Potamogeton distinctus</i>	Pot	0.051282051	Pot	0.051282051
<i>Potamogeton perfoliatus</i>	Pot	0.051282051	Pot	0.051282051
<i>Groenlandia densa</i>	Pot	0.051282051	Pot	0.051282051
<i>Lepilaena australis</i>	Pot	0.051282051	Pot	0.051282051
<i>Zannichellia palustris</i>	Pot	0.051282051	Pot	0.051282051
<i>Pedicellarum paiei</i>	Poh	0.010891089	Poh	0.05
<i>Pothoidium lobbianum</i>	Poh	0.010891089	Poh	0.05
<i>Pothos scandens</i>	Poh	0.010891089	Poh	0.05
<i>Alocasia macrorrhizos</i>	Aro	0.050291159	Pro	0.05720339
<i>Alocasia odora</i>	Aro	0.050291159	Pro	0.05720339
<i>Ariopsis peltata</i>	Aro	0.050291159	Pro	0.05720339
<i>Arisaema dracontium</i>	Aro	0.050291159	Pro	0.05720339
<i>Arisaema speciosum</i>	Aro	0.050291159	Pro	0.05720339
<i>Arisaema tortuosum</i>	Aro	0.050291159	Pro	0.05720339
<i>Arum cylindraceum</i>	Aro	0.050291159	Pro	0.05720339
<i>Arum pictum</i>	Aro	0.050291159	Pro	0.05720339
<i>Biarum tenuifolium</i>	Aro	0.050291159	Pro	0.05720339
<i>Colocasia esculenta</i>	Aro	0.050291159	Pro	0.05720339
<i>Colocasia fallax</i>	Aro	0.050291159	Pro	0.05720339

<i>Dracunculus vulgaris</i>	Aro	0.050291159	Pro	0.05720339
<i>Eminium spiculatum</i>	Aro	0.050291159	Pro	0.05720339
<i>Englerarum hypnosum</i>	Aro	0.050291159	Pro	0.05720339
<i>Helicodiceros muscivorus</i>	Aro	0.050291159	Pro	0.05720339
<i>Lazarum brownii</i>	Aro	0.050291159	Pro	0.05720339
<i>Leucocasia gigantea</i>	Aro	0.050291159	Pro	0.05720339
<i>Lorenzia umbrosa</i>	Aro	0.050291159	Pro	0.05720339
<i>Pinellia pedatisecta</i>	Aro	0.050291159	Pro	0.05720339
<i>Pistia stratiotes</i>	Aro	0.050291159	Pro	0.05720339
<i>Protarum sechellarum</i>	Aro	0.050291159	Pro	0.05720339
<i>Remusatia pumila</i>	Aro	0.050291159	Pro	0.05720339
<i>Remusatia vivipara</i>	Aro	0.050291159	Pro	0.05720339
<i>Sauromatum venosum</i>	Aro	0.050291159	Pro	0.05720339
<i>Steudnera kerrii</i>	Aro	0.050291159	Pro	0.05720339
<i>Theriophonum dalzellii</i>	Aro	0.050291159	Pro	0.05720339
<i>Typhonium blumei</i>	Aro	0.050291159	Pro	0.05720339
<i>Gymnostachys anceps</i>	Gym	1	Pra	0.444444444
<i>Lysichiton americanus</i>	Oro	0.375	Pra	0.444444444
<i>Orontium aquaticum</i>	Oro	0.375	Pra	0.444444444
<i>Symplocarpus foetidus</i>	Oro	0.375	Pra	0.444444444
<i>Scheuchzeria palustris</i>	Sch	1	Sch	1
<i>Asterostigma lividum</i>	Aro	0.050291159	Spa	0.130434783
<i>Bognera recondita</i>	Aro	0.050291159	Spa	0.130434783
<i>Croatiella integrifolia</i>	Aro	0.050291159	Spa	0.130434783
<i>Dieffenbachia aglaonematifolia</i>	Aro	0.050291159	Spa	0.130434783
<i>Gearum brasiliense</i>	Aro	0.050291159	Spa	0.130434783
<i>Gorgonidium intermedium</i>	Aro	0.050291159	Spa	0.130434783
<i>Incarum pavonii</i>	Aro	0.050291159	Spa	0.130434783
<i>Mangonia tweedieana</i>	Aro	0.050291159	Spa	0.130434783
<i>Spathanthemum orbignyanum</i>	Aro	0.050291159	Spa	0.130434783
<i>Spathicarpa hastifolia</i>	Aro	0.050291159	Spa	0.130434783
<i>Synandropadix vermitoxicus</i>	Aro	0.050291159	Spa	0.130434783
<i>Taccarum weddellianum</i>	Aro	0.050291159	Spa	0.130434783
<i>Holochlamys beccarii</i>	Mon	0.097765363	Spt	0.08
<i>Spathiphyllum cannifolium</i>	Mon	0.097765363	Spt	0.08
<i>Spathiphyllum commutatum</i>	Mon	0.097765363	Spt	0.08
<i>Spathiphyllum wallisii</i>	Mon	0.097765363	Spt	0.08
<i>Amorphophallus angolensis</i>	Aro	0.050291159	Tho	0.015228426
<i>Amorphophallus paeoniifolius</i>	Aro	0.050291159	Tho	0.015228426
<i>Pseudodracontium lanceolatum</i>	Aro	0.050291159	Tho	0.015228426
<i>Ambrosina bassii</i>	Aro	0.050291159	Thy	0.35
<i>Arisarum vulgare</i>	Aro	0.050291159	Thy	0.35
<i>Arophyton buchettii</i>	Aro	0.050291159	Thy	0.35

<i>Carlephyton glaucophyllum</i>	Aro	0.050291159	Thy	0.35
<i>Colletogyne perrieri</i>	Aro	0.050291159	Thy	0.35
<i>Peltandra virginica</i>	Aro	0.050291159	Thy	0.35
<i>Typhonodorum lindleyanum</i>	Aro	0.050291159	Thy	0.35
<i>Isidrogalvia sessiliflora</i>	Tof	0.178571429	Tof	0.178571429
<i>Pleea tenuifolia</i>	Tof	0.178571429	Tof	0.178571429
<i>Tofieldia coccinea</i>	Tof	0.178571429	Tof	0.178571429
<i>Tofieldia pusilla</i>	Tof	0.178571429	Tof	0.178571429
<i>Triantha glutinosa</i>	Tof	0.178571429	Tof	0.178571429
<i>Gonatopus angustus</i>	Zam	0.09375	Zam	0.09375
<i>Stylochaeton bogneri</i>	Zam	0.09375	Zam	0.09375
<i>Zamioculcas zamiifolia</i>	Zam	0.09375	Zam	0.09375
<i>Zantedeschia aethiopica</i>	Aro	0.050291159	Zan	0.25
<i>Zantedeschia albomaculata</i>	Aro	0.050291159	Zan	0.25
<i>Phyllospadix iwatensis</i>	Zos	0.136363636	Zos	0.136363636
<i>Zostera marina</i>	Zos	0.136363636	Zos	0.136363636
<i>Zostera tasmanica</i>	Zos	0.136363636	Zos	0.136363636

TABLE S3. Morphological and ecological traits used in the ancestral state reconstructions. NA represents missing information.

Genus	Life form1	Life form2	Life Form3	Distribution	Life cycle	Flower		Perianth	Pollen_exine	Pollen_type
						sexuality				
<i>Acorus</i>	Terrestrial	Rhizomato us	Terrestrial	Temperate Zones Tropics-	Evergreen	bisexual		present	present	monosulcate
<i>Alisma</i>	Rooted_aqu atic	Rhizomato us	Aquatic	Temperate Zones	Evergreen	bisexual		present	present	pantoporate
<i>Astonia</i>	Rooted_aqu atic	?	Aquatic	Tropics Temperate Zones	NA	bisexual		present	NA	NA
<i>Baldellia</i>	Rooted_aqu atic	Rhizomato us	Aquatic		NA	bisexual		present	present	pantoporate
<i>Buonomopsis</i>	Rooted_aqu atic	Caullescent	Aquatic	Tropics Tropics- Temperate Zones	Evergreen	bisexual		present	present	pantoporate
<i>Caldesia</i>	Rooted_aqu atic	Rhizomato us	Aquatic		NA	bisexual		present	present	pantoporate
<i>Damasonium</i>	Rooted_aqu atic	Rhizomato us	Aquatic	Temperate Zones	NA	bisexual		present	present	pantoporate
<i>Echinodorus</i>	Rooted_aqu atic	Rhizomato us	Aquatic	Tropics Tropics- Temperate Zones	NA	bisexual		present	present	pantoporate
<i>Helanthium</i>	Rooted_aqu atic		Aquatic		NA	bisexual		present	NA	NA
<i>Hydrocleys</i>	Rooted_aqu atic	Caullescent Rhizomato	Aquatic	Tropics	NA	bisexual		present	present	pantoporate
<i>Limncharis</i>	Rooted_aqu atic	us Rhizomato	Aquatic	Tropics	Evergreen	bisexual		present	present	pantoporate
<i>Linnophyton</i>	Rooted_aqu atic	us Rhizomato	Aquatic	Tropics Temperate Zones	NA	NA		present	present	pantoporate
<i>Luronium</i>	Rooted_aqu atic	us Rhizomato	Aquatic		NA	bisexual		present	present	pantoporate
<i>Randalsma</i>	Rooted_aqu atic	Rhizomato us	Aquatic	Tropics	Evergreen	bisexual		present	present	pantoporate
<i>Sagittaria</i>	Rooted_aqu atic	Rhizomato	Aquatic	Tropics-	NA	unisexual		present	present	pantoporate

	atic	us	Temperate Zones					
<i>Wiesneria</i>	Rooted_aquatic	Rhizomatous	Aquatic	Tropics	NA	bisexual	present	present
<i>Aponogeton</i>	Rooted_aquatic	Tuberous	Aquatic	Tropics	NA	unisexual	present	present
<i>Adelonema</i>	Terrestrial	?	Terrestrial	Tropics	Evergreen	unisexual	absent	reduced
<i>Aglaodorum</i>	Rooted_aquatic	Rhizomatous	Aquatic	Tropics	Evergreen	unisexual	absent	reduced
<i>Aglaonema</i>	Terrestrial	Caullescent	Terrestrial	Tropics	Evergreen	unisexual	absent	reduced
<i>Alocasia</i>	Terrestrial	us	Terrestrial	Tropics	Evergreen	unisexual	absent	reduced
<i>Ambrosina</i>	Terrestrial	Tuberous	Terrestrial	Tropics	Evergreen	unisexual	absent	reduced
<i>Amorphophallus</i>	Terrestrial	Tuberous	Terrestrial	Tropics	Seasonall	unisexual	absent	reduced
<i>Anchomanes</i>	Terrestrial	Rhizomatous	Terrestrial	Tropics	Seasonall	unisexual	absent	reduced
<i>Anubias</i>	Rooted_aquatic	Rhizomatous	Aquatic	Tropics	Seasonall	unisexual	absent	reduced
<i>Apoballis</i>	Terrestrial	us	Terrestrial	Tropics	Evergreen	unisexual	absent	reduced
<i>Aridarum</i>	Terrestrial	Caullescent	Terrestrial	Tropics	Evergreen	unisexual	absent	reduced
<i>Ariopsis</i>	Terrestrial	Tuberous	Terrestrial	Tropics	Seasonall	unisexual	absent	reduced
<i>Arisaema</i>	Terrestrial	Tuberous	Terrestrial	Tropics	Seasonall	unisexual	absent	reduced
<i>Arisarum</i>	Terrestrial	Tuberous	Terrestrial	Tropics	Seasonall	unisexual	absent	reduced
<i>Arophyton</i>	Terrestrial	Rhizomatous	Terrestrial	Tropics	Seasonall	unisexual	absent	reduced
<i>Arum</i>	Terrestrial	Tuberous	Terrestrial	Tropics	Seasonall	unisexual	absent	reduced
<i>Asterostigma</i>	Terrestrial	Tuberous	Terrestrial	Tropics	Seasonall	unisexual	absent	reduced
<i>Bakoa</i>	Terrestrial	Rhizomatous	Terrestrial	Tropics	Evergreen	unisexual	absent	reduced

<i>Furtadaa</i>	Terrestrial	Caulscent	Terrestrial	Tropics	Evergreen Seasonall	unisexual	absent	reduced	inaperturate
<i>Gearum</i>	Terrestrial	Tuberous	Terrestrial	Tropics	y dormant Seasonall	unisexual	absent	reduced	inaperturate
<i>Gorgonidium</i>	Terrestrial	Tuberous	Terrestrial	Tropics	y dormant Seasonall	unisexual	absent	reduced	inaperturate
<i>Hapaline</i>	Terrestrial	Tuberous	Terrestrial	Tropics Temperate Zones	y dormant Seasonall	unisexual	absent	reduced	inaperturate
<i>Helicodiceros</i>	Terrestrial	Tuberous Rhizomato	Terrestrial		y dormant Seasonall	unisexual	absent	reduced	inaperturate
<i>Hestia</i>	Terrestrial	us Rhizomato	Terrestrial	Tropics	Evergreen	unisexual	absent	reduced	inaperturate
<i>Homalomena</i>	Terrestrial	us	Terrestrial	Tropics	Evergreen	unisexual	absent	reduced	inaperturate
<i>Hottarum</i>	Terrestrial	Caulscent	Terrestrial	Tropics	Evergreen Seasonall	unisexual	absent	reduced	inaperturate
<i>Incarum</i>	Terrestrial Rooted_aqu	Tuberous	Terrestrial	Tropics	y dormant	unisexual	absent	reduced	inaperturate
<i>Jasarum</i>	Rooted_aquatic	Caulscent Rhizomato	Aquatic	Tropics	Evergreen	unisexual	absent	reduced	inaperturate
<i>Lagenandra</i>	Rooted_aquatic	us	Aquatic	Tropics	Evergreen Seasonall	unisexual	absent	reduced	inaperturate
<i>Lazarum</i>	Terrestrial	Tuberous	Terrestrial	Tropics	y dormant	unisexual	absent	reduced	inaperturate
<i>Leucocasia</i>	Terrestrial	Tuberous Rhizomato	Terrestrial	Tropics	Evergreen	unisexual	absent	reduced	inaperturate
<i>Lorenzia</i>	Terrestrial	us	Terrestrial	Tropics	Evergreen Seasonall	unisexual	absent	reduced	inaperturate
<i>Mangonia</i>	Terrestrial	Tuberous	Terrestrial	Tropics	y dormant	unisexual	absent	reduced	inaperturate
<i>Montrichardia</i>	Rooted_aquatic								
<i>Nepthhytis_Africa</i>	Terrestrial	Caulscent Rhizomato	Aquatic	Tropics	Evergreen Seasonall	unisexual	absent	reduced	inaperturate
<i>Nepthhytis_Asia</i>	Terrestrial	us	Terrestrial	Tropics	y dormant Seasonall	unisexual	absent	reduced	inaperturate
<i>Ooia</i>	Terrestrial	Tuberous Rhizomato	Terrestrial	Tropics	y dormant	unisexual	absent	reduced	inaperturate
<i>Pelandra</i>	Terrestrial Rooted_aqu	us Rhizomato	Terrestrial Aquatic	Tropics Temperate	Evergreen Seasonall	unisexual	absent	reduced	inaperturate

	atic	us	Zones	y dormant				
<i>Philodendron</i>	Epiphytic	Climbing herbs	Terrestrial	Tropics	Evergreen	unisexual	absent	reduced
<i>Philotion</i>	Terrestrial	Rhizomato us	Terrestrial	Tropics	Evergreen	unisexual	absent	reduced
<i>Phymatarum</i>	Terrestrial	Caulscent Rhizomato	Terrestrial	Tropics	Evergreen	unisexual	absent	reduced
<i>Pichinia</i>	Terrestrial	us	Terrestrial	Tropics Temperate Zones	Evergreen Seasonall	unisexual	absent	reduced
<i>Pinellia</i>	Terrestrial	Tuberous	Terrestrial	Tropics	Evergreen Seasonall	unisexual	absent	reduced
<i>Piptospatha</i>	Terrestrial	Caulscent	Terrestrial	Tropics	Evergreen	unisexual	absent	reduced
<i>Pistia</i>	Free- floating	Free- floating	Aquatic	Tropics	Evergreen Seasonall	unisexual	absent	reduced
<i>Protarum</i>	Terrestrial	Tuberous	Terrestrial	Tropics	Seasonall y dormant	unisexual	absent	reduced
<i>Pseudohydros me</i>	Terrestrial	Tuberous	Terrestrial	Tropics	Seasonall y dormant	unisexual	absent	reduced
<i>Remusatia</i>	Epiphytic	Tuberous	Terrestrial	Tropics	Seasonall y dormant	unisexual	absent	reduced
<i>Sauomatum</i>	Terrestrial	Tuberous	Terrestrial	Tropics	Seasonall y dormant	unisexual	absent	reduced
<i>Scaphispatha</i>	Terrestrial	Tuberous	Terrestrial	Tropics	Seasonall y dormant	unisexual	absent	reduced
<i>Schismaoglot tis</i>	Terrestrial	Rhizomato us	Terrestrial	Tropics	Evergreen	unisexual	absent	reduced
<i>Schottariella</i>	Terrestrial	Rhizomato us	Terrestrial	Tropics	Evergreen	unisexual	absent	reduced
<i>Schottarum</i>	Terrestrial	Rhizomato us	Terrestrial	Tropics	Evergreen Seasonall	unisexual	absent	reduced
<i>Spathanthemum</i>	Terrestrial	Tuberous	Terrestrial	Tropics	Seasonall y dormant	unisexual	absent	reduced
<i>Spathicarpa</i>	Terrestrial	Tuberous	Terrestrial	Tropics	Seasonall y dormant	unisexual	absent	reduced
<i>Sieudhera</i>	Terrestrial	Caulscent	Terrestrial	Tropics	Evergreen	unisexual	absent	reduced
<i>Synandropad ix</i>	Terrestrial	Tuberous	Terrestrial	Tropics	Seasonall y dormant	unisexual	absent	reduced

	Climbing									
<i>Syngonium</i>	Epiphytic	herbs	Terrestrial	Tropics	Evergreen Seasonal	unisexual	absent	reduced	inaperturate	
<i>Taccarum</i>	Terrestrial	Tuberous	Terrestrial	Tropics	Evergreen Seasonal	unisexual	absent	reduced	inaperturate	
<i>Therophorum</i>	Terrestrial	Tuberous	Terrestrial	Tropics	Evergreen Seasonal	unisexual	absent	reduced	inaperturate	
<i>Typhonium</i>	Terrestrial	Tuberous	Terrestrial	Tropics	Evergreen Seasonal	unisexual	absent	reduced	inaperturate	
<i>Typhonodorum</i>	Terrestrial	Tuberous	Terrestrial	Tropics	Evergreen Seasonal	unisexual	absent	reduced	inaperturate	
<i>Ulearum</i>	Terrestrial	Rhizomatus	Aquatic	Tropics	Evergreen	unisexual	absent	reduced	inaperturate	
<i>Xanthosoma</i>	Terrestrial	Tuberous	Terrestrial	Tropics	Evergreen Seasonal	unisexual	absent	reduced	inaperturate	
<i>Zantedeschia</i>	Terrestrial	Tuberous	Terrestrial	Tropics	Evergreen Seasonal	unisexual	absent	reduced	inaperturate	
<i>Zomicarpa</i>	Terrestrial	Rhizomatus	Terrestrial	Tropics	Evergreen Seasonal	unisexual	absent	reduced	inaperturate	
<i>Zomicarpella</i>	Terrestrial	Rhizomatus	Terrestrial	Tropics	Evergreen Seasonal	unisexual	absent	reduced	inaperturate	
<i>Buomus</i>	Terrestrial	Rhizomatus	Aquatic	Tropics	Evergreen	bisexual	present	present	monosulcate	
<i>Amphibolis</i>	Terrestrial	Rhizomatus	Aquatic	Tropics	Evergreen	unisexual	absent	reduced	inaperturate	
<i>Cymodocea</i>	Terrestrial	Rhizomatus	Aquatic	Tropics	Evergreen	unisexual	absent	reduced	inaperturate	
<i>Halodule</i>	Terrestrial	Rhizomatus	Aquatic	Tropics	Evergreen	unisexual	absent	reduced	inaperturate	
<i>Ruppia</i>	Terrestrial	Rhizomatus	Aquatic	Tropics	Evergreen	bisexual	absent	present	triaperturate	
<i>Syringodium</i>	Terrestrial	?	Aquatic	Tropics	Evergreen	unisexual	absent	NA	NA	
<i>Thalassodendron</i>	Terrestrial	Rhizomatus	Aquatic	Tropics	Evergreen	unisexual	absent	reduced	inaperturate	
<i>Gymnostachys</i>	Terrestrial	Rhizomatus	Terrestrial	Tropics	Evergreen	bisexual	present	present	monosulcate	

<i>Apalanthe</i>	Rooted_aqu atic	Caulscent	Aquatic	Tropics	NA	bisexual	present	present	inaperturate
<i>Blyxa</i>	Rooted_aqu atic	Caulscent	Aquatic	Tropics Tropics- Temperate	NA	unisexual	present	present	inaperturate
<i>Egeria</i>	Rooted_aqu atic	Caulscent	Aquatic	Temperate Zones	Evergreen	unisexual	present	present	inaperturate
<i>Elodea</i>	Rooted_aqu atic	Caulscent	Aquatic	Tropics- Temperate Zones	Evergreen	unisexual	present	present	inaperturate
<i>Enhalus</i>	Rooted_aqu atic	Rhizomato us	Aquatic	Temperate Zones	Evergreen	unisexual	present	reduced	inaperturate
<i>Halophila</i>	Rooted_aqu atic	Caulscent	Aquatic	Tropics Tropics- Temperate Zones	NA	unisexual	present	reduced	inaperturate
<i>Hydrilla</i>	Rooted_aqu atic	Caulscent	Aquatic	Tropics- Temperate Zones	Evergreen	unisexual	present	reduced	inaperturate
<i>Hydrocharis</i>	Rooted_aqu atic	Caulscent	Aquatic	Temperate Zones	Evergreen	unisexual	present	present	inaperturate
<i>Lagarosiphon</i>	Rooted_aqu atic	Caulscent	Aquatic	Tropics Tropics- Temperate Zones	Evergreen	unisexual	present	present	inaperturate
<i>Limnobium</i>	Rooted_aqu atic	Caulscent	Aquatic	Temperate Zones	Evergreen	unisexual	present	present	inaperturate
<i>Najas</i>	Rooted_aqu atic	Caulscent	Aquatic	Tropics- Temperate Zones	Seasonall y dormant	unisexual	absent	reduced	inaperturate
<i>Nechamandra</i>	Rooted_aqu atic	Caulscent	Aquatic	Tropics	Seasonall y dormant	unisexual	present	present	inaperturate
<i>Ottelia</i>	Rooted_aqu atic	?	Aquatic	Tropics Temperate Zones	NA	unisexual	present	present	inaperturate
<i>Stratiotes</i>	Rooted_aqu atic	Caulscent	Aquatic	Temperate Zones	Evergreen	unisexual	present	present	inaperturate
<i>Thalassia</i>	Rooted_aqu atic	Rhizomato us	Aquatic	Tropics	Evergreen	unisexual	present	reduced	inaperturate
<i>Vallisneria</i>	Rooted_aqu atic	Caulscent	Aquatic	Tropics	NA	unisexual	present	reduced	inaperturate

	atic								
	Rooted_aquatic	?	Aquatic	NA Temperate Zones	NA	bisexual	present	NA	inaperturate
<i>Cynogeron</i>	Rooted_aquatic								
<i>Tetroncium</i>	atic	Caulscent	Aquatic	Tropics- Temperate Zones	Evergreen	bisexual	present	NA	inaperturate
<i>Triglochin</i>	Rooted_aquatic	Caulscent	Aquatic		Evergreen Seasonall	bisexual	present	present	inaperturate
<i>Anaphyllopsis</i>	Terrestrial	Rhizomatus	Terrestrial	Tropics	y dormant	bisexual	present	present	monosulcate
<i>Anaphyllum</i>	Terrestrial	Rhizomatus	Terrestrial	Tropics	Evergreen	bisexual	present	present	monosulcate
<i>Cyrtosperma</i>	Rooted_aquatic	Rhizomatus	Aquatic	Tropics	Evergreen Seasonall	bisexual	present	present	monosulcate
<i>Dracontioides</i>	Rooted_aquatic	Rhizomatus	Aquatic	Tropics	y dormant Seasonall	bisexual	present	present	monosulcate
<i>Dracontium</i>	Terrestrial	Tuberous	Terrestrial	Tropics	y dormant	bisexual	present	present	monosulcate
<i>Lasia</i>	Rooted_aquatic	Rhizomatus	Aquatic	Tropics	Evergreen	bisexual	present	present	monosulcate
<i>Lasimorpha</i>	Terrestrial	Rhizomatus	Terrestrial	Tropics	Evergreen	bisexual	present	present	monosulcate
<i>Podolasia</i>	Rooted_aquatic	Rhizomatus	Aquatic	Tropics	Evergreen Seasonall	bisexual	present	present	monosulcate
<i>Pycnospatha</i>	Terrestrial	Tuberous	Terrestrial	Tropics	y dormant	bisexual	present	present	monosulcate
<i>Urospatha</i>	Rooted_aquatic	Rhizomatus	Aquatic	Tropics	Evergreen	bisexual	present	present	monosulcate
<i>Landolphia</i>	Free-floating	Free-floating	Aquatic	Tropics- Temperate Zones	Evergreen	bisexual	absent	present	ulcerate
<i>Lemna</i>	Free-floating	Free-floating	Aquatic	Tropics- Temperate Zones	Evergreen	bisexual	absent	present	ulcerate
<i>Spirodela</i>	Free-floating	Free-floating	Aquatic	Tropics- Temperate Zones	Evergreen	bisexual	absent	present	ulcerate

<i>Wolffia</i>	Free-floating	Free-floating	Aquatic	Tropics-Temperate Zones	Evergreen	bisexual	absent	present	ulcerate
<i>Wolffella</i>	Free-floating	Free-floating	Aquatic	Tropics-Temperate Zones	Evergreen	bisexual	absent	present	ulcerate
<i>Mauandia</i>	Rooted_aquatic	Rhizomatous	Aquatic	Temperate Zones	NA	bisexual	present	present	inaperturate
<i>Alloschemone</i>	Epiphytic	Climbing herbs	Terrestrial	Tropics	Evergreen	bisexual	absent	present	
<i>Amydrium</i>	Epiphytic	Climbing herbs	Terrestrial	Tropics	Evergreen	bisexual	absent	present	zonate
<i>Anadendrum</i>	Epiphytic	Climbing herbs	Terrestrial	Tropics	Evergreen	bisexual	present	present	inaperturate
<i>Epipremnum</i>	Epiphytic	Climbing herbs	Terrestrial	Tropics	Evergreen	bisexual	absent	present	zonate
<i>Heropsis</i>	Epiphytic	Climbing herbs	Terrestrial	Tropics	Evergreen	bisexual	absent	present	zonate
<i>Holochlamys</i>	Terrestrial	Caulscent Climbing	Terrestrial	Tropics	Evergreen	bisexual	present	present	polyplicate
<i>Monstera</i>	Epiphytic	herbs	Terrestrial	Tropics	Evergreen	bisexual	absent	present	zonate
<i>Rhaphidophora</i>	Epiphytic	Climbing herbs	Terrestrial	Tropics	Evergreen	bisexual	absent	present	zonate
<i>Rhodospatha</i>	Epiphytic	Climbing herbs	Terrestrial	Tropics	Evergreen	bisexual	absent	present	zonate
<i>Scindapsus</i>	Epiphytic	herbs	Terrestrial	Tropics	Evergreen	bisexual	absent	present	zonate
<i>Spathiphyllum</i>	Terrestrial	Caulscent Climbing	Terrestrial	Tropics	Evergreen	bisexual	present	present	polyplicate
<i>Stenospermation</i>	Epiphytic	herbs	Terrestrial	Tropics	Evergreen	bisexual	absent	present	zonate
<i>Lysichiton</i>	Rooted_aquatic	Rhizomatous	Aquatic	Temperate Zones	Seasonally dormant	bisexual	present	present	monosulcate
<i>Orontium</i>	Rooted_aquatic	Rhizomatous	Aquatic	Temperate Zones	Seasonally dormant	bisexual	present	present	monosulcate
<i>Symplocarpus</i>	Rooted_aquatic	Rhizomatous	Aquatic	Temperate	Seasonally dormant	bisexual	present	present	monosulcate

	atic	us	Zones	y dormant				
	Rooted_aquatic	Rhizomatous	Temperate Zones	Evergreen	bisexual	absent	reduced	inaperturate
<i>Posidonia</i>	Rooted_aquatic	?	Temperate Zones	NA	bisexual	present	present	inaperturate
<i>Groenlandia</i>	Rooted_aquatic	?	NA	NA	unisexual	present	present	inaperturate
<i>Lepilaena</i>	Rooted_aquatic	?	NA	NA	unisexual	present	present	inaperturate
<i>Potamogeton</i>	Rooted_aquatic	?	Tropics-Temperate Zones	Evergreen	bisexual	present	present	inaperturate
<i>Stuckenia</i>	Rooted_aquatic	?	NA	NA	bisexual	present	present	inaperturate
<i>Zannichellia</i>	Rooted_aquatic	?	Tropics-Temperate Zones	NA	unisexual	absent	present	inaperturate
<i>Anthurium</i>	Epiphytic	Climbing herbs	Tropics	Evergreen	bisexual	present	present	forate
<i>Pedicularum</i>	Epiphytic	Climbing herbs	Tropics	Evergreen	bisexual	present	present	monosulcate
<i>Potheidium</i>	Epiphytic	Climbing herbs	Tropics	Evergreen	bisexual	present	present	monosulcate
<i>Pothos</i>	Epiphytic	Climbing herbs	Tropics	Evergreen	bisexual	present	present	monosulcate
<i>Scheuchzeria</i>	Rooted_aquatic	Rhizomatous	Tropics Temperate Zones	NA	bisexual	present	present	inaperturate
<i>Isidrogavia</i>	Rooted_aquatic	Caullescent	NA	NA	bisexual	present	present	monosulcate
<i>Pilea</i>	Terrestrial	Caullescent	NA	NA	bisexual	present	present	diaperturate
<i>Tofeldia</i>	Terrestrial	Caullescent	NA	NA	bisexual	present	present	diaperturate
<i>Triantha</i>	Terrestrial	Caullescent	NA	NA	bisexual	present	present	diaperturate
<i>Gonatopus</i>	Terrestrial	Tuberous Rhizomatous	Tropics	Seasonally dormant	unisexual	present	present	zone
<i>Strylochaeton</i>	Terrestrial	Rhizomatous	Tropics	Seasonally dormant	unisexual	present	present	inaperturate
<i>Zamioculcas</i>	Terrestrial	Rhizomatous	Tropics	Seasonally dormant	unisexual	present	present	zone

<i>Phyllospadix</i>	Rooted_aquatic	Rhizomatous	Aquatic	Temperate Zones-Tropics-Temperate Zones	Evergreen	unisexual	absent	reduced	inaperturate
<i>Zostera</i>	Rooted_aquatic	Rhizomatous	Aquatic		Evergreen	unisexual	absent	reduced	inaperturate

Chapter 2

Systematics, biogeography and morphological character evolution of the hemi-epiphytic subfamily Monsteroideae (Araceae)

Alejandro Zuluaga¹, Ken Cameron¹

¹ Department of Botany, University of Wisconsin-Madison

ABSTRACT

The subfamily Monsteroideae (Araceae) is the third richest clade in the family with about 500 species. It comprises mostly hemi-epiphytic or epiphytic species restricted to the tropics, with three continental tropical disjunctions. Using a dataset representing all twelve genera in Monsteroideae (ca. 100 taxa), and five plastid and two nuclear markers, we studied the systematics and historical biogeography. We found high support for the monophyly of the three major clades (Spathiphyllae, *Heteropsis* clade and *Rhaphidophora* clade), and for seven of the genera within Monsteroideae. However, we found low rates of DNA variation, and the lack of molecular markers suitable for species level phylogenies in the group. Seed shape, size, number of seeds, and presence of endosperm showed utility in the classification of genera in Monsteroideae. We estimated ancestral ranges using a dispersal-extinction-cladogenesis model as implemented in the R package BioGeoBEARS and found evidence for a Laurasian origin of the clade. One tropical disjunction (*Monstera* sister to *Amydrium* + *Epipremnum*) was found to be the product of a previous Boreotropical distribution. Two other disjunctions are more recent and probably due to long distance dispersal. *Spathiphyllum* (with *Holochlamys* nested within) represents dispersal from South America to the islands in the Pacific of South East Asia; and *Rhaphidophora* represents dispersal from Asia to Tropical Africa.

Keywords

Araceae, BioGeoBEARS, biogeography, character evolution, tropical disjunction, Monsteroideae

INTRODUCTION

The family Araceae belongs to the early diverging monocot order Alismatales and comprises ca. 122 genera and ca. 3500 species —although ca. 4000 are estimated by Boyce & Croat (2014); they are distributed worldwide (Mayo *et al.* 1997; Cusimano *et al.*, 2011a). Approximately 90% of the genera and 95% of the species of Araceae are found in the tropics, with approximately half of the species believed to be exclusively Neotropical (Croat, 1979; Nauheimer *et al.*, 2012). Araceae is considered one of the dominant tropical families of angiosperms, and one of the most important components of humid forests, where aroids are characterized for both high species diversity and local endemism (Croat 1983, 1988, 1992). Despite extensive work by many fine taxonomists (see Croat, 1998 for a detailed taxonomic history of the family), systematics of tropical Araceae remains a field of active research with about 280 new species described in the last four years alone. In addition, several molecular phylogenetic studies in Araceae have been published recently. Although these have focused mostly on the family level, species level studies of Old World and temperate taxa have been conducted as well (e.g., Grob *et al.*, 2004; Renner *et al.*, 2004; Mansion *et al.*, 2008; Wong *et al.*, 2010). In spite of the fact that tropical America holds ca. 2000 species of aroids, only six species-level phylogenetic studies focused on Neotropical taxa have been published to date (*Anthurium* [Carlsen *et al.*, 2013]; *Philodendron* [De Oliveira *et al.*, 2014; Gauthier *et al.*, 2008], Spathicarpeae [Gonçalves *et al.*, 2007], and Caladieae [Loh *et al.*, 2000]).

The most recent classification of Araceae divides the family into eight clear subfamilies, although some issues of delimitation remain, especially in the largest

subfamily Aroideae (Cabrera et al., 2008; Cusimano et al., 2011a, Chartier et al., 2014; Henriquez et al., 2014). The subfamilies Monsteroideae and Pothoideae form a clade that is sister to the rest of ‘true Araceae’ (Cabrera et al., 2008; Cusimano et al., 2011a; Henriquez et al., 2014). The Monsteroideae is one of the most taxonomically challenging clades within Araceae (Tam et al., 2004; Zuluaga et al., 2015 **See Appendix 1**), and is the third richest subfamily with some 400 described species (and many still undescribed). As part of one of the earliest diverging lineages in the family, Monsteroideae may help to provide a clearer picture of the evolution within the ecologically important family Araceae. It comprises mostly hemiepiphytic and epiphytic climbers (Croat, 1988; Madison, 1977; Croat & Boyce, 2014) restricted to tropical America and Asia, except for one or two species that are native to tropical Africa (this intercontinental disjunction is interesting in its own right). The subfamily has a long taxonomic history, with *Monstera* being one of the first genera described in Araceae by Adanson in 1763. Schott (1860) first grouped together various genera from Monsteroideae based on floral morphology. Engler (1876) and Engler & Krause (1908) carried out two early taxonomic revisions of the group in which they circumscribed Monsteroideae to have two tribes and 12 genera based on leaf and inflorescence morpho-anatomical characters. Since then, many studies using different approaches and datasets have attempted to propose a robust classification of Araceae, including a classification for Monsteroideae (e.g., Bunting, 1962; Madison, 1977; Grayum, 1990; Bogner & Nicholson, 1991; Hay & Mabberley, 1991; French et al., 1995; Mayo et al., 1997; Carvell, 1989; Keating, 2003, 2004a, 2004b). However, a review of the historical systematic treatments of Monsteroideae has revealed the lack of consistent morphological characters used to delimit the genera and species (Tam et al.

2004). For instance, different authors have recognized 2-12 genera using the same set of characters (e.g., Bunting, 1962; Madison, 1977; Carvell, 1989; Grayum, 1990).

Vegetative characters, in particular, (e.g., leaf shape) have been shown to be polymorphic within species, and convergent among genera (Madison, 1977). For this reason, currently, non-vegetative characters such as placentation, seed shape and size, septation of the ovary, and geography are regularly in use to define the genera (Bunting, 1962; Madison, 1977; Madison & Tiffany, 1976; Mayo et al., 1997). The current circumscription of Monsteroideae was not established until French et al. (1995) and Mayo et al. (1997) initiated phylogenetic work at the family level using AFLPs and morphology respectively. They found evidence to include *Heteropsis*, *Anadendrum*, and the tribe Spathiphyllae (*Spathiphyllum* + *Holochlamys*) within Monsteroideae. Tam et al. (2004) proposed the first molecular phylogenetic hypothesis for Monsteroideae using the plastid *trnL-trnF* spacer, and found strong support for the subfamily, but not much support for clades within the subfamily. Family-level molecular phylogenetic studies by Cabrera et al. (2008) and Cusimano et al. (2011a), based also exclusively on plastid DNA (*trnL-trnF* spacer, *rbcL* and *matK*), found the same clades as Tam et al. (2004) with slightly better support, but the monophyly of the genera was not tested. As a result, considerable debate has persisted around the circumscription of several genera of Monsteroideae such as *Amydrium*, *Epipremnum*, *Monstera*, *Rhaphidophora* and *Scindapsus*.

The current classification of Monsteroideae by Cusimano et al. (2011a) recognizes 12 genera grouped into three clades: 1) a Tropical America clade formed by *Alloschemone*, *Heteropsis*, *Rhodspatha* and *Stenospermation*, sister to the rest of the

subfamily; 2) the tribe Spathiphyllae comprising *Holochlamys* and *Spathiphyllum*; and 3) a mostly Old world clade comprising *Amydrium*, *Anadendrum*, *Epipremnum*, *Monstera*, *Scindapsus*, and *Rhaphidophora*. The first and third of these clades have been informally named the ‘*Heteropsis*’ and ‘*Rhaphidophora*’ Clades, respectively, by the same authors, and we will follow these names for simplicity. More recent studies using nuclear, chloroplast, and even plastome datasets have instead found Spathiphyllae to be sister to the rest of the Monsteroideae (Chartier et al., 2014, Henriquez et al., 2014; Zuluaga et al., 2015), as opposed to the *Heteropsis* Clade being sister to the rest.

Monsteroideae exhibits three unusual intercontinental geographic disjunctions: 1) *Spathiphyllum*, found primarily in Tropical America, also contains three species in Southeast Asia; these are in a clade with *Holochlamys becarii* from New Guinea; 2) *Monstera*, an exclusively Tropical American genus, is nested in a clade of otherwise Old World genera; and 3) *Rhaphidophora*, mostly from Asia, has one (or two) species in Africa. This pattern of sister groups being disjunct between tropical regions of the New and Old Worlds is shared with several angiosperm groups (e.g., Malpigiaceae, Rubiaceae (Nie et al., 2013), Magnoliaceae (Azuma et al., 2001), Leguminosae and Lauraceae (Renner, 2004)). The pattern has been explained by several competing hypotheses: 1) vicariance involving the break up of the Gondwanan supercontinent (Raven and Axelrod, 1974; Conti et al., 2002); 2) existence of a Boreotropical flora with subsequent extinction/migration across land bridges during Eocene/Oligocene (Davis et al., 2002); and 3) long-distance dispersal (LDD) (Renner et al., 2001; Givnish et al., 2004; Renner, 2004; Clayton et al., 2009; Christenhusz & Chase, 2013). Disjunction

between the tropics of Africa and Asia include two additional hypothesis: 1) an overland dispersal from eastern or south west Asia to Africa during the Miocene (Kappelman et al., 2003); and 2) dispersal from India to Madagascar through the ‘Lemurian stepping stones’ across the western Indian Ocean, and then to Africa during the Eocene-Oligocene (Schatz, 1996). Nauhmeier et al. (2012) investigated the history of geographic distribution in Araceae and suggested that many of the actual tropical disjunctions are products of extinction in Eurasia and North America, and migration to the tropics during the Oligocene (Boreotropical hypothesis) or the Quaternary. A trans-Laurasian range expansion has been suggested for the current distribution of *Spathiphyllum*. Madison (1977) and van Steenis (1962) mentioned that the Boreotropical flora hypothesis might explain the current distribution of *Monstera* and *Epipremnum*. Until now, however, there has been no objective attempt to explain disjunctions in Araceae using dated phylogenetic reconstructions.

Unlike most angiosperm families, the fossil record of Araceae is extensive, and represents (along with some other members of Alismatales) the oldest fossil records in the monocots (Stockey et al., 1997, 2007; Friis et al., 2004, 2006, 2010a; Bogner et al., 2005, 2007; Wilde et al., 2005). However, it is often difficult to assign fossils to particular subfamilies or extant genera within Araceae (Friis et al., 2010a; 2010b). Fossils that with certainty can be attributed to Monsteroideae come from the Cretaceous in India (Bonde 2000), and from the Pleistocene in southwestern Europe (Madison & Tiffney, 1976).

Here we expand previously published taxon and gene sampling across the subfamily Monsteroideae in order to test alternative phylogenetic and historical biogeographic hypotheses. One objective is to assess the utility of different molecular markers in reconstructing phylogenetic hypothesis down to the species level in Araceae, and we compare our results to previous classification systems. We also estimate ancestral ranges, and investigate the origin of the three unusual tropical geographic disjunctions in Monsteroideae. Finally, we evaluate the evolution of several key morphological characters traditionally used in taxonomic revisions of the subfamily.

METHODS

Sampling. A total of 126 accessions from subfamily Monsteroideae and members of the other seven subfamilies were sampled for this study. We included all 12 genera of the subfamily Monsteroideae, and available sequences in GenBank for 14 species of genera in the subfamily Pothoideae (*Anthurium*, *Pothos*, *Pedicellarum* and *Pothoideum*), and ten species representing the other six aroid subfamilies as outgroups (**Table 1**). Monsteroideae sampling included: one accession of the genus *Alloschemone*, five of *Amydrium*, three of *Anadendrum*, seven of *Epipremnum*, the monotypic genus *Holochalmys*, 28 of *Monstera*, 24 of *Rhaphidophora*, eight of *Rhodospatha*, four of *Scindapsus*, nine of *Spathiphyllum* (both Old and New World spp.), and nine of *Stenospermation* (**Table 1**). DNA samples from Neotropical genera were obtained from fieldwork (especially in Colombia, Ecuador and Panama), herbarium specimens, and/or the Araceae Greenhouse at the Missouri Botanical Garden in St. Louis, Missouri. Most of the DNA samples for species from Tropical Asia (e.g., *Rhaphidophora*, *Epipremnum*)

were obtained from the Royal Botanical Gardens Kew DNA Bank, and from the Jardin Botanique de Lyon.

DNA extraction, amplification and sequencing. DNA was extracted from silica-dried plant material, herbarium specimens, and/or living collections using the DNeasy Plant Mini Kit (Qiagen, Valencia, CA, USA) according to the manufacturer's specifications. Some recalcitrant samples were extracted using a CTAB method adapted from Doyle and Doyle (1987), or using the Promega, Maxwell[®] 16 platform.

Three plastid markers (*matK*, *rbcL* and *trnL-trnF* spacer) used in previous aroid studies were amplified (Cabrera et al., 2008; Cusimano et al., 2011a). Primers for *rbcL* came from Kress & Erickson (2007), for *matK* from Cuénoud et al. (2002), and for the *trnL-trnF* spacer from Taberlet et al. (1991). Additionally, ten other plastid markers highlighted in Shaw et al. (2005) were screened in a subset of samples across Monsteroideae in order to determine their nucleotide variability. The plastid regions *psbB-psbH*, *petA-psbJ*, *psbE-petL*, *rps16-trnQ*, and *trnT-psbD* spacers, and *ndhA* gene were amplified and sequenced successfully, but showed little variation, and were not used further in this study. The plastid markers *trnC-petN* spacer and partial *ycfI* showed the highest variation, and were amplified for the rest of the samples. We sequenced about 1000 bp of the first exon of the nuclear Phytochrome C (*PHYC*) gene using primers from Chartier et al. (2014) and new primers designed based on our sequences. *PHYC* sequences for several of the species in the *Rhaphidophora* clade presented many ambiguous nucleotides in the same positions among all the samples, suggesting the

presence of multiple copies in this clade. We performed BLAST searches of the *PHYC* gene sequences in the transcriptomes of *Anthurium andreanum* and *Landoltia punctata*, and in the *Spirodela polyrhiza* genome available in GenBank, and found a single match to the *PHYC* gene. Although previous studies in Araceae do not report the detection of multiple copies for the *PHYC* gene (Chartier et al. 2014), and our limited BLAST searches support this fact, we excluded all *PHYC* sequences from samples of the *Rhaphidophora* clade in our analysis since allele homology could not be verified. Finally, we developed primers for a region of about 700 bp in the second intron of the nuclear gene Chalcone Synthase (*CHS*), based on sequences from Carlsen & Croat (2013) and other monocots available in GenBank. We performed the same blast searches as in *PHYC*, and did not find multiple matches for these *CHS* sequences. Specific primers developed for *ycfI*, *PHYC*, and *CHS* are shown in **Table 2**.

PCR amplification was performed following the protocols from Cabrera et al. (2008). Nuclear marker amplification was enhanced by using 1-3 ul of betaine (10%) in a 25-ul reaction, and/or by adding up to 2 ul of DNA. The PCR products were purified using ExoSap-It (Cleveland, OH). Sequencing reactions used the BigDye Terminator Cycle Sequencing Kit (Applied Biosystems, Foster City, CA, USA), and were cleaned using Agencourt CleanSeq (Beverly, MA) magnetic beads. Samples were sequenced at the University of Wisconsin–Madison Biotechnology Center.

Molecular analysis. All sequences were assembled and manually edited in Geneious v8.0 (Biomatters, <http://www.geneious.com/>). Every gene was aligned using a

combination of MAFFT v7 (Katoh et al., 2002; Katoh & Standley, 2013) and MUSCLE v3.8. (Edgar, 2004). An initial alignment was performed in MAFFT with two subsequent refinements using MUSCLE. All alignments were checked manually in Geneious v8.0 using protein translation for the coding regions. We used PartitionFinder (Lanfear et al., 2012) to select the best-fit partitioning schemes and models. For the initial data blocks in PartitionFinder we allocated nucleotides from the protein-coding genes (*rbcL*, *matK*, *PHYC*, and *CHS*) according to whether they belonged to the first, second or third codon position, and all the other markers as independent data blocks, including the partial *trnK* intron flanking the *matK* gene, and the partial *ycf1* gene. The *ycf1* sequences were not allocated into codon positions because of the difficulty in establishing the reading frame due to the presence of a 37bp inversion. We performed an initial PartitionFinder run allowing all models, and then a subsequent run allowing only models available in RAxML and MrBayes respectively. We used Bayesian information criteria (BIC) to choose the best-fit models. PartitionFinder suggested three partitions for the plastid dataset (**Table 3**): (1) the entire *matK* gene, the second codon position of *rbcL*, the *trnL-trnF* spacer and the partial *ycf1*; (2) the *trnK* intron, and the *trnc-petN* spacer; and (3) the first and third positions of *rbcL*. The nuclear PHYC data set had two partitions: (1) first and second codons positions; and (2) the third codon position. The nuclear CHS data set had three partitions, one for each codon position. Individual phylogenetic analyses for each marker were performed under Maximum Likelihood (ML) using RAxML v8.0 (Statamakis, 2006), and Bayesian Inference (BI) using MrBayes v3.2.3 (Ronquist et al., 2012) as implemented in the CIPRES portal (Miller et al., 2010). For RAxML, we used the best scheme for PartitionFinder, and GTRGAMMA model in all cases. We performed

100 independent tree searches and 1000 independent bootstraps analyses. For MrBayes we used the best scheme and models suggested by PartitionFinder. Bayesian analyses were performed using two independent MCMC runs with four chains and 20 million generations. A combined plastid analysis and also a “total evidence” analysis were performed in the same way. Additional unpartitioned analyses were performed for every individual and variously combined dataset.

Molecular dating. We employed the five-plastid markers dataset for Bayesian molecular dating using BEAST 1.8 (Drummond et al., 2012), and three calibrations points as used in Zuluaga et al. (**Chapter 1**) following recommendations of Sytsma et al. (2014). We used a set of fossils from the early Cretaceous (110 Ma) in Portugal described by Friis et al. (2010a) and Friis et al. (2010b) to constrain the crown node of a clade comprising Lemnoideae and the rest of Araceae (sensu Cusimano et al., 2011a). These fossils correspond to bisexual and unisexual inflorescences that can be assigned with certainty to the family, but their precise affinities within Araceae need more study. Secondly, we assigned *Petrocardium cerrejoense* from the Mid to Late Paleocene (56-61.6 Ma) in Colombia (Herrera et al., 2008) for the crown node of Pothoideae. Finally, for the stem node of the tribe Monstereae (subfamily Monsteroideae) we assigned *Rhodospathodendron tomlinsonii*, from the Late Cretaceous (66 Ma) from India (Bonde, 2000). We used lognormal priors with a mean=10 and SD=0.75 on the fossil calibrations. For the root of the tree we used a Uniform prior (110-135 Ma), based on previous dates calculated for the family Araceae (Nauheimer et al. 2012; Zuluaga et al. unpublished data)

Ancestral range estimation. We used the chronogram from our BEAST analysis as an input for the biogeography analysis. The chronogram was pruned to contain one sample per operational taxonomical unit (OTU). We chose the package BioGeoBEARS (Matkze, 2013) in R (R Development Core Team, 2008) for ancestral geographic area reconstruction because it allows us to compare several models and include parameters such as unequal rates of dispersal between areas and different dispersal rates through time. However, we focused on the comparison between the DEC model (Ree et al., 2005; Ree and Smith, 2008) and the DEC+j model, which incorporates founder-event speciation and allows for dispersal without range expansion (Matzke, 2012).

Initially we defined eight geographic areas modified from Kissling et al. (2012) to accommodate the distribution of the genera in Monsteroideae and outgroups. Nearctic (A), representing North America; Central America (B) including Southern Mexico, and the Caribbean islands; South America (C); Palearctic (D), including Europe and Temperate Asia; Africa (E); Southeast Asia (F); Australasia (G) including, Australia, New Zealand, New Guinea and the Pacific islands; and India (H). An additional set of analyses were performed merging Neotropical areas B and C. Areas occupied for each OTU were defined based on data provided by the Monocot CheckList (WCSP, 2014), herbarium specimens, and taxonomic monographs. *Rhodspatha* and *Stenospermation* were coded as present in Central and South America, since our sampling did not allow for finer separation. *Heteropsis* has only one species extending north into Central America, which is absent from our phylogenetic analysis, so this genus was coded as South

America only. Finally, *Monstera* species were coded according to their current distributions, either Central or South America or both.

We used a time-slice model, and different dispersal multipliers among areas, to try to represent connectivity among different areas at different periods of time. We performed a second analysis giving all areas equal dispersal probability through time to evaluate the effect of the time-slice model. The time slices used were 0-10, 10-50, 50-90 and 90-140 Ma. For the 0-10 Ma slice all main landmasses were in their current position. Palearctic and Nearctic were connected by the Beringian land bridge, and the connection between the Nearctic and Tropical America was completed. From 10 to 50 Ma main landmasses were close to their current position. India connected with Southeast Asia and the Palearctic. From 50-90 Ma (the Late Cretaceous to the Eocene and across the K-T boundary), India was moving towards Asia and the Palearctic, and Australasia separated from Antarctica. Finally, from 90-140 Ma (Early to Late Cretaceous), West Gondwana (South America + Africa) began to separate, as well as India and Madagascar, as both masses started to drift away from East Gondwana (including Australasia).

Since models used are nested, we compared them using a Likelihood ratio test (LRT) as implemented in BioGeoBEARS. To account for uncertainty in the phylogenetic reconstructions we used 100 randomly selected trees from the posterior distribution from the BEAST analysis after burn-in. We ran the DEC model (the preferred model according to LRT) on these 100 trees and counted the ancestral area estimation with highest frequency in nodes of particular interest. The nodes evaluated were the most common

recent ancestors (MRCA) of: *Rhaphidophora africana*, the genus *Monstera*, *Rhaphidophora* + *Anadendrum*, the *Rhaphidophora* clade, the *Heteropsis* clade, the Spathiphyllae, a clade comprising the so-called *Heteropsis* and *Rhaphidophora* clades, Monsteroideae, and a clade comprising Pothoideae plus Monsteroideae.

Character evolution. From the literature, herbarium specimens, spirit collections, and fieldwork observations we coded a set of characters traditionally used by aroid systematists in the delimitation of genera in Monsteroideae. The characters coded were: number of ovules, number of locules, presence of endosperm, seed length in lateral view, and seed width in lateral view (**Table 8**). The same species as the ones in our phylogenetic tree were used when possible, however the reduced number of collections and the lack of taxonomic clarity in genera like *Rhodospatha* and *Rhaphidophora* made it impossible to match our molecular and morphological sampling. In addition to discrete characters, images of seeds in lateral view for several species of each genus were obtained. These 2D-images were processed using the software Gimp v.2.8 to obtain outlines of the shape, and measurements of length and width in lateral view for each seed. Shapes were quantified using elliptic Fourier analysis in the R package Momocs v1.0 (Bonhomme et al., 2014), and principal component analysis (PCA) was performed on the harmonics produced by the outline analysis. Discrete characters were reconstructed using stochastic mapping (Bollback, 2006), and the rerooting method in the R package phytools (Revell, 2012). Because of the issues matching our phylogenetic tree with our seed data, continuous characters (seed size and PC components from pollen shape) were not evaluated in a phylogenetic framework.

RESULTS

Phylogenetic reconstructions. The concatenated alignment of the five plastid markers represents 6192 bp, and contains 38% gaps or missing data (**Table 4**). Variable sites vary between 8.3% (*rbcL*) and 51.7% (*matK*) for the ingroup. However, most of this variation is among genera, whereas variation within genera is low, ranging from only 1-10%. The partial *ycfI* has an indel of about 200 bp in the genus *Stenospermation*, and presents an inversion of 37 bp in several species across the subfamily; this region was excluded of the analyses. *trnL-trnF* spacer contains several indels, and a region of ca. 250 bp with high TA content and repeats was also excluded from the analyses. Finally, the *trnC-petN* spacer is considerably reduced in some species of the genus *Rhaphidophora* (only 300-340 bp) relative to the other taxa. CHS and PHYC alignments contain 692 bp and 1093 bp with 0.3 and 16.8 % missing data respectively (**Table 4**).

In general, Maximum Likelihood and Bayesian analyses of the individual plastid and nuclear datasets produced topologies largely lacking in resolution within the genera of Monsteroideae. ML bootstrap (BS) values > 75% and posterior probabilities from Bayesian inference (PP) > 0.95 are considered as “strong” support for the following results and discussion. The three major clades within Monsteroideae discussed earlier are recovered with strong support in all individual and combined analyses with plastid and nuclear markers, except with the nuclear marker CHS (**Table 5, Figures. 1-2**). There are no significant topology differences among ML partitioned and “unpartitioned” analyses, but support values varied, especially for the CHS marker (**Table 5, Fig 1-2**). ML and BI

topologies vary slightly for some species relationships in nodes with low support.

Concatenated analyses resemble the CHS gene tree for the relationships among genera.

Intergeneric relationships within Monsteroideae. The tribe Spathiphyllae is sister to the rest of Monsteroideae in all cases, but the support for this position is low (BS < 75; PP<0.95) for CHS analysis (**Figure 1-5**). Hypotheses for the generic relationships from different analyses are summarized in **Figure 5**. The plastid analyses recovered eight genera as monophyletic, seven of them with strong support (**Table 5, Figure 5**). However, only the genera *Anadendrum*, *Monstera*, *Rhodspatha*, *Scindapsus*, and *Stenospermation* are recovered as monophyletic consistently through all the analyses with high support (**Table 5, Figures 1-5**). The genera *Amydrium*, *Epipremnum*, *Heteropsis*, *Rhaphidophora*, *Spathiphyllum* and *Holochlamys*, appeared as not monophyletic in at least one analysis (**Table 5**). *Epipremnum* comprises at least two clades, the first one sister to the genus *Scindapsus*, and the second one to the genus *Amydrium*. *Rhaphidophora* is recovered as monophyletic in the combined plastid analyses (with no support), but not in the nuclear and total evidence analyses, where some species are sister to *Anadendrum*. Finally, the monotypic genus *Holochlamys* is nested within the genus *Spathiphyllum* with the plastid datasets. Nuclear dataset did not include enough samples to test *Spathiphyllum* vs. *Holochlamys* monophyly.

Molecular dating. Divergence time estimation analysis found an age of 68.83 Ma (95% confidence interval (CI) 64.20-76.73) for the crown age of Monsteroideae corresponding to the Late Cretaceous (**Figure 6**). The ages for the crown of the three

main Monsteroideae clades are considerably younger than the crown of the subfamily, all falling into the Oligocene, but with the 95% confidence intervals extending into the Miocene and Eocene. The crown age for Spathiphyllae is 26.97 Ma (95% CI 14.64-42.9), 38.44 Ma (95% CI 27.68-50.88) for the *Rhaphidophora* clade, and 38.56 Ma (95% CI 26.92-51.60) for the *Heteropsis* clade (**Table 6, Figure 6**).

Ancestral range estimation. The models with the addition of the time period slices conferred better likelihoods to the data (DEC+time periods LnL=-196.8 vs DEC LnL=-189, $p=7.6e-05$). Addition of the founder event (DEC + J) did not increase significantly the likelihood in any of the analyses ($p=0.55-0.66$), so we chose the simpler model to estimate the historical ranges within Monsteroideae. We will focus on the results of the seven-area analyses with time periods (DEC ln L=-190.81, DECj ln L=-190.91, $p=0.66$) for the results and discussion. The ancestor of Monsteroideae and the *Heteropsis* clade were estimated to range from Tropical America (BC) with higher probability ($P=0.64$ and $P=0.99$ respectively), and our analysis taking into account uncertainty recovered the range BC as the most probable ($P>0.5$) over the 100 trees (**Table 7, Figures 7 & 8**). The ancestral range of the *Rhaphidophora* clade was estimated to be in South East Asia (F) ($P=0.54$); however, this range was only recovered in 24 trees with $P>0.50$ ($P=0.31$, $SD=0.22$). Alternatively a range comprising areas BC (Tropical America) and F (South East Asia) has $P=0.09$ in the single chronogram, but its average probability over 100 trees is 0.23. The ancestral range of the Spathiphyllae was estimated to include Tropical America and Australasia (BC+G) ($P=0.77$) and the same range was recovered in 93 trees in the uncertainty analyses ($P=0.24$, $SD=0.19$).

Character evolution. ML and Bayesian estimations of the ancestral states for the discrete characters produced very similar results (**Figures 9 & 10**). The characters traditionally used for taxonomic purposes have evolved several times in Monsteroideae. Perianth was lost in *Rhaphidophora* + *Heteropsis* clades and regained in the genus *Anadendrum*. Endosperm is absent or very reduced in *Monstera*, *Heteropsis* and *Anadendrum*. Ancestral seed number is >10 with reductions to smaller numbers in several genera. PC1 of the seed shape analysis explain 69.2% of the variation, whereas PC2 and PC3 explained 13 % and 9.1 % respectively (**Figure 11**). There is not a clear separation among clades ($p=0.1503$) or genera ($p=0.78$) in Monsteroideae regarding seed shape, only the genus *Monstera* seems to have a distinctive in shape (**Figure 11**).

DISCUSSION

Monsteroideae classification and character evolution. Species level molecular studies of tropical aroids have been limited mainly because of the difficulty in obtaining samples, the lack of genomic tools, and the low variation of the molecular markers available. Taxonomic identification is another issue that remains problematic when working within the family Araceae. Our study explored the utility of several plastid and nuclear genes, and confirmed the lack of genetic informative regions for phylogenetic reconstruction at the species level in Tropical Araceae. The three main clades within Monsteroideae have been supported since the early molecular work (Tam et al., 2004; Cabrera et al., 2008). However, the position of Spathiphyllleae as sister to rest of

Monsteroideae is only recovered in the most recent studies with nuclear (Chartier et al., 2014), and plastome datasets (Henriquez et al., 2014). Our results strongly support the position of Spathiphyllae as sister to the rest of the Monsteroideae, in contrast to the topology used for the most recent aroid classification (Cusimano et al., 2011a); highlighting the importance of dense sampling for more accurate phylogenetic reconstructions when few molecular markers are available. The authors of that study proposed a set of synapomorphies to define the entire subfamily Monsteroideae that are, in fact, more appropriate for diagnosing the clade containing all Monsteroideae except Spathiphyllae (Zuluaga et al., 2015). These synapomorphies and the alternative characters present in Spathiphyllae are: 1) pollen zonate (i.e., ring-like or hamburger shape) (vs. pollen multiaperturate with exine surface striate/polyplicate in Spathiphyllae); 2) spathe soon deciduous with a distinct basal abscission (vs. spathe marcescent in Spathiphyllae); 3) large trichosclereids not in bundles (vs. smaller trichosclereids in bundles in Spathiphyllae). Additionally, another character that could be added is the hemiepiphytic or epiphytic habit (vs. terrestrial plants growing in shaded, wet places for Spathiphyllae). The only synapomorphy remaining for the entire Monsteroideae is the presence of trichosclereids, as proposed by Nicolson (1960). However, it is important to mention that detailed anatomical studies have found that trichosclereids are absent in the genera *Heteropsis* and *Anadendrum* (Carvell, 1989; Seubert, 1997; Keating, 2002, 2004a); they might be considered secondarily lost in these genera. Furthermore, trichosclerids are very occasionally present in at least four other species outside of Monsteroideae, for example in the genera *Pothos* (Nicolson, 1960), *Anthurium* (Keating, 2002), and *Montrichardia* (Solerder & Meyer, 1928). The last one is

a dubious report, and has never been confirmed with modern studies of *Montrichardia* tissues (Keating, 2002). Flower and seed characters investigated showed different levels of homoplasy. The clade Spathiphyllae is defined by having ovary trilocular. There seems to be convergence between *Heteropsis*, *Monstera* and *Anadendrum*, all having 1-2 large seeds, lacking endosperm; and between *Stenospermation* and *Rhaphidophora* with >10 small seeds with very similar shapes. However, a more dense morphological sampling is needed to test this hypothesis.

Spathiphyllae Clade. The support for the position of *Holochlamys becarri* within *Spathiphyllum* is low and therefore not conclusive, but the molecular analyses indicate that this species may in fact belong to the genus *Spathiphyllum*. *Holochlamys becarri* and the Asian species of *Spathiphyllum* form a monophyletic group (along with one accession of a South American species, *S. cannifolium*, that is nested in this clade as well). Species of this group are more closely related to the species of *Spathiphyllum* section *Massowia* than to the rest of *Spathiphyllum* (Zuluaga et al. 2015). *Holochlamys* and *Spathiphyllum* are very similar in habit and morphology (Bunting, 1960), stem vasculature (French & Tomlinson, 1981), palynological features (Grayum, 1992), and floral anatomy (Carvell, 1989). Moreover, *Holochlamys* shares with *Spathiphyllum* section *Massowia* the presence of fused perianth parts forming a cup around the pistil (Bunting, 1960; Carvell, 1989). Zuluaga et al. (2015) listed in detail the morpho-anatomical similarities between these two genera, but pointed out the need for wider sampling and well supported molecular data before drawing any conclusion about the taxonomic status of *Holochlamys*. At this point in time, all lines of evidence suggest that the two genera should be synonymized.

***Heteropsis* clade.** Despite the fact that genera in this group form a clade, the relationships among them remain unclear. Only *Heteropsis* and *Alloschemone*, the most morphologically unique genera in Monsteroideae regarding morphology, form a consistent clade through all the analyses. There are no apparent similarities between these two groups, besides their geographic distribution (restricted to the Amazon Basin), and so their shared ancestry was unexpected. *Alloschemone* is an enigmatic taxon; only two species with very few collections are known, and they grow far apart in the Amazon (Madison, 1976; Bogner et al., 2001). *Alloschemone* was previously considered to be part of the old world genus *Scindapsus* (Madison, 1976) based on the presence of unilocular ovaries. However, its combination of morphological and anatomical characters does not resemble that of any other Monsteroideae. There are some superficial similarities shared with the flower anatomy of *Rhodspatha*, *Stenospermation* and *Monstera* (Carvell, 1989; see Bogner et al., 2001 for a throughout revision of *Alloschemone* morpho-anatomical characters), but our molecular data do not support a close relationship to any of these other genera. Fruits of *Alloschemone* have never been seen, and their pollen grains have been described as problematic to interpret, with a very inconspicuous sulcus, whereas all other Monsteroideae have fully zonate aperture pollen. Most interestingly, Monsteroideae has a particular chromosome number evolution. The haploid number for the subfamily seems to be 15, with a genome duplication in the *Rhaphidophora* clade, $n=30$ (Cusimano et al., 2011b). Additionally, there are counts of $n=56$ in *Rhodspatha*, and $n=56$ and 84 in *Alloschemone*. Bogner et al. (2001) commented on the possibility of *Alloschemone* being a hexaploid. This polyploid origin, probably through hybridization,

could explain the unusual morphology, pollen anomalies, and the absence of fruit in *Alloschemone*, although better field collections and more molecular data are needed to support this hypothesis. Based on our results, future researchers are encouraged to more thoroughly compare *Alloschemone* with *Heteropsis*, and to consider the latter as putative parent for this unusual aroid genus.

***Rhaphidophora* clade.** This group represents the most taxonomically challenging within Monsteroideae (Boyce, 2001), and the status of several genera within it has been challenged as a result of our study. *Rhaphidophora* is monophyletic in the plastid dataset, but there is no support for this hypothesis. Additional data will be needed to confirm it. Only the genera *Monstera* and *Anadendrum* are clearly monophyletic, and morphological characters and geographical distribution support their status as well. *Monstera* presents with flowers with one locule, producing generally one seed, with a distinctive shape and lacking endosperm; whereas *Anadendrum*, present flowers with perianth and fruits with one seed, lacking endosperm. In contrast, the genus *Epipremnum* is polyphyletic, with some species related to *Scindapsus* and others to *Amydrium*. Seubert (1997) studied trichosclereids in the vegetative tissues of Monsteroideae and found that *Amydrium* species and *E. pinnatum* do not have trichosclereids in the aerial roots and lamina. Additionally, *Amydrium* species and *E. pinnatum* share adult pinnatifid leaves, although we concede that vegetative characters shown high plasticity in the *Rhaphidophora* clade (Bogner et al., 2001). Nevertheless, these features support the results of the gene tree topologies, and the taxonomy of *Epipremnum* may need to be revisited.

Molecular dating and ancestral range estimation. Our time divergence estimation agrees with that of Zuluaga et al. (unpublished data, **Chapter 1**), which included greater taxon sampling for Araceae, Alismatales, and more fossil calibrations. Because of the already mentioned disagreements in tree topology with Nauheimer et al. (2012), and the use of a different set of fossils, our date estimates differ significantly for the crown age of Monsteroideae and the major clades within the subfamily. Nauheimer et al. (2012) estimated an age of 56.3 Ma (vs. 68.8 in our analysis) for Monsteroideae; 34.16 Ma (vs 21.3) for Spathiphyllae; 21.6 Ma (vs. 38.5) for the *Heteropsis* clade; and 29.7 (vs. 38.4) for the *Rhaphidophora* clade. We considered our estimates reflect closer the current knowledge of fossil record of Araceae and in particular of Monsteroideae. Recently described and other undescribed fossils (Friis et al., 2004; Friis et al., 2010a; 2010b; Coiffard et al., 2013; Kvaček & Smith, 2015) show that the Araceae, and very clearly the subfamily Orontioideae, was widespread by the end of the Early Cretaceous (ca. 110 Ma). Most of the other subfamilies are already in the fossil record by the beginning of the Cenozoic (Friis et al., 2004), but it is probable they were already present by the late Cretaceous as indicated by previous studies and our molecular dating analysis. (Nauheimer et al., 2012; Zuluaga Unpublished data, **Chapter 1**). The affinity of some of the older fossils to Monsteroideae is inconclusive. *Mayoa portugallica* (110–120 Ma) resembles pollen from *Spathiphyllum* and *Holochlamys* (Friis et al., 2004; Friis et al., 2010), but these affinities have been challenged recently (Nauheimer et al., 2012a). Accurate fossil remains unquestionably assigned to Monsteroideae are from the Late Cretaceous (Bonde, 2000) in India, and from the Oligocene in Europe (Zetter et al., 2001; Madison & Tiffney, 1976). Other fossils from the Late Cretaceous in Europe resemble

Monsteroideae but they remain unpublished and were not used in our analyses (Friis et al. 2004).

Our biogeographic analyses estimated a Gondwanan ancestral range for the subfamily Monsteroideae, particularly in Tropical America. However, the fossil record and current distributions of extant taxa are difficult to explain completely with our biogeographic model. Additionally, the estimation for range the crown node of Pothoideae + Monsteroideae is ambiguous in our analyses (**Table 7**). The fossil record (albeit incomplete and poorly sampled from tropical areas of the world today) suggests that many members of Monsteroideae were part of the Boreotropical Flora, being widespread in the northern hemisphere from the late Cretaceous at least until the Oligocene (Madison & Tiffney, 1976), and then migrated to their current distributions in the tropics. This is not in conflict with a Gondwanan origin, but the inclusion of fossil distribution into the ancestral range estimation could change estimation drastically. Independently if the subfamily originated or arrived early to Tropical America during the late Cretaceous, we found evidence for independent dispersals events between tropical continental landmasses in the Monsteroideae. There are several cases of tropical disjunctions between sister groups in Araceae at the suprageneric level. For example, the Neotropical genera *Anthurium*, *Philodendron*, and *Jasurum* are sister to the Paleotropical Asian taxa *Pothos* (plus *Pothoidium* and *Pedicellarum*), *Homalomena*, and *Hapaline* respectively (Mayo et al., 1997; Grayum, 1992; Nauheimer et al., 2012). Many of these disjunctions have been explained by an early widespread distribution of Araceae in the Northern Hemisphere (Boreotropical Flora hypothesis) with migration southward to the

tropics of Central and South America, Asia, and Africa, and subsequent extinction in North America and Europe (Nauheimer et al., 2012). In our case only the disjunction between *Monstera* and the rest of the *Rhaphidophora* clade can be explained by this hypothesis. Madison (1977) and van Steenis (1962) already proposed this based solely on fossils and morphology affinities. The sister taxa of *Monstera* (*Amydrium* +some *Epipremnum*) are represented by several Oligocene and Pleistocene fossils from Europe (Madison & Tiffney, 1976). Additionally, our ancestral range estimation using Central and South America as independent areas showed that *Monstera* most probably migrated from North and Central to South America in the last 25 Ma (**Figure 6**).

In contrast, at the infrageneric level, tropical disjunctions are less common in Araceae. Species in the genera *Homalomena*, *Schismatoglottis* and *Spathiphyllum* were believed to be present in both the Neotropics and Asian Paleotropics (van Steenis, 1962; Grayum, 1990; Mayo et al., 1997). However, recent studies (Barabé et al., 2002, Gauthier et al., 2008, Wong et al., 2013; Zuluaga et al., 2015) have challenged the monophyly of both *Homalomena* and *Schismatoglottis* indicating that they do not contain disjunct species at all. This leaves *Spathiphyllum* as the only genus of Araceae exhibiting a New and Old World tropical disjunction. A second monsteroidean genus, *Rhaphidophora*, exhibits a less dramatic tropical disjunction between Asia and Africa. In any event, our ancestral range estimation and molecular dating suggests that both of these generic tropical disjunctions (*Spathiphyllum* and *Rhaphidophora*) correspond to more recent events. The divergence time estimates for the crown of Spathiphyllae (29.9 Ma) and current species distribution suggest long distance dispersal (LDD) between Tropical

America and Asia probably through the Pacific. Transoceanic dispersal in Tropical Araceae is uncommon (Nauheimer et al. 2012); however, one species of *Spathiphyllum*, (*S. leavae* -- which is morphologically very similar to *S. cannifolium* and the other Asian species) is present on Cocos Island, 342 mi from the Pacific shore of Costa Rica. Moreover, all the Asian species in the Spathiphyllaeae are distributed on islands in Australasia with only one species, *S. commutatum*, actually reaching continental Asia. Alternatively, if fossils like *Mayoa* (100-110 Ma) were confirmed to belong to Spathiphyllaeae, it will place this clade as distributed in the northern hemisphere, and a completely different hypothesis for the origin and dispersal of Monsteroideae (and even Araceae) may be needed.

Divergence times for *Rhaphidophora africana* (14 Ma) also suggest LDD from South East Asia to Africa. LDD between Tropical Asia and Africa is more common, and has been described to the genus level in several angiosperm groups (e.g. Nie et al., 2012; Nie et al. 2013), and so is less remarkable than the case of *Spathiphyllum*. However, dispersal from Asia to Africa seems to be less common than the opposite direction (Nie et. al, 2012). Since sampling among the Asian species of the *Rhaphidophora* clade is small and resolution within genera is poor and weakly supported in our tree, we cannot speculate further in terms of the patterns of historical biogeography within this diverse clade in Asia. We described a complex pattern of geographical distribution and morphological evolution in Monsteroideae. Future research based on stronger phylogenetic reconstructions and complete morphological datasets is needed to explore

the details of speciation and migration within and among areas of Asia, and movements between islands and continental landmasses.

REFERENCES

- Adanson, Michel (1763). Familles des plantes. Paris: Vincent. pp. I: 1–199; II: 1–640.
- Azuma, H., García-Franco, J.G., Rico-Gray, V., Thien, L.B., 2001. Molecular phylogeny of the Magnoliaceae: the biogeography of tropical and temperate disjunctions. *Am. J. Bot.* 88, 2275–2285.
- Barabé, D., Bruneau, A., Forest, F., Lacroix, C., 2002. The correlation between development of atypical bisexual flowers and phylogeny in the Aroideae (Araceae). *Plant Syst. Evol.* 232, 1–19.
- Bogner, J., Nicolson, D.H., 1991. A revised classification of Araceae with dichotomous keys. *Willdenowia* 21, 35–50.
- Bogner, J., Boyce, P.C., Sakuragui, C.M., 2001. A revision of *Alloschemone* Schott (Araceae: *Monstereae*). *Aroideana* 24, 80–93.
- Bogner, J., Hoffman, G.L., Aulenback, K.R., 2005. A fossilized aroid infructescence, *Albertarum pueri* gen. nov. et sp. nov., of Late Cretaceous (Late Campanian) age from the Horseshoe Canyon Formation of southern Alberta, Canada. *Can. J. Bot.* 83, 591–598.
- Bogner, J., Johnson, K.R., Kvaček, Z., Upchurch, G.R., 2007. New fossil leaves of Araceae from the Late Cretaceous and Paleogene of western North America. *Zitteliana* 47, 133–147.
- Bollback, J. P. (2006) Stochastic character mapping of discrete traits on phylogenies. *BMC Bioinformatics*, 7, 88.
- Bonde, S.D., 2000. *Rhodospathodendron tomlinsonii* gen. et sp. nov., an araceous viny axis from the Nawargaon intertrappean beds of India. *Palaeobotanist* 49, 85–92.

- Bonhomme, V., Picq S., Gaucherel C., Claude J., 2014. Momocs: outline Analysis using R. J Stat. Software 56:1–24.
- Boyce, P.C., 2001. The genus *Rhaphidophora* Hassk.(Araceae-Monsteroideae-Monstereae) in Borneo. Gard. Bull. Singapore 53, 19–74.
- Boyce, P.C., Croat, T., 2014. The Überlist of Araceae: totals for published and estimated number of species in aroid genera. Available via <http://www.aroid.org/genera/140601uberlist.pdf>. Accessed 5 May 2015
- Bunting, G.S., 1960. The genus *Schismatoglottis* (section *Philonotion*) in America. Ann. Missouri Bot. Gard. 47, 69–71.
- Bunting, G.S., 1962. Generic delimitation in the Araceae, subfamily Monsteroideae. Bailey 10, 21–31.
- Cabrera, L.I., Salazar, G.A., Chase, M.W., Mayo, S.J., Bogner, J., Dávila, P., 2008. Phylogenetic relationships of aroids and duckweeds (Araceae) inferred from coding and noncoding plastid DNA. Am. J. Bot. 95, 1153–1165.
- Carlsen & Croat 2013, Carlsen, M.M., Croat, T.B., 2013. A molecular phylogeny of the species-rich Neotropical genus *Anthurium* (Araceae) based on combined chloroplast and nuclear DNA. Syst. Bot. 38, 576–588. doi: <http://dx.doi.org/10.1600/036364413X670287>
- Carvell, W.N., 1989. Floral anatomy of the Pothoideae and Monsteroideae (Araceae). Amer. J. Bot. 76 (Suppl.): 231.
- Chartier, M., Gibernau, M., Renner, S.S., 2014. The evolution of pollinator-plant interaction types in the Araceae. Evolution 68, 1533–1543.

- Clayton, J.W., Soltis, P.S., Soltis, D.E., 2009. Recent long-distance dispersal overshadows ancient biogeographical patterns in a pantropical angiosperm family (Simaroubaceae, Sapindales). *Syst. Biol.* 58, 395–410. doi: 10.1093/sysbio/syp041.
- Christenhusz, M.J.M., Chase, M.W., 2013. Biogeographical patterns of plants in the Neotropics – dispersal rather than plate tectonics is most explanatory. *Bot. J. Linn. Soc.* 171, 277–286. doi:10.1111/j.1095-8339.2012.01301.x.
- Coiffard, C., Mohr, B.A.R., Bernardes-de-Oliveira, M.E.C., 2013. The Early Cretaceous Aroid, *Spixiarum kipea* gen. et sp. nov., and implications on early dispersal and ecology of basal monocots. *Taxon* 62, 997–1008. doi:10.12705/625.21.
- Conti, E., Eriksson, T., Schönenberger, J., Sytsma, K.J., Baum, D A., 2002. Early Tertiary out-of-India dispersal of Crypteroniaceae: evidence from phylogeny and molecular dating. *Evolution* 56, 1931–1942.
- Croat, T., 1979. The distribution of the Araceae. In: *Tropical botany*, eds., Larsen, K., Holm-Nielsen, L.B., Academic Press, London, pp. 291–308.
- Croat, T., 1983. A revision of the genus *Anthurium* (Araceae) of Mexico and Central America. Part 1: Mexico and Middle America. *Ann. Missouri Bot. Gard.* 70: 211–417.
- Croat, T., 1988. The ecology and life forms of Araceae. *Aroideana* 11(3–4): 4–56.
- Croat, T., 1992. Species diversity of Araceae in Colombia —A preliminary survey. *Ann. Mo. Bot. Gard.* 79, 17–28. doi:10.2307/2399806.
- Croat, T., 1998. History and current status of systematic research with Araceae. *Aroideana*, 21, 26-145.

- Cuénoud, P., Savolainen, V., Chatrou, L.W., Powell, M., Grayer, R.J., Chase, M.W., 2002. Molecular phylogenetics of Caryophyllales based on nuclear 18S rDNA and plastid *rbcL*, *atpB*, and *matK* DNA sequences. *Am. J. Bot.* 89, 132–144.
- Cusimano, N., Bogner, J., Mayo, S. J., Boyce, P.C., Wong, S.Y., Hesse, M., Hetterscheid, W.L.A., Keating, R.C., French, J.C., 2011a. Relationships within the Araceae: comparison of morphological patterns with molecular phylogenies. *Am. J. of Bot.* 98, 654–668.
- Cusimano, N., Sousa, A., Renner, S.S., 2011b. Maximum likelihood inference implies a high, not a low, ancestral haploid chromosome number in Araceae, with a critique of the bias introduced by ‘x’. *Ann. Bot.* 109, 681-692. doi: 10.1093/aob/mcr302.
- Davis, C.C., Bell, C.D., Mathews, S., Donoghue, M.J., 2002. Laurasian migration explains Gondwanan disjunctions: Evidence from Malpighiaceae. *Proc. Natl. Acad. Sci. U.S.A.* 99, 6833–6837. doi:10.1073/pnas.102175899.
- De Oliveira, L.L., Calazans, L.S.B., de Morais, É.B., Mayo, S.J., Schrago, C.G., Sakuragui, C.M., 2014. Floral Evolution of *Philodendron* Subgenus *Meconostigma* (Araceae). *PLoS One* 9, e89701. doi:10.1371/journal.pone.0089701.
- Doyle, J.J., Doyle, J.L., 1987. A rapid DNA isolation procedure for small quantities of fresh leaf tissue. *Phytochem. Bull.* 19, 11–15.
- Drummond, A.J., Suchard, M.A., Xie, D., Rambaut, A., 2012. Bayesian phylogenetics with BEAUti and the BEAST 1.7. *Mol. Biol. Evol.* 29, 1969–1973. doi: 10.1093/molbev/mss075.

- Edgar, R.C., 2004. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res.* 32, 1792–1797.
- Engler, A., 1876 . Vergleichende Untersuchungen ü ber die morphologischen Verh ä ltnisse der Araceae. I. Nat ü rliches System der Araceae. *Nova Acta Academiae Caesareae Leopino - Carolinae Germanicae Naturae Curiosorum* 39, 133–155.
- Engler, A., Krause, K., 1908. Araceae-Monsteroideae. In: *Das Pflanzenreich. IV. 23B (Heft 37)*, ed., Engler, A., Leipzig.
- French, J.C., Tomlinson, P.B., 1981. Vascular patterns in stems of Araceae: subfamily Pothoideae. *Am. J. Bot.* 68, 713–729.
- French, J.C., Chung, M.G., Hur, Y.K., 1995. Chloroplast DNA phylogeny of the Ariflorae. In: *Monocotyledons: Systematics and evolution, vol. I*, eds., Rudall, P.J., Cribb, P.J., Cutler, D.F., Humphries, C.J., Royal Botanic Gardens, Kew, UK, 255–275.
- Friis, E.M., Pedersen, K.R., Crane, P.R., 2004. Araceae from the Early Cretaceous of Portugal: Evidence on the emergence of monocotyledons. *Proc. Natl. Acad. Sci. U.S.A.* 101, 16565–16570. doi:10.1073/pnas.0407174101.
- Friis, E.M., Pedersen, K.R., Crane, P.R. 2006. Cretaceous angiosperm flowers: innovation and evolution in plant reproduction. *Palaeogeogr Palaeoclimatol Palaeoecol* 232(2), 251-293.
- Friis, E.M., Pedersen, K.R., Crane, P.R., 2010a. Diversity in obscurity: fossil flowers and the early history of angiosperms. *Philos. T. Roy. Soc. B.* 365, 369–382. doi:10.1098/rstb.2009.0227.

- Friis, E.M., Pedersen, K.R., Crane, P.R., 2010b. Cretaceous diversification of angiosperms in the western part of the Iberian Peninsula. *Review of Palaeobotany and Palynology*, Iberian Floras through Time: Land of Diversity and Survival 162, 341–361. doi:10.1016/j.revpalbo.2009.11.009
- Gauthier, M.P.L., Barabe, D., Bruneau, A., 2008. Molecular phylogeny of the genus *Philodendron* (Araceae): delimitation and infrageneric classification. *Bot. J. Linean Soc.* 156, 13–27. doi: 10.1111/j.1095-8339.2007.00733.x.
- Givnish, T.J., Millam, K.C., Evans, T.M., Hall, J.C., Pires, J.C., Berry, P.E., Sytsma, K.J., 2004. Ancient vicariance or recent long-distance dispersal? inferences about phylogeny and South American–African disjunctions in Rapateaceae and Bromeliaceae based on *ndhF* sequence data. *Int. J. Plant Sci.* 165(S4), S35–S54.
- Gonçalves, E.G., Mayo, S.J., Van Sluys, M.A., Salatino, A., 2007. Combined genotypic–phenotypic phylogeny of the tribe Spathicarpeae (Araceae) with reference to independent events of invasion to Andean regions. *Mol. Phylogenet. Evol.* 43, 1023–1039. doi:10.1016/j.ympev.2007.01.008.
- Grayum, M.H., 1990. Evolution and phylogeny of the Araceae. *Ann. Mo. Bot. Gard.* 77, 628–697.
- Grayum, M.H., 1992 Comparative external pollen ultrastructure of the Araceae and putatively related taxa. *Monogr. Syst. Botan.*, vol 43. Allen Press, Lawrence, Kans.
- Grob, G.B.J., Gravendeel, B., Eurlings, M.C.M., 2004. Potential phylogenetic utility of the nuclear FLORICAULA/LEAFY second intron: comparison with three

- chloroplast DNA regions in *Amorphophallus* (Araceae). *Mol. Phylogenet. Evol.* 30, 13–23. doi:10.1016/S1055-7903(03)00183-0.
- Hay, A., Mabberley, D.J., 1991. Transference of function and the origin of aroids: their significance in early angiosperm evolution. *Bot. Jb*, 113, 339–428.
- Henriquez, C.L., Arias, T., Pires, J.C., Croat, T.B., Schaal, B.A., 2014. Phylogenomics of the plant family Araceae. *Mol. Phylogenet. Evol.* 75, 91–102. doi:10.1016/j.ympev.2014.02.017.
- Herrera, F.A., Jaramillo, C.A., Dilcher, D.L., Wing, S.L., Gómez-N, C., 2008. Fossil Araceae from a Paleocene neotropical rainforest in Colombia. *Am. J. Bot.* 95, 1569–1583. doi:10.3732/ajb.0800172.
- Kappelman, J., Rasmussen, D.T., Sanders, W.J., Feseha, M., Bown, T., Copeland, P., Crabaugh, J., Fleagle, J., Glantz, M., Gordon, A., Jacobs, B., Maga, M., Muldoon, K., Pan, A., Pyne, L., Richmond, B., Ryan, T., Seiffert, E.R., Sen, S., Todd, L. Wiemann, M.C., Winkler, A., 2003. Oligocene mammals from Ethiopia and faunal exchange between Afro-Arabia and Eurasia. *Nature*, 426(6966), 549–552.
- Katoh, K., Misawa, K., Kuma, K., Miyata, T., 2002. MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. *Nucl. Acids. Res.* 30, 3059–3066. doi:10.1093/nar/gkf436.
- Katoh, K., Standley, D.M., 2013. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Mol. Biol. Evol.* 30, 772–780. doi:10.1093/molbev/mst010.
- Keating, R.C., 2002. Leaf anatomical characters and their value in understanding morphoclines in the Araceae. *Bot. Rev.* 68, 510–523.

- Keating, R.C., 2003. *Anatomy of the monocotyledons: Volume IX, Acoraceae and Araceae*. Clarendon Press.
- Keating, R.C., 2004a. Vegetative anatomical data and its relationship to a revised classification of the genera of Araceae. *Ann. Mo. Bot. Gard.* 91, 485–494.
- Keating, R.C., 2004b. Systematic occurrence of raphide crystals in Araceae. *Ann. Mo. Bot. Gard.* 91, 495–504.
- Kissling, W.D., Baker, W.J., Balslev, H., Barfod, A.S., Borchsenius, F., Dransfield, J., Govaerts, R., Svenning, J.-C., 2012a. Quaternary and pre-Quaternary historical legacies in the global distribution of a major tropical plant lineage. *Global Ecol. Biogeogr.* 21, 909–921. doi:10.1111/j.1466-8238.2011.00728.x.
- Kissling, W.D., Eiserhardt, W.L., Baker, W.J., Borchsenius, F., Couvreur, T.L.P., Balslev, H., Svenning, J.-C., 2012b. Cenozoic imprints on the phylogenetic structure of palm species assemblages worldwide. *Proc. Natl. Acad. Sci. U.S.A.* 109, 7379–7384. doi:10.1073/pnas.1120467109.
- Kress, W.J., Erickson, D.L., 2007. A Two-Locus Global DNA Barcode for Land Plants: The coding *rbcL* gene complements the non-coding *trnH-psbA* spacer region. *PLoS One* 2, e508. doi:10.1371/journal.pone.0000508
- Kvaček, J., Smith, S.Y., 2015. *Orontiophyllum*, a new genus for foliage of fossil Orontioideae (Araceae) from the Cretaceous of central Europe. *Bot. J. Linn. Soc.* doi:10.1111/boj.12256.
- Lanfear, R., Calcott, B., Ho, S.Y.W., Guindon, S., 2012. PartitionFinder: Combined selection of partitioning schemes and substitution models for phylogenetic analyses. *Mol. Biol. Evol.* 29, 1695–1701. doi:10.1093/molbev/mss020.

- Loh, J.P., Kiew, R., Hay, A., Kee, A., Gan, L.H., Gan, Y.-Y., 2000. Intergeneric and Interspecific Relationships in Araceae tribe Caladieae and Development of Molecular Markers using Amplified Fragment Length Polymorphism (AFLP). *Ann Bot* 85, 371–378. doi:10.1006/anbo.1999.1074
- Madison, M., 1976. *Alloschemone* and *Scindapsus* (Araceae). *Selbyana* 1, 325–327.
- Madison, M., 1977. A revision of *Monstera* (Araceae). *Contrib. Gray Herb. Harvard Univ.* 207, 3–100.
- Madison, M., Tiffney, B.H., 1976. The seeds of Monstereae: their morphology and fossil record. *J. Arnold Arbor.* 57, 185–204.
- Mansion, G., Rosenbaum, G., Schoenenberger, N., Bacchetta, G., Rosselló, J.A., Conti, E., 2008. Phylogenetic analysis informed by geological history supports multiple, sequential invasions of the Mediterranean Basin by the angiosperm family Araceae. *Syst. Biol.*, 57, 269–285. doi: 10.1080/10635150802044029.
- Matzke, N., 2012. Founder-event speciation in BioGeoBEARS package dramatically improves likelihoods and alters parameter inference in Dispersal-Extinction-Cladogenesis (DEC) analyses. *Front. Biogeogr* 4(suppl. 1): 210
- Matzke, N., 2013. *BioGeoBEARS: BioGeography with Bayesian (and Likelihood) Evolutionary Analysis in R Scripts*. University of California, Berkeley, Berkeley, CA. <http://CRAN.R-project.org/package=BioGeoBEARS>. Accessed Dec 2014
- Mayo, S.J., Bogner, J., Boyce, P.C., 1997. *The genera of Araceae*. Kew: Royal Botanic Gardens, Kew.
- Miller, M.A., Pfeiffer, W., Schwartz, T., 2010. Creating the CIPRES Science Gateway for inference of large phylogenetic trees. In: *Proceedings of the Gateway*

- Computing Environments Workshop (GCE 2010)*. New Orleans, LA: Institute of Electrical and Electronics Engineers (IEEE). Pp. 1-8.
- Nauheimer, L., D. Metzler, Renner, S.S., 2012. Global history of the ancient monocot family Araceae inferred with models accounting for past continental positions and previous ranges based on fossils. *New Phytol.* 195, 938–950.
- Nicolson, D.H., 1960. The occurrence of trichosclereids in the Monsteroideae (Araceae). *Am. J. Bot.* 47, 598-602.
- Nie, Z.-L., Deng, T., Meng, Y., Sun, H., Wen, J., 2013. Post-Boreotropical dispersals explain the pantropical disjunction in *Paederia* (Rubiaceae). *Ann. Bot.* 111, 873–886. doi:10.1093/aob/mct053.
- R Development Core Team, 2008. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. ISBN 3-900051-07-0, URL <http://www.R-project.org>.
- Raven, P.H., Axelrod, D.E., 1974. Angiosperm biogeography and past continental movements. *Ann. Missouri Bot. Gard.* 61, 539–673.
- Ree, R.H., Smith, S.S., 2008. Maximum-likelihood inference of geographic range evolution by dispersal, local extinction, and cladogenesis. *Syst. Biol.* 57, 4–14.
- Ree, R.H., Moore, B.R., Webb, C.O., Donoghue, M.J., 2005. A likelihood framework for inferring the evolution of geographic range on phylogenetic trees. *Evolution* 59, 2299–2311.
- Renner, S., 2004. Plant Dispersal across the Tropical Atlantic by Wind and Sea Currents. *Int. J. Plant Sci.* 165, S23–S33. doi:10.1086/383334.

- Renner, S.S., Clausen, G., Meyer, K., 2001. Historical biogeography of Melastomataceae: the roles of Tertiary migration and long-distance dispersal. *Am. J. Bot.* 88, 1290–1300.
- Renner, S.S., Zhang, L.-B., Murata, J., 2004. A chloroplast phylogeny of *Arisaema* (Araceae) illustrates Tertiary floristic links between Asia, North America, and East Africa. *Am. J. Bot.* 91, 881–888
- Revell, L.J., 2012. phytools: an R package for phylogenetic comparative biology (and other things). *Methods Ecol. Evol.* 3, 217–223. doi:10.1111/j.2041-210X.2011.00169.x.
- Ronquist, F., Teslenko, M., van der Mark, P., Ayres, D.L., Darling, A., Höhna, S., Larget, B., Liu, L., Suchard, M.A., Huelsenbeck, J.P., 2012. MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice across a large model space. *Syst. Biol.* 61, 539–542.
- Schatz, G.E., 1996. Malagasy/Indo-Australo-Malesian phytogeographic connections. In: *Biogeography of Madagascar*, ed., Lourenco, R., ORSTOM editions. Paris.
- Steenis, C. G. G. J. van; 1996. The land-bridge theory in botany. *Blumea*. 1962, 11, 235-372.
- Schott, H.W., 1860. *Prodromus systematis aroidearum*. Congregationis Mechitharisticae, Vienna, Austria.
- Seubert, E., 1997. A comparative study on the seeds of Lasieae (Araceae). *Bot. Jahrb. Syst. Pflanzengesch Pflanzengeogr* 119, 407–426.
- Shaw, J., Lickey, E.B., Beck, J.T., Farmer, S.B., Liu, W., Miller, J., Siripun, K.C., Winder, C.T., Schilling, E.E., Small, R.L., 2005. The tortoise and the hare II:

- relative utility of 21 noncoding chloroplast DNA sequences for phylogenetic analysis. *Am. J. Bot.* 92, 142–166.
- Solerder H. & Meyer F. J. 1930; Systematische Anatomie der Monokotyledonen Heft. VI Microspermae. Gebrüder Borntraeger-Berlin.
- Stamatakis, A., 2006. RAxML-VI-HPC: maximum likelihood-based phylogenetic analyses with thousands of taxa and mixed models. *Bioinformatics* 22, 2688–2690.
- Stockey, R., Hoffman, G., Rothwell, G., 1997. The fossil monocot *Limnobiophyllum scutatum*: resolving the phylogeny of Lemnaceae. *Am. J. Bot.* 84, 355–355.
- Stockey, R.A., Rothwell, G.W., Johnson, K.R., 2007. *Cobbania corrugata* gen. et comb. nov.(Araceae): a floating aquatic monocot from the Upper Cretaceous of western North America. *Am. J. Bot.*, 94, 609–624.
- Sytsma, K.J., Spalink, D., Berger, B., 2014. Calibrated chronograms, fossils, outgroup relationships, and root priors: re-examining the historical biogeography of Geraniales. *Biol. J. Linnean Soc.* 113, 29–49.
- Taberlet, P., Gielly, L., Pautou, G., Bouvet, J., 1991. Universal primers for amplification of three non-coding regions of chloroplast DNA. *Plant Mol. Biol.* 17, 1105–1109.
- Tam, S.M., Boyce, P.C., Upson, T.M., Barabé, D., Bruneau, A., Forest, F., Parker, J.S., 2004. Intergeneric and infrafamilial phylogeny of subfamily Monsteroideae (Araceae) revealed by chloroplast trnL-F sequences. *Am. J. Bot.* 91, 490–498.
- WCSP, 2015. World Checklist of Selected Plant Families. Facilitated by the Royal Botanic Gardens, Kew. Published on the Internet. <http://apps.kew.org/wcsp/> Accessed July 2014.

- Wilde, V., Kvaček, Z., Bogner, J., 2005. Fossil leaves of the Araceae from the European Eocene and notes on other aroid fossils. *Int. J. Plant Sci.* 166, 157–183.
- Wong, S.Y., Boyce, P.C., Othman, A.S.B., Pin, L.C. 2010. Molecular phylogeny of tribe Schismatoglottideae (Araceae) based on two plastid markers and recognition of a new tribe, Philonotieae, from the Neotropics. *Taxon* 59, 117–124.
- Wong, S.Y. T.P. Jean, N.K. Kiaw, A.S.B. Othman, L.H. Boon, F.B. Ahmad & P.C. Boyce. 2013. Phylogeny of Asian *Homalomena* (Araceae) based on the ITS Region Combined with Morphological and Chemical Data. *Syst. Bot.* 38(3): 589-599.
- Zetter, R., Hesse, M., Frosch-Radivo, A., 2001. Early Eocene zona-aperturate pollen grains of the Proxapertites type with affinity to Araceae. *Rev. Palaeobot. Palynol.* 117, 267–279.

TABLES AND FIGURES

Table 1. Voucher information and Genbank accession numbers. NOTE: **This table would be completed with GENBANK accession numbers for publication.**

Species	Voucher	<i>matK</i>	<i>rb</i> <i>cl</i>	<i>trnC-petN</i> spacer	<i>trnL-trnF</i> spacer	<i>ycf</i> <i>1</i>	<i>PH</i> <i>YC</i>	<i>CH</i> <i>S</i>
<i>Alloschemone</i>								
<i>occidentalis</i>	Chase, M. 9996							
<i>Amydrium</i> sp	Hahn 3295							
<i>Amydrium</i>								
<i>hainanense</i>	Chase, M. 9975							
<i>Amydrium</i>								
<i>humile</i>	Chase, M. 9974							
<i>Amydrium</i>								
<i>medium</i>	Chase, M. 9976							
<i>Amydrium</i>								
<i>zippelianum</i>	Zuluaga, A. s.n.							
<i>Anadendrum</i>	Scherberich, D.							
<i>affine</i>	50950							
<i>Anadendrum</i>								
<i>angustifolium</i>	Chase, M. 9985							
<i>Anadendrum</i>								
<i>microstachyum</i>	Chase, M. 9983							
<i>Anthurium_aca</i>								
<i>ule</i>	GENBANK							
<i>Anthurium_and</i>								
<i>raeanum</i>	GENBANK							
<i>Anthurium_bak</i>								
<i>eri</i>	GENBANK							
<i>Anthurium_bren</i>								
<i>esii</i>	GENBANK							
<i>Anthurium_inte</i>								
<i>rruptum</i>	GENBANK							
<i>Anthurium</i>								
<i>ravenii</i>	GENBANK							
<i>Anthurium_scan</i>								
<i>dens</i>	GENBANK							
<i>Anthurium_sche</i>								
<i>rzerianum</i>	GENBANK							
<i>Anthurium_spec</i>								
<i>tabile</i>	GENBANK							
<i>Dracontioides_</i>								
<i>desciscens</i>	GENBANK							
<i>Epipremnum</i>								
<i>aureum</i>	Zuluaga, A s.n							
<i>Epipremnum</i>	Scherberich, D.							
<i>carolinense</i>	010363A							

<i>Epipremnum falcifolium</i>	Chase, M. 9979
<i>Epipremnum papuanum</i>	Chase, M. 9978
<i>Epipremnum pinnatum</i>	Chase, M. 9977
<i>Epipremnum pinnatum</i>	Zuluaga, A. s.n.
<i>Epipremnum amplissimum</i>	GENBANK
<i>Gymnostachys_anceps</i>	GENBANK
<i>Heteropsis flexuosa</i>	Croat, T. 103206
<i>Heteropsis robusta</i>	Betancur, J. 13955
<i>Heteropsis spruceana</i>	Betancur, J. 13851
<i>Heteropsis steyermarkii</i>	Croat, T. 103150
<i>Holochlamys beccarii</i>	Chase, M. 9992
<i>Lasia spinosa</i>	GENBANK
<i>Lemna minor</i>	GENBANK
<i>Lysichiton_americanus</i>	Zuluaga, A. 919
<i>Monstera acacoyaguensis</i>	Croat, T. 71913
<i>Monstera acuminata</i>	Zuluaga, A s.n.
<i>Monstera adansonii</i>	Betancur, J. 13211
<i>Monstera adansonii</i>	Betancur, J. 16021
<i>Monstera aureopinnata</i>	Betancur, J. 13979
<i>Monstera buseyi</i>	Zuluaga, A. 901
<i>Monstera sp nov integrifolia</i>	Zuluaga, A. 916
<i>Monstera deliciosa</i>	Zuluaga, A. 902
<i>Monstera dissecta</i>	Zuluaga, A. 911
<i>Monstera dubia</i>	Zuluaga, A. s.n.
<i>Monstera egregia</i>	Croat, T. 100357a
<i>Monstera epipremnoides</i>	Croat, T. 69750
<i>Monstera filamentosa</i>	Zuluaga, A. 893
<i>Monstera florescanoana</i>	Zuluaga, A. s.n.
<i>Monstera gracilis</i>	Zuluaga, A. 463
<i>Monstera</i>	Zuluaga, A. 311

<i>lechleriana</i>	
<i>Monstera</i>	
<i>obliqua</i>	Clavijo, L 1565
<i>Monstera</i>	
<i>oreophila</i>	Zuluaga, A. 886
<i>Monstera_pichi</i>	
<i>nchensis</i>	Croat, T. 96437
<i>Monstera</i>	
<i>pinnatipartita</i>	Zuluaga, A. 874
<i>Monstera</i>	
<i>pittieri</i>	Zuluaga, A. 894
<i>Monstera</i>	
<i>planadensis</i>	Zuluaga, A. 351
<i>Monstera</i>	
<i>punctulata</i>	Zuluaga, A. 887
<i>Monstera</i>	
<i>siltepecana</i>	Croat, T. 90214
<i>Monstera</i>	
<i>spruceana</i>	Zuluaga, A. 899
<i>Monstera</i>	
<i>subpinnata</i>	Zuluaga, A. 417
<i>Monstera</i>	
<i>tuberculata</i>	Zuluaga, A s.n.
<i>Monstera</i>	
<i>xanthospatha</i>	Zuluaga, A. 884
<i>Orontium_aqua</i>	
<i>ticum</i>	GENBANK
<i>Pedicellarum_p</i>	
<i>aiei</i>	GENBANK
<i>Pothoidium_lob</i>	
<i>bianum</i>	GENBANK
<i>Pothos_chinensi</i>	
<i>s</i>	GENBANK
<i>Pothos_scanden</i>	
<i>s</i>	GENBANK
<i>Pothos_tener</i>	GENBANK
<i>Rhaphidophora</i>	
<i>africana</i>	Chase, M. 8595
<i>Rhaphidophora</i>	
<i>sp</i>	Chase, M. 8597
<i>Rhaphidophora</i>	
<i>angustata</i>	Chase, M. 8628
<i>Rhaphidophora</i>	
<i>beccarii</i>	Chase, M. 8626
<i>Rhaphidophora</i>	
<i>crassicaulis</i>	Chase, M. 8630
<i>Rhaphidophora</i>	
<i>decursiva</i>	Chase, M. 8601
<i>Rhaphidophora</i>	
<i>guamensis</i>	Chase, M. 8625
<i>Rhaphidophora</i>	
<i>hongkongensis</i>	Chase, M. 8002
<i>Rhaphidophora</i>	
<i>hookeri</i>	Chase, M. 8632
<i>Rhaphidophora</i>	Chase, M. 8604

<i>lobbii</i>	
<i>Rhaphidophora</i>	
<i>luchunensis</i>	Chase, M. 8634
<i>Rhaphidophora</i>	
<i>maingayi</i>	Chase, M. 8599
<i>Rhaphidophora</i>	
<i>megaphylla</i>	Chase, M. 8627
<i>Rhaphidophora</i>	
<i>montana</i>	Croat, T. 59140b
<i>Rhaphidophora</i>	
<i>peepla</i>	Chase, M. 7382
<i>Rhaphidophora</i>	
<i>schlechteri</i>	Chase, M. 8637
<i>Rhaphidophora</i>	
<i>spathacea</i>	Chase, M. 8624
<i>Rhaphidophora</i>	
<i>spuria</i>	Chase, M. 8608
<i>Rhaphidophora</i>	
<i>sylvestris</i>	Chase, M. 7398
<i>Rhaphidophora</i>	
<i>tetrasperma</i>	Chase, M. 8629
<i>Rhaphidophora</i>	
<i>korthalsii</i>	Chase, M. 8635
<i>Rhaphidophora</i>	
<i>latevaginata</i>	Chase, M. 8633
<i>Rhaphidophora</i>	
<i>neoguineensis</i>	Chase, M. 8636 Betancur, J. 15319
<i>Rhodospatha</i>	
<i>Rhodospatha</i>	
<i>moritziana</i>	Zuluaga, A. 565 Betancur, J. 14578
<i>Rhodospatha</i>	
<i>Rhodospatha</i>	Zuluaga, A. 567
<i>Rhodospatha</i>	Zuluaga, A. 567
<i>Rhodospatha</i>	
<i>latifolia</i>	Zuluaga, A. 460
<i>Rhodospatha</i>	
<i>venosa</i>	Croat, T. 104784
<i>Rhodospatha</i>	
<i>wendlandii</i>	Zuluaga, A. 551
<i>Scindapsus</i>	
<i>hederaceus</i>	Chase, M. 9986
<i>Scindapsus</i>	
<i>rupestris</i>	Chase, M. 8623
<i>Spathiphyllum</i>	
<i>cannifolium</i>	Zuluaga, A. 532
<i>Spathiphyllum</i>	
<i>cochlearispathu</i>	
<i>m</i>	Croat, T. 73887
<i>Spathiphyllum</i>	
<i>minor</i>	Betancur, J. 14022
<i>Spathiphyllum</i>	
<i>solomonense</i>	Lavarack 31268
<i>Spathiphyllum</i>	Croat, T. 78313

<i>wallisii</i>	
<i>Spathiphyllum</i> <i>cannifolium</i>	Betancur, J. 14976
<i>Spathiphyllum</i> <i>commutatum</i>	Medecilo, M. 498
<i>Spathiphyllum</i> <i>patulinervum</i>	GENBANK
<i>Spathiphyllum</i> <i>pygmaeum</i>	GENBANK
<i>Spirodela</i> <i>polyrhiza</i>	GENBANK
<i>Stenospermatio</i> <i>n</i>	Zuluaga, A. 371
<i>Stenospermatio</i> <i>n</i>	Zuluaga, A. 756
<i>Stenospermatio</i> <i>n angustifolium</i>	Zuluaga, A. 553
<i>Stenospermatio</i> <i>n</i>	Zuluaga, A. 900
<i>Stenospermatio</i> <i>n andreanum</i>	Zuluaga, A. 321
<i>Stenospermatio</i> <i>n ellipticum</i>	Zuluaga, A. 379
<i>Stenospermatio</i> <i>n</i>	Zuluaga, A. 518
<i>Stenospermatio</i> <i>n escobariae</i>	Zuluaga, A. 496
<i>Stenospermatio</i> <i>n multiovulatum</i>	GENBANK

Table 2. Primers design for *ycf1*, PHYC and CHS.

Region	Primer sequence
<i>ycf1</i> Forward	TCCRYGATATGGTCCATTCAA
<i>ycf1</i> Reverse	TYMTGGATACGGCAAAATCA
phyC Forward	GCCTGTGAATTTTTGATGCA
phyC Reverse	AAGAACGTGGATGCATCTTC
CHS Forward	GCCCAMTAAAAATAACAATGCG
CHS Reverse	CCTGCTGGTACATCATGAG

Table 3. DNA substitutions models and partitioning scheme and partition subsets resulting for RAXML and BEAST analyses from partition analyses conducted in PartitionFinder using the BIC criterion.

	Partition number	Best Model	Partition subsets
RAxML	1	GTR+I+G	matK_pos1, matk_pos2, matK_pos3, rbcL_pos2, trnL-F, ycf1
	2	GTR+G	matk-trnK intron, trnC-petN
	3	GTR+G	rbcL_pos1, rbcL_pos3
MrBayes	1	GTR+I+G	matK_pos1, matk_pos2, matK_pos3, rbcL_pos2, trnL-F, ycf1
	2	GTR+G	matk-trnK intron, trnC-petN
	3	K80+I+G	rbcL_pos1, rbcL_pos3

Table 4. Information about the different plastid, nuclear, and combined datasets and alignments. Number of taxa, the length of the alignment, the number of characters excluded, and percentage of missing data and variable sites (for all taxa and the ingroup) are shown.

	Plastid					Nuclear				
	<i>rbcl</i>	<i>matK</i>	<i>trnC-petN</i>	spacer	<i>trnL-trnF</i>	partial <i>ycfI</i>	Concatenated plastid	PHYC	CHS	Concatenated
Number of taxa	98	99	107		106	90	125	79	72	133
Alignment length (bp)	1376	1810	1071		1198	878	6192	1093	692	7977
Excluded characters	0	0	0		101	40	141	0	54	NA
% gaps and missing data	8	14.8	30.1		32	10.2	38	16.8	0.3	49
% variable sites (ingroup)	8.3	51.7	48.7		32.5	47	90.7	34.5	35.8	89.9
% variable sites (all)	20.7	68.2	92.8		74.6	55.6	96.6	54.2	40.8	92.6

Table 5. Support information for the genera and clades within Monsteroideae with different datasets from different phylogenetic analyses. Supports higher than 70 for Maximum likelihood (ML) partition and unpartition analysis, and higher than 0.9 posterior probabilities for Bayesian inference (BI) are shown. Clades represented by one species are considered not tested unless they are nested in another clade.

Genus	Combined plastid	<i>CHS</i>	<i>PHYC</i>	Concatenated
<i>Alloschemone</i>	not tested	not tested	not tested	not tested
<i>Amydrium</i>	ML: 89, 90; BI: 0.99	not monophyletic	not monophyletic	ML: 79, 61
<i>Anadendrum</i>	ML: 100, 99; BI: 1	ML: 100, 100; BI: 1	ML: 100, 100; BI: 1	ML: 100, 100
<i>Epipremnum</i>	not monophyletic	not monophyletic	ML: 91, 91; BI: 1	not monophyletic
<i>Heteropsis</i>	not supported	ML: 98, 100; BI: 1	not monophyletic	not supported
<i>Holochlamys</i>	Nested in Spathiphyllum	not tested	not tested	Nested in Spathiphyllum
<i>Monstera</i>	ML: 94, 96; BI: 1	not supported	not tested	ML: 97, 94
<i>Rhaphidophora</i>	ML: not supported; BI: 0.99	not monophyletic	not monophyletic	not monophyletic
<i>Rhodospatha</i>	ML: 82, 79; BI: 1	ML: not supported, 92; BI: 0.99	not supported	ML: 94, 94
<i>Scindapsus</i>	ML: 96, 94; BI: 0.99	not tested	ML: 100, 100; BI: 1	ML: 96, 96
<i>Spathiphyllum</i>	not monophyletic	not tested	ML: 100, 100; BI: 1	not monophyletic
<i>Stenospermation</i>	ML: 100, 100; BI: 1	ML: 77, 86; BI: 0.99	ML: 95, 94; BI: 1	not supported
<i>Spathiphyllaeae</i>	ML: 100, 100; BI: 1	ML: 100, 100; BI: 1	ML: 100, 100; BI: 1	ML: 100, 100
<i>Heteropsis clade</i>	ML: 100, 100; BI: 1	ML: not supported, 88; BI: 0.99	ML: 100, 100; BI: 1	ML: 99, 100
<i>Rhaphidophora clade</i>	ML: 100, 100; BI: 1	ML: not supported, 72; BI: 0.99	ML: 100, 100; BI: 1	ML: 100, 100
<i>Rhaphidophora clade + Heteropsis clade</i>	ML: 99, 100; BI: 1	not supported	ML: 99, 100; BI: 1	ML: 100, 100
Monsteroideae	ML: 100, 100; BI: 1	not supported	ML: 100, 100; BI: 1	ML: 100, 100; BI: 1

Table 6. Information about BEAST analysis. Node number according to **Figure 4**, age in Ma, and 5-95 % Highest Posterior Density (HPD) intervals are shown.

Node	Age	CI (5-95%)	Node	Age	CI (5-95%)
126	125.44	(114.85-134.99)	170	28.92	(13.78-48.95)
127	115.24	(109.2-125.83)	189	21.99	(8.77-39.33)
128	108.84	(94.67-121.62)	190	59.79	(28.55-87.76)
129	90.77	(78.26-104.08)	169	125.44	(114.85-134.99)
130	68.83	(64.2-76.73)	192	115.24	(109.2-125.83)
131	59.08	(45.75-70.5)	132	59.08	(45.75-70.5)
132	38.45	(27.68-50.89)	194	38.45	(27.68-50.89)
133	36.75	(26.02-48.07)	195	36.75	(26.02-48.07)
134	28.01	(18.71-38.85)	196	28.01	(18.71-38.85)
135	18.67	(11.63-27.07)	194	6.3	(3.51-9.99)
136	15.69	(9.76-22.92)	131	0.85	(0.1-2.23)
137	9.8	(5.78-14.71)	199	0.5	(0-1.63)
138	7.79	(4.61-11.85)	200	0.23	(0-1.13)
139	6.3	(3.51-9.99)	201	0.47	(0-1.84)
140	4.72	(2.16-8.03)	202	4.25	(1.59-7.46)
141	1.74	(0.45-3.71)	203	1.55	(0.19-3.8)
142	0.85	(0.1-2.23)	204	5.18	(2.29-8.57)
143	0.5	(0-1.63)	203	8.19	(4.4-12.9)
141	5.18	(2.29-8.57)	201	2.14	(0.56-4.51)
139	8.19	(4.4-12.9)	207	2.2	(0.06-6.72)
146	6.26	(2.86-10.39)	200	3.98	(1.07-8.16)
138	2.2	(0.06-6.72)	209	17.13	(9.54-26.27)
148	14.57	(8.26-21.81)	210	12.48	(6.33-20.16)
149	8.35	(3.92-14.07)	211	10	(4.67-16.73)
148	12.48	(6.33-20.16)	212	7.06	(2.75-12.67)
137	2.67	(0.32-6.48)	211	32.29	(23.06-43.58)
152	8.53	(2.99-16.05)	209	19.9	(13.19-28.61)
153	1.72	(0.15-4.94)	199	9.84	(5.48-15.44)
154	32.29	(23.06-43.58)	216	7.02	(3.5-11.66)
155	29.05	(20.49-39.42)	217	5.5	(2.16-10.1)
153	13.15	(7.32-20.02)	218	3.26	(0.78-6.86)
135	0.86	(0-3.55)	130	12.4	(6.47-20.19)
158	3.23	(0-9.3)	220	9.13	(4.17-16.39)
159	13.96	(6.69-21.91)	221	7.01	(2.91-12.62)
160	3.15	(0.57-7.64)	222	3.11	(0.69-7.02)
134	11.66	(4.93-20.87)	222	5.72	(1.66-11.95)
162	8.01	(3.04-15.48)	221	17.03	(8.05-29.26)
163	5.72	(1.66-11.95)	225	5.57	(1.8-11.45)
164	7.73	(2.48-16.03)	226	2.33	(0.48-5.52)
165	5.09	(1.26-11.25)	220	21.17	(13.53-30.62)
162	38.56	(26.93-51.61)	129	11.5	(6.13-18.32)
167	34.86	(24.06-47.71)	229	7	(2.65-12.74)
133	11.5	(6.13-18.32)	230	3.65	(0.43-9.63)
169	7	(2.65-12.74)	231	12.67	(5.52-21.82)
170	3.65	(0.43-9.63)	232	5.85	(1.62-12.26)

171	12.67	(5.52-21.82)	233	23.68	(12.38-36.67)
172	5.85	(1.62-12.26)	231	5.95	(1.65-12.16)
173	23.68	(12.38-36.67)	235	0.89	(0.01-3.66)
174	18.11	(8.99-29.81)	230	0.38	(0-2.04)
175	10.11	(4.3-18.29)	229	11.14	(4.66-19.79)
176	1.56	(0.14-4.51)	238	6.06	(0.69-13.5)
177	0.34	(0-1.62)	239	4.59	(1.2-9.83)
178	5.95	(1.65-12.16)	239	2.22	(0-8.43)
179	0.89	(0.01-3.66)	128	19.53	(10.39-30.64)
176	21.36	(11.75-34.36)	242	14.76	(6.74-24.7)
173	7.66	(2.43-15.16)	243	9.33	(2.89-17.74)
182	2.22	(0-8.43)	242	10.2	(3.01-20.24)
171	14.76	(6.74-24.7)	127	17.18	(7.13-29.04)
184	9.33	(2.89-17.74)	246	11.65	(3.47-22.46)
185	3.34	(0-10.34)	126	59.79	(28.55-87.76)
186	12.5	(4.08-23.44)	248	73.42	(53.71-90.98)
187	5.71	(0.8-14.25)	249	34.09	(16.42-57.14)

Table 7. Information from selected nodes about ancestral range estimation over 100 trees from BEAST posterior distribution using seven areas and DEC model in BioGeoBears. Columns show the most probable range using the chronogram from BEAST; the probability of that range; the frequency of the range as the most probable (P>50) over 100 trees chosen from posterior distribution from BEAST; and the average and standard deviation (SD) for the probability of that range over 100 trees. Clades in bold were constrained in the BEAST analysis.

Node	Clade	Most probable range	Probability on chronogram	Frequency over 100 trees P>50	Average and SD over 100 trees
126	Monsteroideae + Pothoideae	BC or BCGH	0.19	0	0.19(0.09)
127	Monsteroideae	BC	0.64	100	0.61(0.09)
128	<i>Rhaphidophora</i> + <i>Heteropsis</i> clades	BC	0.57	58	0.51(0.13)
129	<i>Rhaphidophora</i> clade	F	0.54	24	0.31(0.22)
132	<i>Monstera</i>	BC	1	100	1(0.001)
177	<i>Rhaphidophora africana</i>	ED	0.68	87	0.64(0.09)
194	Heteropsis clade	BC	0.99	100	0.99(0.002)
215	Spathiphyllae	BCG	0.77	93	0.74(0.19)
164	<i>Rhaphidophora</i> + <i>Anandendrum</i>	F	0.96	100	0.91(0.08)

Table 8. Morphological characters used in the ancestral state reconstruction.

Pearianth and endosperm are coded as absent-present; locule number as 1, 2, 3; and seed number as 1, 2, 3 (representing 3-8 seeds) and 4 (representing >8 seeds).

Species	Perianth	SEED		SEED_Number
		endosperm	# Locules	
<i>Alloschemone_occidentalis</i>	Absent	NA	1	NA
<i>Amydrium_sp</i>	Absent	NA	NA	NA
<i>Amydrium_hainanense</i>	Absent	Present	1	2
<i>Amydrium_humile</i>	Absent	Present	1	2
<i>Amydrium_medium</i>	Absent	Present	1	2
<i>Amydrium_zippelianum</i>	Absent	Present	1	2
<i>Anadendrum_affine</i>	Present	Absent	1	1
<i>Anadendrum_angustifolium</i>	Present	Absent	1	1
<i>Anadendrum_microstachyum</i>	Present	Absent	1	1
<i>Anthurium_acaule</i>	Present	Present	2	4
<i>Anthurium_andraeanum</i>	Present	Present	2	4
<i>Anthurium_bakeri</i>	Present	Present	2	4
<i>Anthurium_brenesii</i>	Present	Present	2	4
<i>Anthurium_interruptum</i>	Present	Present	2	4
<i>Anthurium_ravenii</i>	Present	Present	2	4
<i>Anthurium_scandens</i>	Present	Present	2	4
<i>Anthurium_scherzerianum</i>	Present	Present	2	4
<i>Anthurium_spectabile</i>	Present	Present	2	4
<i>Dracontioides_desciscens</i>	Present	Present	2	2
<i>Epipremnum_aureum</i>	Absent	Present	1	2
<i>Epipremnum_carolinense</i>	Absent	Present	1	4
<i>Epipremnum_falcifolium</i>	Absent	Present	1	2
<i>Epipremnum_papuanum</i>	Absent	Present	1	4
<i>Epipremnum_pinnatum</i>	Absent	Present	1	4
<i>Epipremnum_pinnatum2</i>	Absent	Present	1	4
<i>Epipremnum_amplissimum</i>	Absent	Present	1	4
<i>Gymnostachys_anceps</i>	Present	Present	1	1
<i>Heteropsis_flexuosa</i>	Absent	Absent	2	2
<i>Heteropsis_robusta</i>	Absent	Absent	2	4
<i>Heteropsis_spruceana</i>	Absent	Absent	2	2
<i>Heteropsis_steyermarkii</i>	Absent	Absent	2	2
<i>Holochlamys_beccarii</i>	Present	Present	1	4
<i>Lasia_spinosa</i>	Present	Present	1	1
<i>Lemna_minor</i>	Absent	Present	1	4

<i>Lysichiton_americanus</i>	Present	Absent	2	4
<i>Monstera_acacoyaguensis</i>	Absent	Absent	2	1
<i>Monstera_acuminata</i>	Absent	Absent	2	1
<i>Monstera_adansonii</i>	Absent	Absent	2	1
<i>Monstera_adansonii2</i>	Absent	Absent	2	1
<i>Monstera_aureopinnata</i>	Absent	Absent	2	1
<i>Monstera_buseyi</i>	Absent	Absent	2	1
<i>Monstera_sp_nov1</i>	Absent	Absent	2	1
<i>Monstera_deliciosa</i>	Absent	Absent	2	1
<i>Monstera_dissecta</i>	Absent	Absent	2	1
<i>Monstera_dubia</i>	Absent	Absent	2	1
<i>Monstera_egregia</i>	Absent	Absent	2	1
<i>Monstera_epipremnoides</i>	Absent	Absent	2	1
<i>Monstera_filamentosa</i>	Absent	Absent	2	1
<i>Monstera_florescanoana</i>	Absent	Absent	2	1
<i>Monstera_gracilis</i>	Absent	Absent	2	1
<i>Monstera_lechleriana</i>	Absent	Absent	2	1
<i>Monstera_obliqua</i>	Absent	Absent	2	1
<i>Monstera_oreophila</i>	Absent	Absent	2	1
<i>Monstera_spnov2</i>	Absent	Absent	2	1
<i>Monstera_pinnatipartita</i>	Absent	Absent	2	1
<i>Monstera_pittieri</i>	Absent	Absent	2	1
<i>Monstera_planadensis</i>	Absent	Absent	2	1
<i>Monstera_punctulata</i>	Absent	Absent	2	1
<i>Monstera_siltepecana</i>	Absent	Absent	2	1
<i>Monstera_spruceana</i>	Absent	Absent	2	1
<i>Monstera_subpinnata</i>	Absent	Absent	2	1
<i>Monstera_tuberculata</i>	Absent	Absent	2	1
<i>Monstera_xanthospatha</i>	Absent	Absent	2	1
<i>Orontium_aquaticum</i>	Present	Absent	1	1
<i>Pedicellarum_paiei</i>	Present	Absent	3	NA
<i>Pothoidium_lobbianum</i>	Present	Absent	1	NA
<i>Pothos_chinensis</i>	Present	Absent	3	NA
<i>Pothos_scandens</i>	Present	Absent	3	NA
<i>Pothos_tener</i>	Present	Absent	3	NA
<i>Rhaphidophora_africana</i>	Absent	Present	1	10
<i>Rhaphidophora_sp</i>	Absent	Present	1	10
<i>Rhaphidophora_angustata</i>	Absent	Present	1	10
<i>Rhaphidophora_beccarii</i>	Absent	Present	1	10
<i>Rhaphidophora_crassicaulis</i>	Absent	Present	1	10
<i>Rhaphidophora_decursiva</i>	Absent	Present	1	10
<i>Rhaphidophora_guamensis</i>	Absent	Present	1	10
<i>Rhaphidophora_hongkongensis</i>	Absent	Present	1	10

<i>Rhaphidophora_hookeri</i>	Absent	Present	1	10
<i>Rhaphidophora_lobbii</i>	Absent	Present	1	10
<i>Rhaphidophora_luchunensis</i>	Absent	Present	1	10
<i>Rhaphidophora_maingayi</i>	Absent	Present	1	10
<i>Rhaphidophora_megaphylla</i>	Absent	Present	1	10
<i>Rhaphidophora_montana</i>	Absent	Present	1	10
<i>Rhaphidophora_peepla</i>	Absent	Present	1	10
<i>Rhaphidophora_schlechteri</i>	Absent	Present	1	10
<i>Rhaphidophora_spathacea</i>	Absent	Present	1	10
<i>Rhaphidophora_spuria</i>	Absent	Present	1	10
<i>Rhaphidophora_sylvestris</i>	Absent	Present	1	10
<i>Rhaphidophora_tetrasperma</i>	Absent	Present	1	10
<i>Rhaphidophora_korthalsii</i>	Absent	Present	1	10
<i>Rhaphidophora_latevaginata</i>	Absent	Present	1	10
<i>Rhaphidophora_neoguineensis</i>	Absent	Present	1	10
<i>Rhodospatha_sp1</i>	Absent	Present	2	10
<i>Rhodospatha_moritziana</i>	Absent	Present	2	10
<i>Rhodospatha_sp2</i>	Absent	Present	2	10
<i>Rhodospatha_sp3</i>	Absent	Present	2	10
<i>Rhodospatha_sp4</i>	Absent	Present	2	10
<i>Rhodospatha_latifolia</i>	Absent	Present	2	10
<i>Rhodospatha_venosa</i>	Absent	Present	2	10
<i>Rhodospatha_wendlandii</i>	Absent	Present	2	10
<i>Scindapsus_hederaceus</i>	Absent	Present	1	1
<i>Scindapsus_rupestris</i>	Absent	Present	1	1
<i>Spathiphyllum_cannifolium</i>	Present	Present	3	10
<i>Spathiphyllum_cochlearispathu</i> <i>m</i>	Present	Present	3	10
<i>Spathiphyllum_minor</i>	Present	Present	3	10
<i>Spathiphyllum_solomonense</i>	Present	Present	3	10
<i>Spathiphyllum_wallisii</i>	Present	Present	3	10
<i>Spathiphyllum_cannifolium2</i>	Present	Present	3	10
<i>Spathiphyllum_commutatum</i>	Present	Present	3	10
<i>Spathiphyllum_patulinervum</i>	Present	Present	3	10
<i>Spathiphyllum_pygmaeum</i>	Present	Present	3	10
<i>Spirodela_polyrhiza</i>	Absent	Present	1	4
<i>Stenospermatum_sp1</i>	Absent	Present	2	10
<i>Stenospermatum_sp2</i>	Absent	Present	2	10
<i>Stenospermatum_angustifolium</i>	Absent	Present	2	10
<i>Stenospermatum_sp3</i>	Absent	Present	2	10
<i>Stenospermatum_andreanum</i>	Absent	Present	2	10
<i>Stenospermatum_ellipticum</i>	Absent	Present	2	10
<i>Stenospermatum_sp4</i>	Absent	Present	2	10

<i>Stenospermation_escobariae</i>	Absent	Present	2	10
<i>Stenospermation_multiovulatum</i>	Absent	Present	2	10

Figure 1. Cladogram depicting the topology obtained from Maximum Likelihood analyses using RAxML for a dataset including five palstid regions (*matK*, *rbcl*, *trnC-petN* spacer, *trnL-trnF* spacer, and partial *ycfI*). Bootstrap support values from partitioned analysis are show above the branches, and from unpartitioned below. Inset phylogram indicates branch lengths.




Figure 2. Cladogram depicting the topology obtained from Maximum Likelihood analyses using RAxML for a dataset including the nuclear gene Chalcone synthase (CHS). Bootstrap support values from partitioned analysis are show above the branches, and from unpartitioned below. Inset phylogram indicates branch lengths.




Figure 3. Cladogram depicting the topology obtained from Maximum Likelihood analyses using RAxML for a dataset including the nuclear gen phytochrome C (PHYC). Bootstrap support values from partitioned analysis are show above the branches, and from unpartitioned below. Inset phylogram indicates branch lengths.




Figure 4. Cladogram depicting the topology obtained from Maximum Likelihood analyses using RAxML for a concatenated dataset (plastid and nuclear regions). Bootstrap support values from partitioned analysis are show above the branches, and from unpartitioned below. Inset phylogram indicates branch lengths.




Figure 5. Phylogenetic relationships among clades and genera within Monsteroideae resulted from plastid and nuclear regions datasets. **A**, Phylogram depicting the topology obtained from Maximum Likelihood analyses using RAxML for a dataset including five plastid regions (*matK*, *rbcl*, *trnC-petN* spacer, *trnL-trnF* spacer, and partial *ycf1*); Clades and genera within Monsteroideae are indicated with bars. **B-C**, reduced phylograms showing the topologies resulted from the nuclear regions CHS and PHYC respectively. Bootstrap support values are indicated by circles; white < 75, grey <=75<95, black >=95. Posterior probabilities from Bayesian inference >= 95 are indicated in grey triangles.




Figure 6. Chronogram from BEAST, indicating node numbers. 95% HPD are indicated with blue bars over the nodes.





Figure 7. Ancestral range estimation using the DEC model, four time periods, and seven areas as implemented in BioGeoBears. **A**, squares represent the most probable range at nodes and corners. **B**, pies represents all probable ranges at nodes and corners. Inset map show the different areas: Nearctic (A), Tropical America (BC; C-B-Central America, C South America in the eight area analysis), Palearctic (D), Africa (E), SouthEast Asia (F), Australasia (G), India (H).







Figure 8. Ancestral range estimation using the DEC model, four time periods, and eight areas as implemented in BioGeoBears. **A**, squares represent the most probable range at nodes and corners. **B**, pies represents all probable ranges at nodes and corners. Inset map show the different areas: Nearctic (A), Tropical America (BC; C-B-Central America, C South America in the eight area analysis), Palearctic (D), Africa (E), SouthEast Asia (F), Australasia (G), India (H).






Figure 9. Ancestral state reconstruction for the characters number of locules (top) and number of seeds (bottom) in the subfamily Monsteroideae. Figures to the left show reconstructions using the rerooting method (ML); figures to the right show reconstruction using 1000 stochastic mappings.




Figure 10. Ancestral state reconstruction for the characters presence of perianth (top) and the presence of endosperm (bottom) in the subfamily Monsteroideae. Figures to the left show reconstructions using the rerooting method (ML); figures to the right show reconstruction using 1000 stochastic mappings.








Figure 11. Morphometric analyses from seed shape and size in Monsteroideae.

A-B, scatter plots showing the PC1 and PC3 of the resulting form a principal component analysis on the harmonics produce from the outline analysis (elliptical fourier analysis). **A** is color code by clade in Monsteroideae and **B** is color code by genus. **C-D**, scatter plots showing the PC1 and PC2 from the same analysis; points are color code in the same way. **E**, panel showing outlines for the species of Monsteroideae used in the outline analysis. **F**, 3D-scatter plot using PC1, PC2 and length of the seed (in lateral view) with ellipsoids and color code by clade.



SUPPLEMENTARY TABLES AND FIGURES

TABLE S1. Area coding for each taxa used in the BioGeoBears analyses. Areas are coded as: Nearctic (A), Tropical America (BC; C-B-Central America, C South America in the eight area analysis), Palearctic (D), Africa (E), SouthEast Asia (F), Australasia (G), India (H).

Species Name	A	BC	D	E	F	G	H
<i>Alloschemone occidentalis</i>	0	1	0	0	0	0	0
<i>Amydrium_sp</i>	0	0	0	0	1	0	0
<i>Amydrium_hainanense</i>	0	0	1	0	1	0	0
<i>Amydrium_humile</i>	0	0	0	0	1	0	0
<i>Amydrium_medium</i>	0	0	0	0	1	0	0
<i>Amydrium_zippelianum</i>	0	0	0	0	1	1	0
<i>Anadendrum_affine</i>	0	0	0	0	1	0	0
<i>Anadendrum_angustifolium</i>	0	0	0	0	1	0	0
<i>Anadendrum_microstachyum</i>	0	0	0	0	1	1	0
<i>Anthurium_acaule</i>	0	1	0	0	0	0	0
<i>Anthurium_andraeanum</i>	0	1	0	0	0	0	0
<i>Anthurium_bakeri</i>	0	1	0	0	0	0	0
<i>Anthurium_brenesii</i>	0	1	0	0	0	0	0
<i>Anthurium_interruptum</i>	0	1	0	0	0	0	0
<i>Anthurium_ravenii</i>	0	1	0	0	0	0	0
<i>Anthurium_scandens</i>	0	1	0	0	0	0	0
<i>Anthurium_scherzerianum</i>	0	1	0	0	0	0	0
<i>Anthurium_spectabile</i>	0	1	0	0	0	0	0
<i>Dracontiodides_desciscens</i>	0	1	0	0	0	0	0
<i>Epipremnum_aureum</i>	0	0	0	0	0	1	0
<i>Epipremnum_carolinense</i>	0	0	0	0	0	1	0
<i>Epipremnum_falcifolium</i>	0	0	0	0	1	0	0
<i>Epipremnum_papuanum</i>	0	0	0	0	0	1	0
<i>Epipremnum_pinnatum</i>	0	0	0	0	1	1	1
<i>Epipremnum_amplissimum</i>	0	0	0	0	0	1	0
<i>Gymnostachys_anceps</i>	0	0	0	0	0	1	0
<i>Heteropsis_flexuosa</i>	0	1	0	0	0	0	0
<i>Heteropsis_robusta</i>	0	1	0	0	0	0	0
<i>Heteropsis_spruceana</i>	0	1	0	0	0	0	0
<i>Heteropsis_steyermarkii</i>	0	1	0	0	0	0	0
<i>Holochlamys_beccarii</i>	0	0	0	0	0	1	0
<i>Lasia_spinosa</i>	0	0	0	0	1	0	0

<i>Lemna_minor</i>	1	1	1	1	1	1	1
<i>Lysichiton_americanus</i>	1	0	1	0	0	0	0
<i>Monstera_acacoyaguensis</i>	0	1	0	0	0	0	0
<i>Monstera_acuminata</i>	0	1	0	0	0	0	0
<i>Monstera_adansonii2</i>	0	1	0	0	0	0	0
<i>Monstera_aureopinnata</i>	0	1	0	0	0	0	0
<i>Monstera_buseyi</i>	0	1	0	0	0	0	0
<i>Monstera_sp_nov1</i>	0	1	0	0	0	0	0
<i>Monstera_deliciosa</i>	0	1	0	0	0	0	0
<i>Monstera_dissecta</i>	0	1	0	0	0	0	0
<i>Monstera_dubia</i>	0	1	0	0	0	0	0
<i>Monstera_egregia</i>	0	1	0	0	0	0	0
<i>Monstera_epipremnoides</i>	0	1	0	0	0	0	0
<i>Monstera_filamentosa</i>	0	1	0	0	0	0	0
<i>Monstera_florescanoana</i>	0	1	0	0	0	0	0
<i>Monstera_gracilis</i>	0	1	0	0	0	0	0
<i>Monstera_lechleriana</i>	0	1	0	0	0	0	0
<i>Monstera_obliqua</i>	0	1	0	0	0	0	0
<i>Monstera_oreophila</i>	0	1	0	0	0	0	0
<i>Monstera_spnov2</i>	0	1	0	0	0	0	0
<i>Monstera_pinnatipartita</i>	0	1	0	0	0	0	0
<i>Monstera_pittieri</i>	0	1	0	0	0	0	0
<i>Monstera_planadensis</i>	0	1	0	0	0	0	0
<i>Monstera_punctulata</i>	0	1	0	0	0	0	0
<i>Monstera_siltepecana</i>	0	1	0	0	0	0	0
<i>Monstera_spruceana</i>	0	1	0	0	0	0	0
<i>Monstera_subpinnata</i>	0	1	0	0	0	0	0
<i>Monstera_tuberculata</i>	0	1	0	0	0	0	0
<i>Monstera_xanthospatha</i>	0	1	0	0	0	0	0
<i>Orontium_aquaticum</i>	1	0	0	0	0	0	0
<i>Pedicellarum_paiei</i>	0	0	0	0	1	0	0
<i>Pothoidium_lobbianum</i>	0	0	0	0	1	0	0
<i>Pothos_chinensis</i>	0	0	1	0	1	0	1
<i>Pothos_scandens</i>	0	0	1	1	1	0	1
<i>Pothos_tener</i>	0	0	0	0	0	1	0
<i>Rhaphidophora_africana</i>	0	0	0	1	0	0	0
<i>Rhaphidophora_sp</i>	0	0	0	0	1	0	0
<i>Rhaphidophora_angustata</i>	0	0	0	0	1	0	0
<i>Rhaphidophora_beccarii</i>	0	0	0	0	1	0	0
<i>Rhaphidophora_crassicaulis</i>	0	0	1	0	1	0	0
<i>Rhaphidophora_decursiva</i>	0	0	1	0	1	0	1
<i>Rhaphidophora_guamensis</i>	0	0	0	0	0	1	0
<i>Rhaphidophora_hongkongensis</i>	0	0	1	0	1	0	0

<i>Rhaphidophora_hookeri</i>	0	0	1	0	1	0	0
<i>Rhaphidophora_lobbii</i>	0	0	0	0	1	0	0
<i>Rhaphidophora_luchunensis</i>	0	0	1	0	0	0	0
<i>Rhaphidophora_maingayi</i>	0	0	0	0	1	0	0
<i>Rhaphidophora_megaphylla</i>	0	0	1	0	1	0	0
<i>Rhaphidophora_montana</i>	0	0	0	0	1	0	0
<i>Rhaphidophora_peepla</i>	0	0	1	0	1	0	1
<i>Rhaphidophora_schlechteri</i>	0	0	0	0	1	1	0
<i>Rhaphidophora_spathacea</i>	0	0	0	0	0	1	0
<i>Rhaphidophora_spuria</i>	0	0	0	0	0	1	0
<i>Rhaphidophora_sylvestris</i>	0	0	0	0	1	1	0
<i>Rhaphidophora_tetrasperma</i>	0	0	0	0	1	0	0
<i>Rhodospatha_sp1</i>	0	0	0	0	1	1	0
<i>Rhodospatha_moritziana</i>	0	0	0	0	1	0	0
<i>Rhodospatha_sp2</i>	0	0	0	0	0	1	0
<i>Rhodospatha_sp3</i>	0	1	0	0	0	0	0
<i>Rhodospatha_sp4</i>	0	1	0	0	0	0	0
<i>Rhodospatha_latifolia</i>	0	1	0	0	0	0	0
<i>Rhodospatha_yenosa</i>	0	1	0	0	0	0	0
<i>Rhodospatha_wendlandii</i>	0	1	0	0	0	0	0
<i>Rhaphidophora_korthalsii</i>	0	1	0	0	0	0	0
<i>Rhaphidophora_latevaginata</i>	0	1	0	0	0	0	0
<i>Rhaphidophora_neoguineensis</i>	0	1	0	0	0	0	0
<i>Scindapsus_hederaceus</i>	0	0	0	0	1	0	0
<i>Scindapsus_rupestris</i>	0	0	0	0	1	0	0
<i>Spathiphyllum_cannifolium</i>	0	1	0	0	0	0	0
<i>Spathiphyllum_cochlearispathum</i>	0	1	0	0	0	0	0
<i>Spathiphyllum_minor</i>	0	1	0	0	0	0	0
<i>Spathiphyllum_solomonense</i>	0	0	0	0	0	1	0
<i>Spathiphyllum_wallisii</i>	0	1	0	0	0	0	0
<i>Spathiphyllum_commutatum</i>	0	0	0	0	1	1	0
<i>Spathiphyllum_patulinervum</i>	0	1	0	0	0	0	0
<i>Spathiphyllum_pygmaeum</i>	0	1	0	0	0	0	0
<i>Spirodela_polyrhiza</i>	1	1	1	1	1	1	1
<i>Stenospermatum_sp1</i>	0	1	0	0	0	0	0
<i>Stenospermatum_sp2</i>	0	1	0	0	0	0	0
<i>Stenospermatum_angustifolium</i>	0	1	0	0	0	0	0
<i>Stenospermatum_sp3</i>	0	1	0	0	0	0	0
<i>Stenospermatum_andreanum</i>	0	1	0	0	0	0	0
<i>Stenospermatum_ellipticum</i>	0	1	0	0	0	0	0
<i>Stenospermatum_sp4</i>	0	1	0	0	0	0	0
<i>Stenospermatum_escobariae</i>	0	1	0	0	0	0	0
<i>Stenospermatum_multiovulatum</i>	0	1	0	0	0	0	0

<i>Stylochaeton_bogneri</i>	0	0	0	1	0	0	0
<i>Urospatha_sagittifolia</i>	0	1	0	0	0	0	0
<i>Wolffiella_lingulata</i>	1	1	1	1	1	1	1
<i>Colocasia_esculenta</i>	0	0	0	0	1	0	1

TABLE S2. Dispersal multipliers matrices used in the time period analyses with BioGeoBEARS. Areas are coded as: Nearctic (A), Tropical America (BC; C-B-Central America, C South America in the eight area analysis), Palearctic (D), Africa (E), South East Asia (F), Australasia (G), India (H). The matrices correspond to the following time periods from top to bottom: 10, 50 90, and 150 Ma.

BC	G	F	E	D	A	H
1	0.001	0.001	0.001	0.001	1	0.001
0.001	1	1	0.001	0.1	0.001	0.1
0.001	1	1	0.1	1	0.1	1
0.001	0.001	0.1	1	0.5	0.001	0.1
0.001	0.1	1	0.5	1	0.5	1
1	0.001	0.1	0.001	0.5	1	0.001
0.001	0.1	1	0.1	1	0.001	1

BC	G	F	E	D	A	H
1	0.1	0.001	0.001	0.5	0.5	0.001
0.1	1	0.001	0.001	0.001	0.001	0.001
0.001	0.001	1	0.001	1	0.01	1
0.001	0.001	0.001	1	0.5	0.001	0.001
0.5	0.001	1	0.5	1	0.5	0.5
0.5	0.001	0.01	0.001	0.5	1	0.001
0.001	0.001	1	0.001	0.5	0.001	1

BC	G	F	E	D	A	H
1	0.1	0.001	0.5	0.001	0.5	0.1
0.1	1	0.001	0.01	0.001	0.001	0.1
0.001	0.001	1	0.001	1	0.001	0.001
0.5	0.01	0.001	1	0.5	0.001	0.5
0.001	0.001	1	0.5	1	0.5	0.001
0.5	0.001	0.001	0.001	0.5	1	0.001
0.1	0.1	0.001	0.5	0.001	0.001	1

BC	G	F	E	D	A	H
1	0.5	0.001	1	0.01	0.5	0.5
0.5	1	0.001	0.5	0.001	0.001	1
0.001	0.001	1	0.001	1	0.001	0.001

1	0.5	0.001	1	0.5	0.01	1
0.01	0.001	1	0.5	1	0.5	0.001
0.5	0.001	0.001	0.01	0.5	1	0.001
0.5	1	0.001	1	0.001	0.001	1

Chapter 3

Using genome skimming to resolve the phylogenetic relationships of the Neotropical genus *Monstera* (Araceae)

Alejandro Zuluaga¹, Monica Carlsen², Claudia Henríquez³, Elizabeth Kellogg⁴ & Ken Cameron¹

¹ Department of Botany, University of Wisconsin-Madison

² Missouri Botanical Garden, St. Louis

³ Department of Biology, Washington University, St. Louis

⁴ Donald Danforth Plant Science Center, St. Louis

ABSTRACT

The reconstruction of phylogenetic hypotheses for tropical aroids has been restricted to the use of traditional PCR amplification and Sanger sequencing of few individual genes. We used Next-Generation Sequencing (NGS) data from a genome skimming approach to recover plastid, mitochondrial, and nuclear ribosomal DNA for 22 samples comprising 18 species in the Neotropical genus *Monstera* and two other close related genera. We recovered complete plastomes for 15 samples, and plastid coding regions for the remaining four samples; 15 mitochondrial coding regions; and the 18S, 5.8S and 26S nrRNA. Plastomes in *Monstera* are collinear with no differences in arrangement or number of genes. Genetic variation among *Monstera* species plastid DNA was low, with only 1% of the complete plastome providing parsimony informative characters for phylogenetic analysis. However, phylogenetic reconstructions showed resolution and high support for the majority of the nodes. Previous infrageneric classifications of *Monstera* are not supported suggesting high levels of homoplasy in vegetative and reproductive structures used for classification. Our phylogenetic hypothesis represents a robust framework for the future studies systematics and evolution in the genus.

Keywords

Araceae, genome skimming, *Monstera*, next-generation sequencing, plastomes

INTRODUCTION

The reconstruction of molecular phylogenetic hypotheses for non-model taxa has been restricted primarily to the use of traditional PCR amplification and Sanger sequencing of individual genes (Cronn et al., 2012; Godden et al., 2012). This approach is both expensive and time consuming, and has been limited to a few plastid markers, nuclear ribosomal loci, or rarely low copy nuclear markers, depending on the availability of genomic tools for the group of interest. Additionally, for many groups, phylogenetic hypotheses of relationships based on few markers lack resolution and support due to the presence of relatively few variable and informative sites (Davis et al. 2005; Whitfield & Lockhart, 2007; Davis & Anderson, 2010). Other limitations for non-model taxa regard the difficulty of obtaining adequate tissue samples and extracting high-quality DNA. Historical herbarium specimens have been useful in some cases, but they often yield degraded DNA, and are usually only applicable for short segments of high copy number DNA regions (Varma et al., 2007), such as nuclear ribosomal and plastid markers.

High throughput DNA sequencing or ‘Next Generation Sequencing’ (NGS) approaches are rapidly improving our ability to produce phylogenetic hypotheses for non-model organisms (Atherton et al. 2010; Egan et al., 2012; Malé et al. 2014; Ripma et al., 2014). Most of the approaches developed so far involve some method of genome reduction by physical means (e.g., Restriction site associated DNA markers), or by targeting selected parts of the genome (e.g., transcriptomes [Wen et al., 2013; Yang et al., 2015], plastomes [e.g., Stull et al., 2013], or pre-determined single copy nuclear loci [e.g., Mandel et al., 2014; Weitemier et al., 2014; Zeng et al., 2014]). Reduction

techniques capturing large numbers of nuclear loci are ideal, but require some basic genomic knowledge (Cronn et al., 2012) or at least initial investment to achieve this goal. Reduction approaches using Restriction site associated DNA markers (RAD) are less expensive, viable for non-model organisms, and particularly useful for population genetics studies (e.g. Hohenlohe et al., 2010; Hohenlohe et al., 2011), but have been used with success for phylogenetic reconstruction at the species level as well (e.g., Eaton & Ree, 2013; Escudero et al., 2014).

On the other hand, high throughput DNA sequencing of total genomic DNA (gDNA) without genome reduction during library preparation, and with shallow depth of coverage (genome skimming method [Straub et al., 2012]), allows for assembly of the high-copy fraction of gDNA. For plants, this method has been successful in recovering the complete plastomes (cpDNA), the nuclear ribosomal cistron (nrDNA), and several regions of the mitochondria (mtDNA) (Ripma et al., 2014). The genome skimming method has been used to produce family-level phylogenies (Malé et al., 2014), species level phylogenies (Parks et al., 2009; Straub et al., 2012), and infra-species phylogenies (Whittall et al., 2010; Kane et al., 2012). Moreover, while directly recovering complete nuclear DNA sequences suitable for phylogenetic reconstruction is not an immediate outcome of these methods, the data from genome skimming studies may help to provide information to later identify low-copy nuclear genes (LCNG) of known identity that could be used for targeted sequencing studies in the absence of genomic tools for the group of interest (Straub et al., 2012; Ripma et al., 2014).

In this study we used genome skimming as a tool to generate a phylogenetic hypothesis in a poorly known genus of the tropical plant family Araceae. Molecular studies at the species and even the genus level in the family Araceae are particularly well known for their lack of resolution (e.g. *Anthurium*, [Carlsen et al., 2013]; *Amorphophallus* [Grob et al., 2004]), and only a handful of commonly used molecular markers have been traditionally employed in these studies (e.g., Grob et al., 2004; Cabrera et al., 2008). Only recently have genomic tools been developed that may help to identify new markers or to design targeted gene sequencing studies across the family. Henriquez et al. (2014), using genome skimming, sequenced gDNA for 32 taxa across Araceae and assembled complete plastid genomes and some mitochondrial coding regions. Additionally, several aroid transcriptomes have been sequenced (e.g., *Arum*, *Anthurium*, *Amorphophallus*), but although raw data are available in public databases the assembled sequences for the transcripts are not. Moreover, a disproportionate amount of work has been focused on the subfamily Lemnoideae (duckweeds) relative to other aroids, due to its bioenergy application potential. Mardanov et al. (2008) and Wang & Messing (2011) sequenced complete plastomes for four genera in the Lemnoideae; and Wang et al. (2014) sequenced and assembled the complete genome of *Spirodela polyrhiza*. These tools combined with the increasing popularity of NGS techniques need to be explored in order to improve our understanding of the relationships and evolution in Araceae.

The genus *Monstera* is a Neotropical member of the aroid subfamily Monsterodieae, and has evolved within a clade of otherwise Old World genera (Tam et

al., 2004; Cusimano et al., 2011; Zuluaga et al., **CHAPTER 2**). Members of the Monsteroideae are mostly hemiepiphytic plants inhabiting wet and moist forests across the tropics (Madison, 1977). The genus *Monstera* ranges from southern Mexico to Brazil and Bolivia, from rain forest at sea level up to cloud forest at 2300 m (Madison et al., 1977). Among other characteristics, *Monstera* is recognized by its remarkable heteroblastic development and the adult leaves with fenestrations (resulting in the common name of ‘Swiss cheese plant’). *Monstera* has a convoluted taxonomic history, and currently comprises 42 described species –significantly more than 23 species that were known at the time of the last taxonomic revision by Madison (1977)– and about 15 new taxa are awaiting description (Croat, 2005; Zuluaga et al., unpublished). No previous molecular studies have focused on the genus itself; only Tam et al. (2004) sequenced the plastid *trnL-trnF* spacer for four species of *Monstera* in a phylogenetic analysis of the subfamily Monsteroideae. Additionally, AFLP analysis was used to investigate taxonomic subtleties in the *Monstera adansonii* species complex in Brazil (Andrade et al., 2007). Our phylogenetic studies (Zuluaga et al., **CHAPTER 2**) have shown a striking lack of molecular variation within the genus despite the fact that it seems to be a relatively old lineage (crown age 20 Ma) with many morphologically unique taxa.

METHODS

Taxonomic sampling. 19 samples of the genus *Monstera* were used for the genome skimming approach, including species from the four sections proposed by Madison (1977). One *Epipremnum* and one *Rhodospatha* species were included to be used as outgroups. One sample of *Spathiphyllum* and one of *Monstera* from Henriquez et

al. (2014) were also included in the phylogenetic analysis. Leaf tissue samples were obtained from fieldwork carried out in Panama, Colombia and Ecuador, and the Araceae Greenhouse at the Missouri Botanical Garden in St. Louis, Missouri. The list of species, vouchers, and GENBANK accession numbers are shown in (**TABLE S1**).

Library preparation and sequencing. 10-30 mg of silica dried leaf tissue was used to extract total genomic DNA using Qiagen DNeasy Minikit (Qiagen, Germantown, Maryland, USA) following the manufacturer's protocol. CTAB method using the protocol from Doyle & Doyle (1987) was used for some recalcitrant samples. The tissue was ground using TissueLyser II (Qiagen, Courtaboeuf, France), and final DNA was eluted using 80 μ l of Qiagen elution buffer. 2 μ l of each extraction were run in agarose gel electrophoresis to confirm the presence of high molecular weight DNA. DNA concentrations were calculated using a fluorescence method via a Qubit fluorometer and the dsDNA HS Assay (Life Technologies, Invitrogen division, Darmstadt, Germany). 20 ng of DNA for each sample were sheared using a Covaris device (Covaris Inc.) to average size of 300-400 bp at the University of Missouri DNA Core Facility. Library preparation was performed in the lab of E. Kellog at the University of Missouri, Columbia, using the NEBNext DNA Library Prep kit for Illumina (New England Biolabs, MA). Briefly, purified fragments were A-tailed and ligated to sequencing indexed adapters. Fragments with an insert size of approximately 200 bp were size selected using AMPure® XP Beads (Beckman Coulter, Inc. #A63881). Libraries were enriched using five cycles of PCR and clean using AMPure® XP Beads (Beckman Coulter, Inc. #A63881) before library quantification and validation. Real time-PCR (qPCR) was

performed on each sample to quantify the relative amount of plastid fragments in each sample. Twenty-two samples were pooled together using the concentrations obtained after qPCR quantification. Sequencing was performed using 100 bp single-end read in one lane of Illumina HiSeq 2000 (Illumina, Inc., San Diego, California) at the University of Missouri DNA Core Facility.

Demultiplexing and quality trimming. Raw fastq reads were demultiplexed, barcodes and adapters were removed, and an initial quality trim was performed using Cutadapt v. 1.7 (Martin, 2011). Subsequent quality trimming was performed using the `fastq_quality_trimmer` function from the FASTX-Toolkit v. 0.0.13 (Gordon, 2010)

Plastid, mitochondria and ribosomal DNA assembly. Demultiplexed reads for each sample were assembled using a hybrid approach combining de novo and reference guided assemblies, similar to the one used by Henriquez et al. (2014) or the one proposed by Wysocki et al. (2014). *De novo* assembly was performed using the software SPAdes v 3.1.1 (Bankevich et al., 2012) and Velvet (Zerbino & Birney, 2008). We used k-mer lengths of 19 for velvet and 19, 21, 33, 55 for SPAdes. SPAdes and Velvet analyses were performed via server clusters at the Center for High Throughput Sequencing (CHTC) at the University of Wisconsin- Madison. An initial reference guide assembly was performed using YASRA v. 2.33 (Ratan, 2009) using an *Anthurium* sample as reference (Carlsen et al., unpublished data), and 85% similarity threshold. One initial *Monstera* plastome (*M. adansonii*-Brazil) was completely assembled and used as the reference for

the rest of the samples in the YASRA runs. This reference only included one copy of the plastome inverted repeat to avoid double matches during the assembly.

Post assembly analysis was performed almost entirely using Geneious 8.0 (Biomatters, <http://www.geneious.com>, Kearse et al., 2012) in a similar way as described by Ripma et al. (2014). Final contigs from both the reference guided and the *de novo* assemblies were transferred to Geneious. Contigs with average coverage <5 were discarded. Two subsequent assemblies runs using the Geneious assembler were performed to create longer contigs. These “final” contigs were mapped back to the initial *Monstera* plastome using the Geneious map reference tool to obtain a plastome draft. Each plastome draft was preliminary annotated based on the *Spirodela polyrhiza* plastid genome using the transfer annotations tool in Geneious. We checked each draft manually looking for regions of no coverage or regions of incongruence among different contigs (i.e., regions with long repeats). To solve these incongruences the 100 bp ends of the contigs were deleted and subsequently extended manually by looking back into the original reads using command line. Extended contigs were assembled back in Geneious. When no unambiguous assembly was found Ns were added to the consensus sequence to represent unknowns. We confirmed start and stop codons of each coding region, and evaluated reading frames using protein translation in Geneious. Finally, a consensus sequence was generated and this complete plastome sequence was exported into a FASTA for final annotation using DOGMA (Wyman et al., 2004).

We mapped *Monstera adansonii*-Brazil contigs from the *de novo* assembly to the ribosomal DNA of *Wolffia* (Lemnoideae) using the Geneious Map to reference tool, and extracted contigs with high similarity and high coverage. These contigs were used as reference for a new YASRA run, to finally assemble one complete ribosomal DNA repeat region containing 26S, 18S, 5.8S and portions of the internal and external transcribed spacers. This complete nrDNA was used as a reference for the rest of the samples. Similarly, *M. adansonii*-Brazil contigs from the *de novo* assembly were mapped to the *Spirodela polyrhiza* mitochondrial genome. The contigs that mapped protein-coding regions were extracted, and used as our reference for all the subsequent mitochondrial assemblies using YASRA and *de novo* contigs. Only 15 protein coding mitochondrial genes were identified, and we included up to 500 bases flanking each side of the regions for our alignments.

Alignment and phylogenetic reconstruction. For the plastomes, individual protein-coding regions, rRNA and tRNA, introns and spacers were aligned independently using one initial MAFFT v7 (Katoh et al., 2002, Katoh & Standley, 2013) run, and two subsequent MUSCLE v3.8.31 (Edgar 2004) runs. We concatenated: 1) all regions, and also 2) only coding regions (exons, rRNA, tRNA), and 3) only non-coding regions (introns and spacers). nrDNA and mtDNA consensus sequences were aligned the same way, and then concatenated. Phylogenetic reconstructions were performed under maximum parsimony (MP) and maximum likelihood (ML). MP analyses were conducted in PAUP* v. d104 (Swofford 2003), with 1000 bootstrap replicates to assess the relative degree of support for individual nodes. ML analyses were conducted in RAxML v. 8.0.9

(Stamatakis 2014), as implemented in the CIPRES Science Gateway (Miller et al. 2010). For each case during the ML analyses we used the GTRCAT model of substitution and performed 20 independent tree searches, and 1000 subsequent bootstraps replicates. To calculate average coverage of each genomic region we averaged the coverage of all the contigs that were mapped to the reference plastome. In that sense, measurements of the mitochondrial and nrDNA are relative only to the regions recovered.

RESULTS

Sequencing and assembly of plastid, mitochondrial and nrDNA. We found large variation in the amount of raw and filtered reads for each sample. *Monstera adansonii* -Colombia produced 1,240,022 raw reads (1,154,270 filtered reads) whereas *Rhodospatha* produce 35,887,362 raw reads (3,3288,989 filtered reads) (**TABLE 2**). The average coverage for the plastid varied between 7.5-157.6. Due to low coverage many contigs from *M. tuberculata*, *M. dubia*, *M. acacoyaguensis*, and *M. adansonii*-Colombia were discarded, and not all plastid regions were recovered, especially the spacer regions (**TABLE 3**). In general, assemblies and coverage of plastid DNA were more homogeneous through the length of the genome, whereas assemblies of mitochondrial DNA were sporadic and variable among species. Using *S. polyrhiza* as a reference for the mitochondrial DNA we were able to assemble up to 15 protein-coding regions including some of the flanking regions (between 50 and 500 bp). Assemblies for nrDNA were more problematic. Although the 18S, 5.8S and 26S nrRNA matched few contigs with high coverage producing consensus sequences without or with very ambiguities, assemblies from external and internal transcribed regions (ETS, ITS) matched hundreds of contigs

producing consensus sequences with many ambiguities that made it nearly impossible to obtain reliable sequences. Despite discarding all contigs with low coverage (<5), sequences for the ETS and ITS remained ambiguous, thus only 18S, 5.8S and 26S were considered for the alignments. The average coverage for the 15 mitochondrial coding regions varied between 6.3 and 76.4, and for the nrDNA between 15.6 and 396.5.

Structure of the plastid genome in *Monstera*. For the species where complete or almost complete plastomes were assembled the sequences were collinear with no differences in arrangement or number of genes. We used data for the species with better coverage in the genus (*M. adansonii*-Brazil) for the purposes of describing plastome structure. The plastid of *M. adansonii*-Brazil is 163,244 nucleotides long including the two inverted repeat regions (IRs) with a GC content of 38 %; whereas the range in *Monstera* species was approximately 163,000-166,000 nucleotides long. The IRs contain 25,721 nucleotides. The boundary between the large single copy region (LSC) and the inverted repeat (IRa) was between *rpl2* and *rpl19*. The boundary between the IRa and the small single copy region (SSC) was between *tRNA-Asn (GUU)* and *ndhF* in all species. The SSC-IRb boundary was between *ycf1* and *tRNA-Asn (GUU)*. Finally, the IRb-LSC boundary was between *tRNA-His (GUG)* and *rpl2* in all species (**Figure 1**).

We identified a total of 77 protein-coding regions (plus seven duplicated in the IR), four rRNA regions, and 30 tRNA regions. Additionally, we identified three regions with high similarity to the protein coding genes *cemA* (*ycf10*), *infA*, and *ycf68*, but we did not find a reading frame for protein translation of them.

DNA alignments and phylogenetic reconstruction. Complete plastid alignments, excluding the IRb were 129,162 bp long with 2.3% representing variable sites and just 1% of the total being parsimony informative characters (PIC) (**TABLE 3**). Alignment of the protein coding regions comprised 79,062 bp (2.2% variable sites), whereas the alignment of the non-coding regions was 50,100 bp (2.4% variable sites). Fifteen spacer regions were excluded from the alignment since they did not contain any variable sites. In general, missing data varied between 1.2 and 9.7 %, with only one sample, *M. tuberculata* falling outside this range and reaching 36 % (**TABLE 2**). Final alignment of mitochondrial regions and nrDNA comprised 21768 bp (0.7 % variable sites), and 5386 bp (2.2 % variable sites) respectively. In general, alignments of mitochondrial coding regions and nrDNA lacked variability within *Monstera*, even when flanking regions were considered for the mitochondria. Phylogenetic reconstructions from these regions do not have any resolution besides supporting the monophyly of the genus *Monstera* so they are not presented or discussed. ML and MP analyses of the plastid matrices produced identical topologies (**Figures 2 & 3**). Analyses from non-coding vs. coding regions produced different topologies (**Figures 2 & 3**), particularly regarding the position of *M. deliciosa* and *M. florescanoana*. However, nodes with incongruence have low bootstrap support (< 75). Three species in section *Marcgraviopsis* from Central America are grouped in a clade, with the monotypic section *Echinospadix* (*M. tuberculata*) nested within (clade 1, **figure 3**). This clade is sister to the rest of *Monstera* in the non-coding and complete plastid datasets, but not in the coding regions analysis. *Monstera deliciosa* and/or *M. florescanoana*, also from Central America appear sister to

the remaining *Monstera* (excluding the clade 1). The remaining species form four well-supported clades (BS ≥ 95 but relationships among them remaining unknown, **figure 3**).

DISCUSSION

Sequencing and assembly of plastid, mitochondrial, and nrDNA. Large differences in read coverage restricted us from recovering complete plastomes for all species (**TABLE 2**); however, we showed that even very shallow sequencing allowed us to recover large amounts of information (e.g., ca. 104,000 nucleotides with 36% missing data for the plastid genome in the sample with less success). The three plastid coding regions that seem not to be functional (*cemA*, *infA*, and *ycf68*) were also reported as problematic by Henriquez et al., (2015) across Araceae. Additionally these genes have been reported as losses or pseudogenes in many other groups of angiosperms (Millen et al., 2001). A study of the plastid evolution within the family Araceae including plastomes from this study is underway (Carlsen et al. Unpublished data) so we will not discuss differences among the three clades in the subfamily Monsteroideae. Briefly, *Spathiphyllum*, *Rhodospatha* and *Monstera* plastid sequences are collinear, without any major rearrangements, the main differences are in the length of spacer regions. For the mitochondrial genome we recovered few coding regions, which had low variation among *Monstera* species (0.06 % PICs). The low amount of coding regions recovered in the mtDNA may be due to size and structure of the *Monstera* mitochondrial genome in comparison with the reference used (*S. polyrhiza*). *Spirodela* represents the most compact monocot mitochondrial genome known so far, with 228,493 bases, containing 35 proteins-coding regions, 3 ribosomal RNAs and 19 transfer RNAs (Wang et al., 2012).

Studies in the chloroplast of Lemnoideae have shown high rates of mutation in comparison with other aroids, so it is probable the same is true for the mitochondria. The closest available mitochondrial genome to Araceae is for the palm *Phoenix dactylifera*, which comprises 715,001 bp and contains 38 proteins, 30 tRNAs, and 3 ribosomal RNAs. Using other mitochondrial genomes as references may yield more and longer regions for our data; but due to the lack of variability found so far we did not pursue this. Similarly, nrDNA lacked variation for *Monstera* (0.4% PICs). ITS and ETS regions have been used once in Araceae for phylogenetic reconstruction (Gauthier et al., 2008), but other studies have failed to use this DNA region and the presence of pseudogenes has been suggested (Carlsen, 2011). Many biological processes like allopolyploidy or incomplete lineage sorting can cause polymorphism and/or pseudogenic sequences in the ITS region (Alvarez & Wendel, 2003; Won & Renner, 2005); however it is difficult to establish which may be causing the problems with the assembly of the nrDNA in *Monstera*.

The amount of variable sites and PICs observed in our plastid DNA dataset (2.3 % and 1 %, **TABLE 3**) are similar to what has been found in other species level studies using complete plastid genomes, (e.g., 3% PIC in *Ipomoea* [Eserman et al., 2014]; 0.48% in *Oreocarya* [Ripma et al., 2014]). However, if we considered only the *Monstera* species (ingroup) the amount of PIC is only 0.5%, again similar to Ripma et al., (2014) for *Oreocarya* (0.32%). In the absence of genomic tools the discovery of molecular markers of phylogenetic utility in tropical Aroids has been evasive. Few chloroplast markers, and even less from nuclear DNA have been used that are suitable for species level phylogenies. In this context, NGS approaches like Genome Skimming offer an alternative

to produce large amounts of data from high copy regions of the DNA in a very efficient and inexpensive manner. Moreover, in a preliminary approach we performed blast searches (E-value < 1e-5) of *Monstera* contigs from the de novo assembly against the 959 single copy genes that are shared in the genomes of *Arabidopsis thaliana*, *Populus trichocarpa*, *Vitis vinifera* and *Oryza sativa* (Duarte et al., 2010). We recovered ca. 100 matches to exons of these genes. The utility of these data to develop nuclear markers either for individual sequencing or for NGS target approaches seems to have great potential.

Phylogenetic reconstruction and implications for the systematics of *Monstera*.

Despite the low variation in the plastid DNA, we were able to recover highly supported phylogenetic reconstructions for the genus *Monstera*. Some incongruence was found among coding and non-coding regions of the plastid genome, but they corresponded to nodes of low support (BS<75), particularly regarding the positions of *M. deliciosa* and *M. florescanoana*. Several studies have shown that the mechanisms of molecular evolution present in non-coding regions of the chloroplast may produce homoplasious patterns that could obscure the actual relationships among taxa (Kelchner, 2000). Despite we tried to exclude regions with ambiguous alignment, inversions, and regions with repeats, other contradictory sequences may persist.

The sectional classification proposed by Madison (1977) was not supported completely in any of our analyses. Phylogenetic relationships of *Monstera* do not reflect the morphological patterns described by Madison (1977), and morphological characters

that are important for taxonomic purposes appear to be homoplasious in *Monstera*. The one that is especially interesting is the presence of shingled leaves, which has been the defining characteristic for the section *Marcgraviopsis* (Madison, 1977). Species from this group appeared in three unrelated high supported clades, implying that the extreme heteroblastic development (i.e., development of a stolon-like seedling, follow by shingled-leave juveniles, and large adult leaves) has evolved more than once in *Monstera*. Although an unexpected result, this type of development has evolved several times in Araceae, especially in the family Monsteroideae (e.g., *Rhaphidophora* and *Scindapsus*), but also in other groups (e.g., *Philodendron*). However the homology of the process resulting in the shingles leaves has not been tested. A group of species from Central America, all having shingled leaves, but very different adult leaf morphologies form a clade (clade 1, **Figure 3**). The neotenic species *M. tuberculata* placed by Madison in a monotypic section (*Echinospadix*) is nested in this clade. Neoteny has evolved several times in Monsteroideae as well, one in *Scindapsus* and at least one other time in *Rhaphidophora* (Boyce & Bogner, 2000). Other species of *Monstera* (e.g., *M. pittieri*, *M. minima*) not included in this study exhibit reduction of the adult leaves, but they are not considered neotenic since adult leaf shapes are slightly different. Evolution of leaf shape in *Monstera* (and in general in Monsteroideae) is an interesting topic for future research. The other defining character of *M. tuberculata*, the echinate spadix has been found in other unrelated species in these analyses (*M. lentii*).

The position of *M. deliciosa* (section *Tornelia*) and *M. florescanoana* varies in the different reconstructions. They appear as sister to the rest of the genus (excluding clade 1,

Figure 3), and they are very distinctive species morphologically. *Monstera florescanoana* is endemic from Veracruz in Mexico, the northern most extreme range for the genus (Croat et al., 2010). This species was assigned to the section *Monstera*, and is the only species in the genus with glaucous leaves in the lower surface. *Monstera deliciosa* also has a unique set of morphological and anatomical characters. Among these unique features we can mention: 1) wide flowers (0.7-1 cm, whereas in all the other species flower width range between 0.2-6 mm); 2) flowers with trichosclereids distributed across the entire flower, whereas in the rest of the species studied the trichosclereids are concentrated in rings around the ovary; and 3) vascular bundles in the flower are distributed in two concentric rings vs. only one a ring of vascular bundles in other species (Zuluaga et al. Unpublished data). Clade 2 comprises four species from Central and South America without extreme heteroblastic development (section *Monstera*); all having lobulated leaves. Clade 3 comprises five species of Central America with a wide morphological variation, including one species with shingled leaves (*M. acuminata*). Clade 4 is composed by two Central American species with the distinctive characters of section *Marcgraviopsis*. These two species are robust plants (with leaves reaching 1.5 m long.) that flower high in the canopy (10-20 m). Finally, clade 5 comprises three species with a wide distribution range. *Monstera adansonii* is the most wide spread species in the genus, from Costa Rica to Brazil and Bolivia, reaching the Caribbean islands.

The phylogenetic reconstructions from plastid genomes in the genus *Monstera* suggest a striking pattern of parallel evolution of morphological characters in vegetative

and reproductive structures. However, the lack of nuclear markers with informative characters prevented us of investigating other hypotheses like the presence of hybridization, and subsequent chloroplast capture. DNA sequences from three nuclear regions (Chalcone synthase, Phytochrome C, and at103 putative protein) showed low or no variation for the species in the genus (Unpublished data). Nauheimer et al. (2012) found incongruences between chloroplast and nuclear markers in the genus *Alocasia*, suggesting hybridization and chloroplast capture in this group. Hybridization has been hypothesized to be common in *Monstera* (Madison, 1977); however there is not evidence supporting this hypothesis. The use of approaches that allow us to recover information from several variable nuclear regions is critical to advance in the understanding of the evolution of the genus *Monstera*, and data generated from this study may help to achieve this goal.

REFERENCES

- Álvarez, I., Wendel, J.F., 2003. Ribosomal ITS sequences and plant phylogenetic inference. *Mol. Phylogenet. Evol.* 29, 417–434. doi:10.1016/S1055-7903(03)00208-2.
- Andrade, I.M., Mayo, S.J., Berg, C. van den, Fay, M.F., Chester, M., Lexer, C., Kirkup, D., 2007. A preliminary study of genetic variation in populations of *Monstera adansonii* var. *klotzschiana* (Araceae) from North-East Brazil, estimated with AFLP molecular markers. *Ann. Bot.* 100, 1143–1154. doi:10.1093/aob/mcm200.
- Atherton, R.A., McComish, B.J., Shepherd, L.D., Berry, L.A., Albert, N.W., Lockhart, P.J., 2010. Whole genome sequencing of enriched chloroplast DNA using the Illumina GAI platform. *Plant methods* 6, 22. doi:10.1186/1746-4811-6-22.
- Bankevich, A., Nurk, S., Antipov, D., Gurevich, A.A., Dvorkin, M., Kulikov, A.S., Lesin, V.M., Nikolenko, S.I., Pham, S., Prjibelski, A.D., Pyshkin, A.V., Sirotkin, A.V., Vyahhi, N., Tesler, G., Alekseyev, M.A., Pevzner, P.A., 2012. SPAdes: A new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19, 455–477. doi:10.1089/cmb.2012.0021.
- Boyce, P.C., Bogner, J., 2000. An account of neotenic species of *Raphidophora* Hassk. (Araceae-Monsteroideae-Monstereae) in New Guinea and Australia. *Gard. Bull. Singapore* 52, 89–100.
- Cabrera, L.I., Salazar, G.A., Chase, M.W., Mayo, S.J., Bogner, J., Dávila, P., 2008. Phylogenetic relationships of aroids and duckweeds (Araceae) inferred from coding and noncoding plastid DNA. *Am. J. Bot.* 95, 1153–1165.

- Carlsen, M.M., 2011. Understanding the origin and rapid diversification of the genus *Anthurium* Schott (Araceae), integrating molecular phylogenetics, morphology and fossils. PhD thesis, University of Missouri–St. Louis, USA.
<https://apps.umsi.edu/webapps/weboffice/ETD/query.cfm?id=r6587>
- Carlsen, M.M., Croat, T.B., 2013. A molecular phylogeny of the species-rich Neotropical genus *Anthurium* (Araceae) based on combined chloroplast and nuclear DNA. *Syst. Bot.* 38, 576–588. doi: <http://dx.doi.org/10.1600/036364413X670287>.
- Croat, T.B., Acebey, A., 2005. New species of Araceae from Bolivia and the tropical Andes. *Novon*, 80–103.
- Croat, T.B., Krömer, T., Acebey, A., 2010. *Monstera florescanoana* (Araceae), a new species from central Veracruz, Mexico. *Revista Mexicana de Biodiversidad* 81: 225–228.
- Cronn, R., Knaus, B.J., Liston, A., Maughan, P.J., Parks, M., Syring, J.V., Udall, J., 2012. Targeted enrichment strategies for next-generation plant biology. *Am. J. Bot.* 99, 291–311.
- Cusimano, N., Bogner, J., Mayo, S.J., Boyce, P.C., Wong, S.Y., Hesse, M., Hetterscheid, W.L.A., Keating, R.C., French, J.C., 2011. Relationships within the Araceae: comparison of morphological patterns with molecular phylogenies. *Am. J. of Bot.* 98, 654–668.
- Davis, C.C., Anderson, W.R., 2010. A complete generic phylogeny of Malpighiaceae inferred from nucleotide sequence data and morphology. *Am. J. Bot.* 97, 2031–2048. doi: 10.3732/ajb.1000146.

- Davis, C.C., Webb, C.O., Wurdack, K.J., Jaramillo, C.A., Donoghue, M.J., 2005. Explosive radiation of Malpighiales supports a mid-cretaceous origin of modern tropical rain forests. *Am Nat* 165, E36–65. doi:10.1086/428296.
- Doyle, J.J., Doyle, J.L., 1987. A rapid DNA isolation procedure for small quantities of fresh leaf tissue. *Phytochem. Bull.* 19, 11–15.
- Duarte, J.M., Wall, P.K., Edger, P.P., Landherr, L.L., Ma, H., Pires, J.C., Leebens-Mack, J., dePamphilis, C.W., 2010. Identification of shared single copy nuclear genes in *Arabidopsis*, *Populus*, *Vitis* and *Oryza* and their phylogenetic utility across various taxonomic levels. *BMC Evol. Biol.* 10, 61. doi:10.1186/1471-2148-10-61.
- Eaton, D.A.R., Ree, R.H., 2013. Inferring phylogeny and introgression using RADseq data: An example from flowering plants (*Pedicularis*: Orobanchaceae). *Syst Biol* 62, 689–706. doi:10.1093/sysbio/syt032.
- Edgar, R.C., 2004. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res.* 32, 1792–1797.
- Egan, J.B., Shi, C.X., Tembe, W., Christoforides, A., Kurdoglu, A., Sinari, S., Middha, S., Asmann, Y., Schmidt, J., Braggio, E., Keats, J.J., Fonseca, R., Bergsagel, P.L., Craig, D.W., Carpten, J.D., Stewart, A.K., 2012. Whole-genome sequencing of multiple myeloma from diagnosis to plasma cell leukemia reveals genomic initiating events, evolution, and clonal tides. *Blood*, 120, 1060–1066. doi: <http://dx.doi.org/10.1182/blood-2012-01-405977>.
- Escudero, M., Eaton, D.A.R., Hahn, M., Hipp, A.L., 2014. Genotyping-by-sequencing as a tool to infer phylogeny and ancestral hybridization: A case study in *Carex*

- (Cyperaceae). *Mol. Phylogenet. Evol.* 79, 359–367.
doi:10.1016/j.ympev.2014.06.026.
- Eserman, L.A., Tiley, G.P., Jarret, R.L., Leebens-Mack, J.H., Miller, R.E., 2014. Phylogenetics and diversification of morning glories (tribe Ipomoeae, Convolvulaceae) based on whole plastome sequences. *Am. J. Bot.* 101, 92–103.
doi:10.3732/ajb.1300207.
- Gauthier, M.P.L., Barabe, D., Bruneau, A., 2008. Molecular phylogeny of the genus *Philodendron* (Araceae): delimitation and infrageneric classification. *Bot. J. Linean Soc.* 156, 13–27. doi: 10.1111/j.1095-8339.2007.00733.x.
- Godden, G. T., Jordon-Thaden, I.E., Chamala, S., Cowl, A.A., García, N., Germain-Aubrey, C. C., García, N., Germain-Aubrey, C.C., Heaneyab, J.M., Latvisab, M., Qiaf, Z., Gitzendanner, M.A., 2012. Making next-generation sequencing work for you: approaches and practical considerations for marker development and phylogenetics. *Plant Ecol. Divers.* 5, 427–450.
- Gordon, A. 2010 *FASTX-toolkit*. Available at http://hannonlab.cshl.edu/fastx_toolkit/index.html. Accessed Nov, 2014.
- Grob, G.B.J., Gravendeel, B., Eurlings, M.C.M., 2004. Potential phylogenetic utility of the nuclear FLORICAULA/LEAFY second intron: comparison with three chloroplast DNA regions in *Amorphophallus* (Araceae). *Mol. Phylogenet. Evol.* 30, 13–23. doi:10.1016/S1055-7903(03)00183-0.
- Henriquez, C.L., Arias, T., Pires, J.C., Croat, T.B., Schaal, B.A., 2014. Phylogenomics of the plant family Araceae. *Mol. Phylogenet. Evol.* 75, 91–102.
doi:10.1016/j.ympev.2014.02.017.

- Hohenlohe, P.A., Amish, S.J., Catchen, J.M., Allendorf, F.W., Luikart, G., 2011. Next-generation RAD sequencing identifies thousands of SNPs for assessing hybridization between rainbow and westslope cutthroat trout. *Mol Ecol Resour* 11, 117–122. doi:10.1111/j.1755-0998.2010.02967.x.
- Hohenlohe, P.A., Bassham, S., Etter, P.D., Stiffler, N., Johnson, E.A., Cresko, W.A., 2010. Population genomics of parallel adaptation in threespine stickleback using sequenced RAD tags. *PLoS Genet* 6, e1000862. doi:10.1371/journal.pgen.1000862.
- Kane, N.C., Sveinsson, S., Dempewolf, H., Yang, J.Y., Zhang, D., Engels, J.M., Cronk, Q., 2012. Ultra-barcoding in cacao (*Theobroma* spp.; Malvaceae) using whole chloroplast genomes and nuclear ribosomal DNA. *Am. J. Bot.* 99, 320–329.
- Katoh, K., Misawa, K., Kuma, K., Miyata, T., 2002. MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. *Nucl. Acids. Res.* 30, 3059–3066. doi:10.1093/nar/gkf436.
- Katoh, K., Standley, D.M., 2013. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Mol. Biol. Evol.* 30, 772–780. doi:10.1093/molbev/mst010.
- Kearse, M., Moir, R., Wilson, A., Stones-Havas, S., Cheung, M., Sturrock, S., Buxton, S., Cooper, A., Markowitz, S., Duran, C., Thierer, T., Ashton, B., Meintjes, P., Drummond, A., 2012. Geneious Basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* 28, 1647–1649. doi:10.1093/bioinformatics/bts199.

- Kelchner, S.A., 2000. The evolution of non-coding chloroplast DNA and its application in Plant Systematics. *Ann. Mo. Bot. Gard.* 87, 482–498. doi:10.2307/2666142.
- Madison, M., 1977. A revision of *Monstera* (Araceae). *Contrib. Gray Herb. Harvard Univ.* 207, 3–100.
- Magallón S, Castillo A., 2009. Angiosperm diversification through time. *Am. J. Bot.* 96: 349–365.
- Malé, P.J.G., Bardon, L., Besnard, G., Coissac, E., Delsuc, F., Engel, J., Lhuillier, E., Scotti-Saintagne, C., Tinaut, A., Chave, J., 2014. Genome skimming by shotgun sequencing helps resolve the phylogeny of a pantropical tree family. *Mol Ecol Resour* 14, 966–975. doi: 10.1111/1755-0998.12246.
- Mandel, J.R., Dikow, R.B., Funk, V.A., Masalia, R.R., Staton, S.E., Kozik, A., Micheltore, R.W., Rieseberg, L.H., Burke, J.M., 2014. A target enrichment method for gathering phylogenetic information from hundreds of loci: An example from the Compositae¹. *Appl Plant Sci* 2. doi:10.3732/apps.1300085
- Mardanov, A.V., Ravin, N.V., Kuznetsov, B.B., Samigullin, T.H., Antonov, A.S., Kolganova, T.V., Skyabin, K.G., 2008. Complete sequence of the duckweed (*Lemna minor*) chloroplast genome: Structural organization and phylogenetic relationships to other Angiosperms. *J. Mol. Evol.* 66, 555–564. doi:10.1007/s00239-008-9091-7.
- Martin, M., 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet.journal* 17, pp. 10–12. doi:10.14806/ej.17.1.200.
- Millen, R.S., Olmstead, R.G., Adams, K.L., Palmer, J.D., Lao, N.T., Heggie, L., Kavanagh, T.A., Hibberd, J.M., Gray, J.C., Morden, C.W., Calie, P.J., Jermini,

- L.S., Wolfe, K.H., 2001. Many parallel losses of *infA* from chloroplast DNA during angiosperm evolution with multiple independent transfers to the nucleus. *Plant Cell* 13, 645–658.
- Miller, M.A., Pfeiffer, W., Schwartz, T., 2010. Creating the CIPRES Science Gateway for inference of large phylogenetic trees. In: *Proceedings of the Gateway Computing Environments Workshop (GCE 2010)*. New Orleans, LA: Institute of Electrical and Electronics Engineers (IEEE). Pp. 1-8.
- Nauheimer, L., Metzler, D., Renner, S.S., 2012. Global history of the ancient monocot family Araceae inferred with models accounting for past continental positions and previous ranges based on fossils. *New Phytol.* 195, 938–950.
- Parks, M., Cronn, R., Liston, A. 2009. Increasing phylogenetic resolution at low taxonomic levels using massively parallel sequencing of chloroplast genomes. *BMC biology* 7, 84. doi:10.1186/1741-7007-7-84.
- Ratan, A., 2009. Assembly algorithms for next-generation sequence data. *PhD thesis*. The Pennsylvania State University, Computer Science and Engineering.
- Ripma, L.A., Simpson, M.G., Hasenstab-Lehman, K., 2014. Geneious! Simplified genome skimming methods for phylogenetic systematic studies: A case study in *Oreocarya* (Boraginaceae). *Appl Plant Sci* 2. doi:10.3732/apps.1400062.
- Stamatakis, A., 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* 30, 1312–1313. doi:10.1093/bioinformatics/btu033.

- Straub, S.C., Parks, M., Weitemier, K., Fishbein, M., Cronn, R.C., Liston, A. 2012. Navigating the tip of the genomic iceberg: Next-generation sequencing for plant systematics. *Am. J. Bot.* 99, 349–364. doi: 10.3732/ajb.1100335.
- Stull, G.W., Moore, M.J., Mandala, V.S., Douglas, N.A., Kates, H.R., Qi, X., Brockington, S.F., Soltis, P.S., Soltis, D.E., Gitzendanner, M.A. 2013. A targeted enrichment strategy for massively parallel sequencing of Angiosperm plastid genomes. *Appl Plant Sci* 1, apps.1200497. doi: 10.3732/apps.1200497.
- Swofford, D.L., 2003. PAUP*. Phylogenetic Analysis Using Parsimony (*and Other Methods). Version 4. Sinauer Associates, Sunderland, Massachusetts, USA.
- Tam, S.M., Boyce, P.C., Upson, T.M., Barabé, D., Bruneau, A., Forest, F., Parker, J.S., 2004. Intergeneric and infrafamilial phylogeny of subfamily Monsteroideae (Araceae) revealed by chloroplast trnL-F sequences. *Am. J. Bot.* 91, 490–498.
- Varma, A., Padh, H., Shrivastava, N., 2007. Plant genomic DNA isolation: an art or a science. *Biotechnol. J.* 2: 386–392.
- Wang, W., Haberer, G., Gundlach, H., Gläßer, C., Nussbaumer, T., Luo, M.C., Lomsadze, A., Borodovsky, M., Kerstetter, R.A., Shanklin, J., Byrant, D.W., Mockler, T.C., Appenroth, K.J., Grimwood, J., Jenkins, J., Chow, J., Choi, C., Adam, C., Cao, X.-H., Fuchs, J., Schubert, I., Rokhsar, D., Schmutz, J., Michael, T.P., Mayer, K.F.X., Messing, J., 2014. The *Spirodela polyrhiza* genome reveals insights into its neotenuous reduction fast growth and aquatic lifestyle. *Nat Commun* 5. doi:10.1038/ncomms4311.

- Wang, W., Messing, J., 2011. High-throughput sequencing of three Lemnoideae (Duckweeds) Chloroplast Genomes from Total DNA. PLoS One 6, e24670. doi:10.1371/journal.pone.0024670.
- Wang, W., Wu, Y., Messing, J., 2012. The Mitochondrial Genome of an Aquatic Plant, *Spirodela polyrhiza*. PLoS One 7, e46747. doi:10.1371/journal.pone.0046747.
- Weitemier, K., Straub, S.C.K., Cronn, R.C., Fishbein, M., Schmickl, R., McDonnell, A., Liston, A., 2014. Hyb-Seq: Combining target enrichment and genome skimming for plant phylogenomics1. Appl Plant Sci 2. doi:10.3732/apps.1400042.
- Wen, J., Xiong, Z., Nie, Z.L., Mao, L., Zhu, Y., Kan, X.Z., Ickert-Bond, S.M., Gerrath, J., Zimmer, E.A., Fang, X.D. 2013. Transcriptome sequences resolve deep relationships of the grape family. PloS One 8, e74394. doi: 10.1371/journal.pone.0074394.
- Whitfield, J.B., Lockhart, P.J., 2007. Deciphering ancient rapid radiations. Trends Ecol. Evol. 22, 258–265. doi:10.1016/j.tree.2007.01.012.
- Whittall, J.B., Syring, J., Parks, M., Buenrostro, J., Dick, C., Liston, A., Cronn, R., 2010. Finding a (pine) needle in a haystack: chloroplast genome sequence divergence in rare and widespread pines. Mol. Ecol. 19(s1), 100–114. doi: 10.1111/j.1365-294X.2009.04474.x.
- Won, H. & Renner, S.S., 2005. The internal transcribed spacer of nuclear ribosomal DNA in the gymnosperm *Gnetum*. Mol. Phylogenet. Evol. 36, 581–597. doi:10.1016/j.ympev.2005.03.011.

- Wyman, S.K., Jansen, R.K., Boore, J.L., 2004. Automatic annotation of organellar genomes with DOGMA. *Bioinformatics* 20, 3252–3255. doi:10.1093/bioinformatics/bth352.
- Wysocki, W.P., Clark, L.G., Kelchner, S.A., Burke, S.V., Pires, J.C., Edger, P.P., Mayfield, D., Triplett, J.K., Columbus, J.T., Ingram, A., Duvall, M.R., 2014. A multi-step comparison of short-read full plastome sequence assembly methods in grasses. *Taxon* 63, 899–910. doi: <http://dx.doi.org/10.12705/634.5>.
- Yang, Y., Moore, M.J., Brockington, S.F., Soltis, D.E., Wong, G.K.-S., Carpenter, E.J., Zhang, Y., Chen, L., Yan, Z., Xie, Y., Sage, R.F., Covshoff, S., Hibberd, J.M., Nelson, M.N., Smith, S.A., 2015. Dissecting molecular evolution in the highly diverse plant clade Caryophyllales using transcriptome sequencing. *Mol Biol Evol* msv081. doi:10.1093/molbev/msv081.
- Zeng, L., Zhang, Q., Sun, R., Kong, H., Zhang, N., Ma, H., 2014. Resolution of deep angiosperm phylogeny using conserved nuclear genes and estimates of early divergence times. *Nat Commun* 5. doi:10.1038/ncomms5956.
- Zerbino, D.R., Birney, E., 2008. Velvet: Algorithms for de novo short read assembly using de Bruijn graphs. *Genome Res.* 18, 821–829. doi:10.1101/gr.074492.107.

TABLES AND FIGURES

Table 1. Species names and voucher information for the samples use in this study. **Note:**

Some voucher numbers still need to be assigned.

Species	Voucher	GenBank accession numbers
<i>Monstera lechleriana</i>	Zuluaga A. s.n. (COL)	
<i>Monstera egregia</i>	Zuluaga A. s.n. (WIS)	
<i>Monstera sp nov anomala</i>	Zuluaga A. 888 (WIS, PMA, MO)	
<i>Monstera oreophila</i>	Zuluaga A. 886 (WIS, PMA, MO)	
<i>Monstera punctulata</i>	Zuluaga A. 887 (WIS, PMA, MO)	
<i>Monstera pinnatiparita</i>	Zuluaga A. 892 (WIS)	
<i>Monstera acuminata</i>	Zuluaga A. s.n. (WIS)	
<i>Monstera florescanoana</i>	Zuluaga A. s.n. (WIS)	
<i>Monstera tuberculata</i>	Zuluaga A. s.n (WIS)	
<i>Monstera dubia</i>	Zuluaga A. s.n. (WIS)	
<i>Monstera acacoyaguensis</i>	Zuluaga A. s.n. (WIS)	
<i>Epipremnum pinnatum</i>	Zuluaga A. s.n (WIS)	
<i>Monstera adansonii-Colombia</i>	Zuluaga A. 400 (COL)	
<i>Monstera adansonii-Brazil</i>	Monsteiro 465	
<i>Monstera gracilis</i>	Zuluaga A. 421 (COL)	
<i>Rhodospatha venosa</i>	Croat, T. 104784 (MO)	
<i>Monstera deliciosa</i>	Zuluaga A. 902 (WIS, PMA, MO)	
<i>Monstera filamentosa</i>	Zuluaga A. 893 (WIS, PMA)	
<i>Monstera sp nov integrigolia</i>	Zuluaga A. 916 (WIS, PMA, MO)	
<i>Monstera lentii</i>	Zuluaga A. 911 (WIS, PMA, MO)	
<i>Monstera spruceana</i>	Zuluaga A. 899 (WIS)	

Table 2. Information from the Illumina sequencing of genomic DNA for 21 samples used in this study. Number of raw reads and number of reads after quality filter are shown.

mtDNA and nrDNA were calculated only for the length of the regions we recovered.

Species	Raw reads	Filter reads	Plastid mean coverage	mtDNA mean coverage	nrDNA mean coverage
<i>Monstera lechleriana</i>	7974607	7303098	94.9	26.3	126.4
<i>Monstera egregia</i>	5059727	4775714	86.0	35.8	99.7
<i>Monstera sp nov anomala</i>	2522940	2340270	20.6	11.5	50.7
<i>Monstera oreophila</i>	3750804	3540264	15.6	11.9	250
<i>Monstera punctulata</i>	2838671	2679330	15.0	8.9	44.6
<i>Monstera pinnatiparita</i>	2078598	1903568	15.1	7	34.5
<i>Monstera acuminata</i>	3320896	3041257	21.0	16.4	48.7
<i>Monstera florescanoana</i>	2847041	2650159	16.0	13.9	55.6
<i>Monstera tuberculata</i>	1845985	1690542	7.5	7.1	18.5
<i>Monstera dubia</i>	1829675	1697200	9.9	6.3	43.5
<i>Monstera acacoyaguensis</i>	1804758	1674087	11.0	6.8	15.6
<i>Epipremnum pinnatum</i>	3824160	3502143	27.3	13.8	66.7
<i>Monstera adansonii</i>	1240022	1154270	8.9	7.5	25.4
<i>Monstera adansonii</i>	23834154	21827177	103.5	45.7	271
<i>Monstera gracilis</i>	15780683	14689397	37.5	15.6	156.4
<i>Rhodospatha venosa</i>	35887362	33288989	157.6	76.4	396.5
<i>Monstera deliciosa</i>	14556713	13502755	46.3	25	234.6
<i>Monstera filamentosa</i>	11103196	10168241	78.3	28.1	125.7
<i>Monstera coloradense</i>	11198974	10570352	49.7	26.4	176.5
<i>Monstera lentii</i>	10060891	9332447	110.7	36.5	101.6
<i>Monstera spruceana</i>	9816048	9137236	75.6	33.2	102.4


Table 3. Alignment information for the different genomic regions; coding and non- coding regions of the plastid, complete plastid, mtDNA, and nr DNA. The length of the alignment, the percentage of missing data, the variable sites and the Parsimony informative characters (PICs) are shown.

Alignment	Length	Missing Data	Variable sites	PICs	PICs(<i>Monstera</i>)
Non-coding regions	50100	8.8%	1236 (2.4%)	543(1.1%)	291(0.6%)
Coding regions	79062	3.2%	1719 (2.2%)	699 (0.88%)	363(0.45%)
Plastids (excluding IR)	129162	5.9%	2955 (2.3%)	1242(1%)	666(0.5%)
nrDNA	5386	1.8 %	123(2.2%)	34(0.63%)	21(0.4%)
mtDNA	21768	29.9 %	153(0.7%)	22(0.1%)	15(0.06%)

Figure 1. Complete annotated chloroplast of *Monstera adansonii*. Genes (including introns) are coded in green, coding regions in yellow, rRNA in red, tRNA in pink, and inverted repeats in light blue.

Figure 2. Phylograms from Maximum Likelihood (RAxML) analysis of coding (left) and non-coding regions (right) from the plastid genome of species in the genus *Monstera*. Bootstrap support values are shown. Grey bars indicate species with positions supported in both trees, black bars indicate species with incongruences.

Figure 3. Cladogram depicting the topology from Maximum Likelihood (ML) analysis of complete plastomes of species in the genus *Monstera* using RAxML. Bootstrap support values less than 100 are shown above the branches for ML, and below for Maximum Parsimony (MP); unsupported node in MP are indicated by n.s. Inset tree indicates branch lengths.



Chapter 4

Taxonomic studies in the Neotropical genus *Monstera* (Araceae)

Part 1. Two new species of *Monstera* (Araceae) with entire leaves from Panama and Costa Rica

Alejandro Zuluaga¹, Thomas B. Croat² & Kenneth Cameron¹

¹ Department of Botany, University of Wisconsin-Madison

² Missouri Botanical Garden, S.t. Louis

ABSTRACT

The genus *Monstera* (Araceae) is a poorly known genus in terms of its systematics and ecology, and historically its taxonomy has been problematic due to the large number of existing names, incorrect use of them, and existence of few complete herbarium specimens. The last taxonomic revision of the genus included 22 species, but now is very much out of date, and there are more species than this recognized today in Panama or Costa Rica alone. Today, among ca. 130 names, only 42 species are recognized, and as many as 10-15 new taxa remain undescribed. A product of field explorations and study of herbarium specimens, in the first part of this chapter we described two new species of *Monstera* from Panama and Costa Rica. In the second part, we developed a web based interactive tool for the identification of the species in the genus using the software Lucid. For this purpose we created a geographic database with ca. 4200 herbarium specimens; and an identification database including 108 ecological, morphological and anatomical characters. So far we have included 42 recognized species plus the two new species described in this study.

Keywords

Araceae, Costa Rica, interactive key, Lucid, *Monstera*, new species, Panama,

INTRODUCTION

Species of the genus *Monstera* (Araceae) are an abundant and important hemiepiphytic component of montane and lowland tropical rainforests (Madison, 1977). The genus is exclusively Neotropical and has two principal centers of diversity: Costa Rica–Panamá and the Colombian Andes. *Monstera* is a poorly known genus in terms of its ecology, and historically its taxonomy has been problematic due to the large number of existing names, incorrect use of them, and prevalence of incomplete herbarium specimens (Madison, 1977). The last taxonomic revision of the genus (Madison, 1977) included 22 species, but now is very much out of data, and there are more species than this recognized today in Panama or Costa Rica alone (Andrade & Mayo, 1998; Grayum, 2003; Croat *et al.*, 2005). Today, among ca. 130 names, only 42 species are recognized, and as many as 10–15 new taxa remain undescribed. Based on morphological characters Madison (1977) proposed four sections within *Monstera*, changing considerably previous classifications (e.g., Engler, 1908). The species of sections *Marcgraviopsis*, *Tornelia*, and *Echinospadix* (ca. 10 species collectively) are easily recognized based on their morphology, growth habits, and habitat preferences. Section *Marcgraviopsis* is characterized by a heteroblastic development with three clear stages: a stolon-like seedling lacking leaves; an early juvenile phase with adpressed leaves (shingled) similar to those of the eudicot genus *Marcgarvia*; and pre-adult and/or adult plants with exserted leaves that are otherwise different in shape from shingled leaves (Madison, 1977; Ray, 1987; Oberbauer & Noudali, 1998). *Monstera* section *Tornelia* (= *M. deliciosa* Liebm. [1849:19]) and section *Echinospadix* (= *M. tuberculata* Lundell [1939:78]) are both monotypic, and recognized by its large pistils and seeds, and for the neotenic adult leaves with a ligule

respectively (Madison, 1977). In contrast, delimitation of species in the fourth section, *Monstera*, is notoriously difficult because of the presence of numerous intermediates connecting them (Madison, 1977). Recent molecular phylogenetic studies (Zuluaga et al., unpublished data) do not support the monophyly of the *Monstera* sections, showing that morphological used in classifications are homoplastic. However, most of the species in section *Marcgraviopsis* do form a clade (albeit with monotypic sections *Echinospadix* and *Tornelia* nested within).

MATERIALS AND METHODS

Field explorations in Panama and study of herbarium collections, especially at the Missouri Botanical Garden (MO), have revealed several undescribed taxa in the genus *Monstera*. Many of these new taxa are being described for the treatment of the forthcoming Flora Mesoamericana (Croat personal communication), and here we describe two new species of *Monstera* bearing entire leaves and lacking fenestrations.

TAXONOMIC TREATMENT

Monstera anomala Zuluaga & Croat *sp. nov.* (Figs 1 & 2)

Diagnosis:—*Monstera anomala* is easy to distinguish from other species in the genus with shingled leaves because of its flowers with a long and constricted stylar region, 4–6 mm long, which is narrower than the ovary. It is most similar to *M. acuminata* K.Koch (1855[App]: 4), but adult plants are more robust, with entire leaves lacking fenestrations, and with secondary and tertiary veins inconspicuous on the upper surface.

Type: —PANAMA. Veraguas: Distrito Santa Fe, corregimiento El Pantano, Parque Nacional Santa Fe, alto Los Gonzales to alto El Viro, 800–1000 m. 8°33.162'N–8°33.625'N, 81°4.586'–81°4.767'W, 16 Jan 2013, *A. Zuluaga, A. Doucette, E. Brantner & E. González 888* (Holotype: WIS!; Isotypes: PMA!, MO!).

Seedling stolon-like, on the ground of the forest, green, 1–3 mm diameter. *Juvenile plants* completely adpressed to the host (i.e., with shingled leaves); petioles less than 6 mm long; blades oblate to circular, 0.5–10 cm in diameter, size increasing gradually with the height on the host, asymmetrical, apex with a short acumen, suddenly switching to the adult shape. *Pre-adult and adult* stems adpressed to the host, circular in cross section, 1.5–2.8 cm across, internodes 3–7 cm long. Pre-adult and adult leaves exerted from the host, alternate, most of the leaves distichous, except for the ones close to the apex of the plant; petioles 20–50 cm long, 0.4–1.2 cm diameter, circular in cross section, departing 45–90 degrees from the host, petiole sheathed for the almost entire petiole (excluding the geniculum); sheath marcescent, the base persistent as fibers; geniculum 1.8–3.8 cm long, 0.4–1.2 diameter; blades entire, very rarely with fenestrations or lobate, perpendicular to the host to slightly pendent, elliptical to oblong, 0.9–1.5 times longer than the petiole, 27–60 cm long, 9–20 cm width, 2.1–4.2 times longer than wider, symmetrical to asymmetrical, apex turning towards one side, discolor, upper surface bright green and glossy, lower surface paler looking whitish, and mate, base obtuse, apex acuminate; main vein flat in the upper surface, raised in the lower surface, primary lateral veins sunken in the upper surface, raised in the lower surface, 10–15 per side, secondary lateral veins

inconspicuous on the upper surface, visible in the lower surface; tertiary veins reticulate, inconspicuous. Plants flowering (6–)10–20 m above the ground, inflorescence axillary, solitary; peduncle 6–14 cm long, 0.6–1.4 cm diameter, circular in cross section, green; spathe white in anthesis, 10–13 cm long, 1–3 cm longer than the spadix; spadix terete in cross section, slightly narrower at the apex, 12–22 cm long, 2.7–5 cm diameter, 1.7–2.8 times longer than the peduncle. Flowers white, 15–22 in the primary spiral, 17–25 in the secondary spiral, 8–18 mm long, cross section at the ovary circular, 3–4 mm diameter, style 3–5 mm long, cross section at the style narrower than the ovary, 2–4 mm diameter, apex of the flower as wide as the ovary, hexagonal viewed from the top, trichosclereids abundant in the stylar region and reaching the ovary; stigma linear, 2–3 mm long; stamens four, filaments laminar, 3–4 mm long; anthers 2–3 mm long; ovary 3–4 mm long, 2-locular, two ovules per locule. Fruiting spadix first green, yellow at maturity, 15–27 cm long, 5–8 cm diameter; stylar cap falling off the fruits exposing the white aril. Seeds one per fruit, green–blue, 4–6 mm long, 3–4 cm wide, 1–2 mm thick; embryo green.

Etymology:— The epithet *anomala* (meaning anomalous or strange), was chosen because of the species unique combination of foliar and flower characters not common in other *Monstera* species.

Distribution and habitat:— *Monstera anomala* ranges from North Central Panama to South Central Costa Rica, at elevations between 100–1200 m. This species grows high (10–20 m) into the canopy of humid premontane and montane forests (**Figure 3**).

Observations:— *Monstera anomala* has all the general characteristics of the species in the section *Marcgraviopsis* sensu Madison (1977). From among this group of species *M. anomala* it is most similar to *M. acuminata* K. Koch, a widespread species from Mexico to Panama. However, *M. anomala* differs from *M. acuminata* in having more coriaceous leaves, larger length/width leaf ratio, inconspicuous secondary lateral veins on the lower surface (vs. conspicuous), and flowers with a long and narrow style (vs. style as wide as the ovary) (**Table 1**). Some specimens of *M. spruceana* (Schott) Engler (1878: 115) with leaves lacking fenestrations could be confused with *M. anomala*; however, *M. spruceana* is a more robust plant with deeply lobate leaves. Grayum and Croat first recognized several specimens of *M. anomala* as a different taxon, and assigned the name *M. skutchii* to several collections from Costa Rica. However, all these specimens were later identified as *M. spruceana* for the Araceae treatment of the Costa Rican Flora (Grayum, 2003). Morphologically *M. spruceana* is a species distinctive from *M. anomala*, and its distribution throughout Central America is not clear.

Additional specimens examined:—**COSTA RICA. Cartago:** Turrialba, along camino Raíz de Hule, south east of Platanillo (Tsipirí), 9°49'12"N, 83°24'W, 1200–1400 m, 1 Jul 1976, *T.B. Croat* 36788 (MO); Turrialba, along Quebrada Platanillo near confluence of Quebrada Sipirí, Platanillo de Chirripó, 9°48'36"N, 83°24'W, 1135 m, 2 Mar 1990, *M.H. Grayum* 9727 (MO). **Heredia:** Sarapiquí, finca El Bejuco, S base of Cerros Sardinal, Chilamate de Sarapiquí. 10°27'N, 84°3'36"W, 70–100 m, 2 Jun 1985, *M.H. Grayum* 5327 (MO); Sarapiquí, finca La Selva, Puerto Viejo de Sarapiquí, ridge near end of Pasos

Perdidos, 10°25'N, 84°2'W, 120 m, 3 Jun 1985, *M.H. Grayum 5356* (DUKE, MO).

Limón: Talamanca, Cantón de Talamanca, Alto Urén, subiendo a Cerro Chun, siguiendo un antiguo camino de Quebrada Chaho a Alto Lari, 9°24'N, 83°19'48"W, 800 m, 24 Jul 1989, *G. Herrera–Chacón 3330* (MO); along road between Puerto Viejo de Talamanca and Manzanillo, stretch from Quebrado Ernesto to Manzanillo, 9°37'48"–09°38'N, 82°40'48"–82°42'W, 5 m, 3 Nov 1984, *M.H. Grayum 4352* (AAU, CR, MO); hills 3.5 airline km S of Islas Buena Vista in the Río Colorado, 16 airline km SW of Barra del Colorado, premontane wet forest on low hills, 10°39'N, 83°40'12"W, 10–120 m, 15 Sep 1986, *G. Davidse 31269* (MO); Refugio Barra del Colorado, forests and pastures between Río Chirripocito and Río Sardina (Sardinal on Chirripó Atlántico quadrangle), 10°37'48"N, 83°45'W, 12 m, 21 Apr 1990, *M.H. Grayum 9809* (MO). **Puntarenas:** Osa, hills north of Palmar Norte, along trail to Jalisco, 8°58'48"N, 83°25'12"W, 50–700 m, 21 May 1976, *T.B. Croat 35187* (MO); along highway to Golfito from Panamerican highway at Río Claro, 2.5 mi SE of Golfito, 8°36'N, 83°3'36"W, 60 m, 14 Sep 1987, *T.B. Croat 67585* (HNMN, MO); along road between Palmar Norte and Panamerican border, 3 km N of turn-off to Rincón, 8°48'39"N, 83°16'18"W, 110 m, 10 Sep 1996, *T.B. Croat 79200* (INB, MO); along road between Rincón de Osa and Rancho Quemado, ca. 10 km W of main Rincón–Pto. Jimenez Road, 8°41'N, 83°32'W, 150–260 m, 3 Mar 1985, *T.B. Croat 59755* (CAS, GB, HNMN, MO); Rincón de Osa, along ridge between quebrada Aparicio and quebrada Aguabuena, 8°42'N, 83°30'36"W, 200–400 m, 7 Oct 1984, *M.H. Grayum 4015* (MO, NY). **San José:** Perez Zeledon, about 1 mile beyond divide between San Isidro del General and coastal town of Dominical, 9°16'12"N, 83°51'36"W, 900 m, 22 May 1976, *T.B. Croat 35304* (MO). **PANAMA. Chiriquí:** San Bartolo Limite, 18 km

W of Puerto Armuelles, rainforest, 8°17'24"N, 82°58'36"W, 450 m, 24 Feb 1973, *P. Busey 604* (MO). **Colón:** Teck Cominco Petaquilla mining concession, forested slopes along ridge road, 8°49'28"N, 80°39'29"W, 110 m, 10 Mar 2008, *G.D. Mcpherson 20496* (MO, PMA). **Panamá:** Capira, cloud forest on Cerro Campana above Su Lin Motel, 8°43'N, 79°54'W, 209 m, 25 May 1971, *T.B. Croat 14767* (MO); middle slopes of Cerro Campana, ca. 1 mi from Interamerican Highway, 8°43'9"N, 79°53'25"W, 150 m, 15 Jun 1976, *T.B. Croat 35989* (MO); Panamá, vicinity of Cerro Jefe, 4.6 km beyond peak on road to Altos de Pacora, 26.3 km from the Inter-American Highway, 9°14'20"N, 79°20'25"W, 600 m, 12 Jun 1976, *T.B. Croat 35914* (MO); Cerro Campana, 6.1 miles above Pan-American highway, 3.2 miles beyond park entrance and Guarda Bosque Station, 8°41'N, 79°56'W, 800 m, 23 Mar 1993, *T.B. Croat 74767* (CM, MO). **Veraguas:** Santa Fé, along 1st branch of Río Santa María, roadside and forest a short way S of river, road from Santa Fé, 8°31'26"N, 81°7'46"W, 600 m, 26 Oct 1975, *J.T. Witherspoon 8903* (MO); NW of Santa Fé, ca. 1km from Escuela Agrícola Alto de Piedra, Pacific slope, 8°31'22"N, 81°7'35"W, 800 m, 21 Dec 1974, *S.A. Mori 4024* (MO); vicinity of Santa Fé, along dirt road from Santa Fé to Río San Luis, past Escuela Circolo Alto de Piedra, at Río Segundo Brazo (2nd stream below school on Atlantic Coast), 8°33'N, 81°8'W, 480 m, 28 Jun 1987, *T.B. Croat 66902* (MEXU, MO).

***Monstera integrifolia* Zuluaga & Croat *sp. nov.* (Figures 4 & 5).**

Diagnosis:—*Monstera integrifolia* has erect and small entire leaf blades (16–31 cm long) lacking fenestrations. It is most similar in appearance to *M. xanthospataha* Madison

(1977: 65–67) but differs in the shorter peduncles (9–14 vs. 13–25 cm), longer spadix (8–10 vs. 5–7 cm), and raised stylar region (vs. flat).

Type: PANAMA. Chiriqui: Distrito Gualaca, corregimiento Hornito, Reserva Forestal Fortuna, trails near to research center Jorge L. Arauz. 1200–1500 m elevation, 8°47'W, 82°12.8'N, 31 Jan 2013, *A. Zuluaga 916* (Holotype: WIS!, Isotypes: PMA!, MO!).

Seedling not seen. *Juvenile* plants with stems adpressed to the host; leaves exerted from the host in an angle less than 45 degrees, leaf blade shape the same as adults leaves but increasing gradually in size. *Pre-adult and adult* plants growing on vertical stems and lateral branches; stem depressed, D shape in cross section, flat side facing the host, 1–3 cm diameter, internodes 4–6 cm long. Leaves erect or almost erect, spreading from an angle less than 45 degrees on a vertical host, or 45–90 degrees when host is horizontal; petioles 5–10 cm long, 1–2 cm diameter, D shape in cross section, petiole sheathed almost its entire length, sheath marcescent, not persistent; geniculum 2–3 cm long; blades ovate to lanceolate, assymetrical, 16–31 cm long, 7–10 cm width, blade 1.2–1.6 times longer than petiole, coriaceous, dark green on both surfaces, paler in the lower surface, base obtuse, apex acuminate; main vein flat in the upper surface, raised in the lower surface, primary lateral veins sunken in the upper surface, raised in the lower surface, 4–8 per side, secondary lateral veins sunken on the upper surface, slightly raised in the lower surface; tertiary veins reticulate, inconspicuous. Plants flowering 3–5 m above the ground, inflorescence solitary; peduncle 9–14 cm long, 0.6–1.2 cm diameter, circular in cross section, green; spathe white in anthesis, 15–18 cm long, 8–9 cm width, 5–8 cm

longer than the spadix; spadix terete, slightly narrower at the apex, 8–10 cm long, 1.7–2 cm diameter, 0.6–0.9 times longer than the peduncle. Flowers white, 12–14 in the primary spiral, 14–17 in the secondary spiral, 4–5 mm long, cross section at the ovary circular, 3–4 mm diameter, apex of style narrower than the ovary, raised, 1–2 mm long; stigma capitated, 1–2–mm long; stamens four; filaments laminar, 3–4 mm long; anthers 2–3 mm long. Infructescence, fruits, and seeds not seen.

Etymology:—The specific epithet *integrifolia* refers to the entire leaves lacking fenestrations, a character not common in *Monstera*.

Distribution and habitat:—*Monstera integrifolia* is only known from four collections from North Panama and the Panama–Costa Rica border. It grows in wet and humid forests at 1000–2000 m above sea level (**Figure 6**).

Observations:—*Monstera integrifolia* is most similar to *M. xanthospatha*, a species endemic to the West and Central Cordilleras of Colombia at elevations between 1500–2300 m. Both species are small plants, flowering when they are less than 10 m high in the host, and inhabit montane cloud forests. *Monstera integrifolia* differs from *M. xanthospatha* in having shorter petioles and peduncles, a longer spadix, and flowers with raised stylar regions (vs. flat) (see **Table 2**.)

Additional specimens examined:—**COSTA RICA. Puntarenas:** Corredores, along road between San Vito de Java and Villa Neily, 8.8 miles north of Villa Neily,

8°42'28"N, 82°56'16"W, 1010 m, 14 Jun 1987, *T.B. Croat 66170* (MO). **PANAMA.**

Bocas del Toro: Cerro Colorado, 9.2 mi W of Chamé, along trail E of road which leads down to stream, 8°35'N, 81°50'W, 1450–1480 m, 6 Jul 1988, *T.B. Croat 69067* (MO, PMA); Changuinola, about 1 mile S of Changuinola on the Changuinola River, forest edge of banana grove, 9°24'06"N, 82°30'W, 19 Apr 1969, *R.L. Lazor 2658* (MO).

Chiriquí: Fortuna Dam site, forest area, 8°44'N, 82°15'W, 1400–1600 m, 15 Sep 1977, *J.P. Folsom 5598* (MO).

Acknowledgments

Thanks to Sarah Friedrich at **Botany Media Studio at University of Wisconsin**

Madison. This study was partly supported by Davis Research Grant and the Vilas Grant from the University of Wisconsin-Madison.

REFERENCES

- Andrade, I.M., Mayo, S.J., 1998. Dynamic Shoot Morphology in *Monstera adansonii* Schott var. *klotzschiana* (Schott) Madison (Araceae). *Kew Bulletin* 53, 399–417.
doi:10.2307/4114505
- Croat, T.B., Acebey, A., 2005. New Species of Araceae from Bolivia and the Tropical Andes. *Novon* 15, 80–103.
- Engler, A., Krause, K., 1908. Araceae–Monsteroideae. In: *Das Pflanzenreich. IV. 23B* (Heft 37), ed., Engler, A., Leipzig.
- Engler, A. 1878. Araceae.. In Martius, K.F.P. von, Eichler, A.W., Endlicher, I.L., Fenzl, E., Mary, B. Oldenburg, R. & Urban, I., (1878-1882). *Flora Brasiliensis, enumeratio plantarum in Brasili hactenus detectarum*. v.3, pt.2, pp 25–224.
Munich & Leipzig, R. Oldenbourg.
- Grayum, M. H. 2003. Araceae. In: *Manual de Plantas de Costa Rica*, B.E. Hammel, M.H. Grayum, C. Herrera & N. Zamora (eds.). *Monogr. Syst. Bot. Missouri Bot. Gard.* 92: 59–200.
- Koch, K. 1855. *Monstera acuminata*. *Index seminum in Horto Botanico Berolinensi*.
Appendix. Genera et speciarum novarum et minus cognitarum. Pp. 4. In
Botanischer Garten (Berlin, Germany).
- Liebmann, F.M. 1849. Om Mexicos Aroider. *Videnskabelige Meddelelser fra Dansk Naturhistorisk Forening i Kjøbenhavn* Nr1-2: 11–25.
- Lundell, C.L. 1939. Studies of Mexican and Central American Plants VII. *Lloydia* 2: 73–108
- Madison, M. 1977. A revision of *Monstera* (Araceae). *Contr. Gray Herb.* 207: 1–100.

- Oberbauer, S., Noudali, M., 1998. Potential carbon gain of shingle leaves in juveniles of the vine *Monstera tenuis* (Araceae) in Costa Rica. *Am. J. Bot.* 85, 850–850.
- Ray, T.S., 1987. Leaf Types in the Araceae. *American Journal of Botany* 74, 1359–1372.
doi:10.2307/2444314

TABLES AND FIGURES

Table 1. Morphological differences between *Monstera anomala* and *M. acuminata*

Species	Leaf	Blade space	Length/width		
	fenestrations	orientation	Secondary veins	leaf ratio	Style shape
<i>M. anomala</i>	absent	Perpendicular	Inconspicuous	1.7–2.8	Constricted,
		to slightly pendent	on the lower surface		narrower than ovary
<i>M. acuminata</i>	present or absent	Pendent	Visible on the lower surface	1.3–2.0	Straight, as wide as ovary

Table 2. Morphological differences between *Monstera integrifolia* and *M. xanthospatha*

Species	Leaf fenestrations	Peduncule length	Spadix length	Flower styler region
<i>M. integrifolia</i>	absent	9–14 cm	8–10 cm	Raised
<i>M. xanthospatha</i>	present or absent	13–25 cm	5–7 cm	Flat

Figure 1. Comparison between *Monstera anomala* sp. nov. (**A–C, E–F**), and *M. acuminata* (**D, G–I**). *Monstera anomala*: **A**, Adult plant showing leaves lacking fenestrations and infrutescence; **B–C**, mature infrutescence showing stylar caps falling off; **E**, Juvenile shingled-leaf plant; **F**, individual fruits showing narrow stylar region of the flower. *Monstera acuminata*: **D**, inflorescence; **G–I**, adult leaves of with and without fenestrations.




Figure 2. Illustration of *Monstera anomala* sp. nov.



Figure 3. Distribution map of *Monstera anomala*.




Figure 4. Comparison between *Monstera integrifolia* sp. nov. (**A–C**) and *M. xanthospatha* (**D–G**). *Monstera integrifolia*: **A**, adult leaf lacking fenestrations; **B**, detail of the periole showing the punctuated pattern; **C**, young spadix showing blunt flowers. *Monstera xanthospatha*: **D–E**, Adult leaves of with and without fenestrations; **F**, young spadix enclosed by the spathe; **G**, infrutescence.




Figure 5. Illustration of *Monstera integrifolia* sp. nov.





Figure 6. Distribution map of *Monstera integrifolia*.



Part 2. A preliminary interactive web tool for the identification of the species in the genus *Monstera* (Araceae)

Alejandro Zuluaga¹, Kenneth Cameron¹

¹ Department of Botany, University of Wisconsin-Madison

INTRODUCTION

New web-based tools are being used to facilitate the gathering, storage, and access of taxonomical, geographical, and ecological information about particular taxonomical groups or for complete inventories from particular regions (Brach et al., 2005; Brach & Song, 2006; Clark et al., 2009). Web-based tools used to document taxonomical groups have many advantages: 1) they can be constantly updated and modified, allowing the inclusion/deletion new taxa, or any information about morphology, geography or ecology; 2) there is not limitation to the amount visual material that can be added for a particular taxon; and 3) they generally free and easily access for anyone with an Internet connection. Many flexible and multifunctional tools have been developed to facilitate this task. The Scratchpad Virtual Research Environment (<http://scratchpads.eu/>) (Smith et al., 2009; Smith et al., 2011) is one of the most flexible and easy to use resources of this kind, with about ca. 680 web-based sites using it to date (<http://scratchpads.eu/>).

The family Araceae was used as a model group for the Creating a Taxonomic e-Science (CATE) project, which goal was to create an E-Taxonomy of the family, including a full taxonomic revision for all genera and species (Haigh et al., 2009). This information was later transferred to the Scratchpad environment, and it is now available at eMonocot Team Araceae (*CATE Araceae* Monocot Portal, <http://araceae.e-monocot.org/>). One of the tools available through the *CATE Araceae*-eMonocot portal is a series of interactive keys for identification of genera in Araceae, Araceae from Africa, *Anthurium* species, among others; all of them built using the software Lucid (www.lucidcentral.com). These

tools are from great utility not only for scientist but also for enthusiasts and growers. Interactive keys and other digital identification tools have become very popular in recent years and several options are available (e.g., Linnaeus II [www.eti.uva.nl/products/linnaeus.php]; Lucid Key Server [www.lucidcentral.com]; Xper2 [<http://lis-upmc.snv.jussieu.fr/lis/?q=en/resources/software/xper2>]. but see Dallwitz et al., and 2013; Farnsworth et al., 2013, for a list and comparison of interactive key tools). The Lucid software offers a series of tools to build interactive and dichotomous keys as well as fact sheets for taxa, and is one of the most used interactive key tools with more than 250 keys available from a variety of organisms and regions around the world (lucidcentral.org). One of the downsides of this software is that the most up to date version is not freely available to the public, although older versions are.

Traditional taxonomy for many groups of Araceae is complex, especially for those from the tropics, and current taxonomic revisions and dichotomous keys are very much out of date, and/or are insufficient for species identification. In some cases like for the genera *Anthurium* and *Philodendron* the main limitation to build efficient taxonomy databases and identification tools is the large number of species (ca. 1000 and 500 respectively) and its constant growing; in other cases like the genus *Monstera* (ca. 50 species) the limitation is the deficient collections, the chaotic taxonomic history, and the general poor knowledge about the natural history of the group. Our goal is to create a complete web-based tool for the genus *Monstera* that includes not only taxonomical information, but also geographical and ecological information in a phylogenetic framework. This will help not only with taxonomic identification, but will build the basis

to answer evolutionary questions about this group and to develop comparative analysis combining several sources of information. Here we present the first step of this undertake, an interactive key for the genus including all species described so far plus some taxa in the process of being described.

METHODS

Sources of information. *Herbarium and fieldwork.* We visited the main aroid collections in the United States, including the herbarium at the Missouri Botanical Garden (MO), the Field Museum (F), Gray Herbarium at Harvard (GH), The New York Botanical Garden (NY), and the National Herbarium (US). We also visited collections in Colombia (COL, COAH, JAUM, MEDEL, HUA, CUVC), Ecuador (QCA, QCNE, QPLS), Panama (PMA, SCZ), and Mexico (MEXU, UJAT, and XAL). Additionally, we studied digital images of the specimens from other herbaria that could not be visited. Type specimens were studied when available or digital images were use instead. Fieldwork was conducted in Colombia, Ecuador, Mexico and Panama.

Floral morphological and anatomical work. Due to the limited amount of vegetative characters of utility for *Monstera* species identification we explored morphological and anatomical characters from reproductive structures, in particular flowers. Anatomical work was conducted in the Plant Anatomy Lab at the National Herbarium (US, Smithsonian Institution) through a short-term visitor grant. We used flowers collected in the field and other available in the spirit collection at the National Herbarium (US) built by Dr. Dan Nicolson and Dr. R Eyde. . In brief, tissues were fixed

in FAA, and then transferred to 70 % ETOH until paraffin infiltration and embedding. We placed individual flowers in 10% Ethylenediamine at 60 °C for 24h to soft the trichosclereids and facilitate the sectioning. Then we infiltrated and embedded the tissues in paraffin wax (POLYFIN) using tert-butyl alcohol (TBA) as the intermediate solvent. All infiltration and embedding procedures we done under vacuum. After paraffin embedding, tissues were sectioned using a rotary microtome. Sections were stained using Toluidine blue and mounted for microscopy observation. Additional slides containing floral anatomy sections were obtained from Dr R. Eyde collection at the National Herbarium (US), and others provide by Dr. W. Carvell (Benedictine University, Lisle, IL)

Geographical database. We built a geographical database comprises a record for each specimen, and it includes collector and collection number, geographic location, and geographic coordinates, and herbarium where the specimen is deposited. Other information that may be present in herbarium collection, and that was collected includes, collection date, phenology and uses. When available digital databases were obtained directly from the herbaria, and then complete manually by the revision of the physical specimen. GBIF database records were used but only collections for which taxonomical identification was confirmed were used. Distribution maps will be built using scripts developed in the software R and the packages maps, ggmap, maptools, and raster (**Figure 1**).

Interactive key. We built a database to record morpho-anatomical information for each taxonomical unit (**Table 1**). The morpho-anatomical database was built using the software Lucid v3.5 (lucidcentral.org). It contains 108 ecological, morphological and anatomical traits from vegetative and reproductive structures and it is divided in three phenological stages, seedling, juvenile and adult plants. Within each stage characters are subdivided in the following order: habit, stem, leaves, inflorescence/infrutescence. The database includes discrete and continuous characters. We included 44 taxonomical units representing the currently 42 described species in the genus and other two new described species (Zuluaga et al. **Chapter 4, part 1**). The traits and their states were introduced to the lucid software using the Lucid Builder tool. Each taxonomical unit was coded or measure for each trait using field observations, literature, living plants, and spirit and herbarium collections deposited in herbaria in previously mentioned.

PRELIMINARY RESULTS

Interactive key implementation. The information in the interactive key includes data from 25 species studied *in vivo* during fieldwork or at botanical gardens, and ca. 4208 herbarium collections, 80% of them with geographic coordinates (**Table 2**). See **Figure 2** for views of the interactive key).

Anatomical work. 16 species were been embedded, and six have been sectioned and mounted. Additionally, slides for other six species were obtained from Dr. W. Carvell. Microscopy work has revealed heretofore underappreciated differences in style

length, stigma surface, distribution of trichosclereids and vascular bundles (see **Figures 3** for examples of anatomical characters).

The taxonomic work in the genus *Monstera* is being actively being developed. We still need to analyze and include all our anatomical studies into the interactive key, as well as describe and include all the potential new taxa for the genus. Our ultimate goal is to develop a database of geographic and morpho-anatomical data that will allow us to answer questions about the evolution of this enigmatic group.

REFERENCES

- Blagoderov V, Brake I, Penev L, Roberts D, Rycroft S, Smith VS. (2010) Blagoderov V, Brake I, Georgiev T, Penev L, Roberts D, Rycroft S, Scott B, Agosti D, Catapano T, Smith VS (2010) Streamlining taxonomic publication: a working example with Scratchpads and ZooKeys. *ZooKeys* 50: 17-28 doi: 10.3897/zookeys.50.539 [PMC free article] [PubMed]
- Brach, A.R., Song, H., 2005. ActKey: A Web-Based Interactive Identification Key Program. *Taxon* 54, 1041–1046. doi:10.2307/25065490
- Brach, A.R., Song, H., 2006. eFlorae: New directions for online floras exemplified by the Flora of China Project. *Taxon* 55, 188–192.
- Clark, B.R., Godfray, H.C.J., Kitching, I.J., Mayo, S.J., Scoble, M.J., 2009. Taxonomy as an eScience. *Philosophical Transactions of the Royal Society of London A: Mathematical, Physical and Engineering Sciences* 367, 953–966. doi:10.1098/rsta.2008.0190
- Dallwitz MJ., *A Comparison of Interactive Identification Programs. Institute of Botany, Chinese Academy of Sciences; 2011. (2 August 2013; <http://delta-intkey.com/www/comparison.htm>).*
- Dallwitz MJ, Paine TA, Zurcher EJ. *Principles of Interactive Keys. Institute of Botany, Chinese Academy of Sciences; 2013. (2 August 2013; <http://delta-intkey.com/www/interactivekeys.htm>).*
- eMonocot Team Araceae *CATE Araceae* <http://araceae.e-monocot.org/taxonomy/term/141> (accessed 20 April 2015)

Farnsworth, E.J., Chu, M., Kress, W.J., Neill, A.K., Best, J.H., Pickering, J., Stevenson, R.D., Courtney, G.W., VanDyk, J.K., Ellison, A.M., 2013. Next-Generation Field Guides. *BioScience* 63, 891–899. doi:10.1525/bio.2013.63.11.8

Haigh, A., Mayo, S.J., Croat, T., Reynolds, L., Pinto, M.M., Boyce, P.C., Lay, L., Bogner, J., Clark, B., Kostelac, C., Hay, A., 2009. Interactive web-taxonomy for the Araceae: www.cate-araceae.org. *Blumea - Biodiversity, Evolution and Biogeography of Plants* 54, 13–15. doi:10.3767/000651909X474032

Smith, V.S., Rycroft, S.D., Brake, I., Scott, B., Baker, E., Livermore, L., Blagoderov, V., Roberts, D., 2011. Scratchpads 2.0: a Virtual Research Environment supporting scholarly collaboration, communication and data publication in biodiversity science. *Zookeys* 53–70. doi:10.3897/zookeys.150.2193

Smith VS, Rycroft SD, Harman KT, Scott B, Roberts D. (2009) Scratchpads: a data-publishing framework to build, share and manage information on the diversity of life *BMC Bioinformatics* 10(Suppl 14): S6:

TABLES AND FIGURES

Table 1. List of geographical, ecological and morpho-anatomical characters and states used for the lucid key. Characters are divided in four sections; ecology and distribution, seedling, juvenile and adult plants. Within morphological section characters are subdivided by structures in the following order: habit, internodes, leaves, inflorescence/infrutescence.

Ecology and distribution		
Geographic distribution		
Height on the tree (adult plant)		
Elevation		
Seedling		
<i>Seedling habit</i>		
stolon-like		
with foliage leaves		
Juvenile plants		
<i>Growth habit</i>		
terrestrial creepers		
scandent		
adpressed climbers		
<i>Internodes</i>		
Internodes length		
Internodes diameter		
Internodes surface		
	smooth	
	verrucose	
	papillate-papillose	
<i>Leaves</i>		
Juvenile leaf host attachment		
	adpressed (shingled)	
	spreading from host	
Juvenile/adult leaf shape differentiation		
	not differentiated	
	differentiated	
Petiole		
	Petiole length in proportion to the leaf blade	
		longer than leaf blade
		shorter than leaf blade
		about as long as leaf blade
	Petiole length	
	Petiole diameter	
	Sheathing section	

Blades	persistence	deciduos marcescent persistent fibers-like
	Sheathing section lenght in proportion to the petiole	less than 1/2 between 1/2 and 3/4 all the petiole
	Sheathing section lenght	
	Symmetry	Equilateral Inaequilateral
	Shape	falcate lanceolate oblong rounded elliptic ovate
	Blade apex	acute gradually acuminate abrutly acuminate
	Blade base	attenuate oblique obtuse cuneate subcordate cordate
	Blade coloration	concolorous bicolorous variegated
	Blade texture	membranaceous chartaceous subcoriaceus coriaceus
	Perforations	present absent
	Blade overall length	
	Blade overall maximum	
	width	
	Adult plants	
	<i>Growth habit</i>	
	vine	
	adpressed climber	
	<i>Stem</i>	
	Internodes length/width	

relation	longer than broad broader than long	
Adult internode length		
Adult internode/Stem diameter		
Leaf scars wide		
Axillary buds		
Stem cross section	subcylindric dorsiventrally flattened flattened laterally	
Stem surface	smooth papillate verrucose striate speckled lineated raised ridged lenticellate	
Stem epidermis (when dry)	flaking no flaking	
<i>Cataphylls</i>		
Cataphyll number		
Cataphyll length		
<i>Leaves</i>		
Spatial disposition	along of the stem apex clustered	
Petiole	Petiole cross-sectional shape	subterete to terete C to D-shaped or broader than thick U-shaped or thicker than broad
	Petiole surface	smooth verrucose glaucous papillate white dotted or flecks green dotted
	Petiole length Petiole diameter Petiole length in proportion to the leaf blade	longer than leaf blade shorter than leaf blade about as long as leaf blade

Blades	Petiole/Blade length ratio		
	Petiole upper (adaxial) surface	sulcate flattened with medial rib convex	
	Petiole lower (abaxial) surface	rounded angular ribbed	
	Geniculum	Geniculum length	
	Sheathing section	Sheathing section length in proportion to the petiole	less than 1/2 between 1/2 and 3/4 more than 3/4
		Persistence	deciduous fibers-like deciduous intact fiber persistent marcescent persistent
		Sheath "vernation"	no margins acutely raised winged obtusely raised
		Sheathing section length sheath extension (beyond geniculum)	
	Shape	lanceolate falcate elliptic ovate oblong rounded	
	Symmetry	equilateral inaequilateral	
	Blade apex	obtuse to weakly emarginate acute mucronate gradually acuminate	

Blade base	abruptly acuminate
	attenuate
	obtuse
	truncate
	cuneate
	subcordate
	oblique
Blade margins	cordate
	entire
	pinnatifid
	lacerate
	lobed
	one side pinnatifid
Blade overall length	
Blade overall maximum width	
Proportion Length/width	
Blade coloration	concolorous
	bicolorous
Blade glossiness on upper surface	
	matte
	semiglossy
	glossy
	velvety
Blade glossiness on lower surface	
	matte
	velvety
	semiglossy
	glossy
Blade texture	
	membranaceous
	chartaceous
	coriaceous
	subcoriaceous
Blade color when dried on upper surface	
	yellowish
	greenish to olive-green
	grayish
	brownish
	dark brown
	blackish
	reddish brown
Blade color when dried on lower surface	
	yellowish
	greenish to olive-green
	grayish
	brownish

	dark brown blackish reddish brown	
Midrib	Midrib shape above	acute convex or round raised flat sunken
	Midrib shape below	acute convex or round flat quadrangular (square to rectangular) multi-ribbed
Primary lateral veins	Number of pairs of primary lateral veins Distance between primary lateral veins Departing angle of primary lateral veins Primary lateral veins appearance above	sunken acute convex flat or obscure
	Primary lateral veins appearance below	flat or obscure acute sunken convex
Secondary veins	reticulate parallel (only reticulate close to the margin)	
Perforations (fenestrations)	Perforations spatial distribution	irregular absent in series
	Number of perforation series	
Number of segments (lobes) per side Distance between pinnae Difference between wider		

	and thinner side	
	Pinnae width	
<i>Flowering shoots</i>	erect	
	pendent	
<i>Inflorescence</i>		
Number per axile		
Spatial position	erect	
	spreading	
	pendent	
Length in proportion to leaves	shorter than leaves	
	longer than leaves	
	about as long as leaves	
Peduncle	Surface	smooth
		papillate
		verrucose
		whitish dotted
	Peduncle length	
	Peduncle diameter	
Spathe	Spathe shape (when flattened)	ovate
		elliptic
		oblong
		obovate
		subcircular
	Spathe color outside	white to cream
		greenish white
		green
		yellow to orange
		light yellow
		pinkish
	Spathe color inside	greenish white
		green
		white to cream
		light yellow
		yellow to orange
		pinkish
	Spathe length	
	Spathe width	
	Length in proportion to spadix	much longer than spadix
		slightly longer than the spadix

Spadix	Spadix length in proportion to peduncle	shorter than peduncle
		longer than peduncle
		about as long as peduncle
	Spadix shape	cylindroid
		apex tapered
		clavate
		base tapered
	Spadix color (Anthesis)	white to cream
		yellow to yellowish
		orange to orangish
		pinkish
Portion of sterile flowers	absent	
	present	
Flowers	Spadix frontal length	
	Spadix posterior length	
	Spadix diameter	
	Length in proportion to peduncle (number)	
	Shape	prismatic
		cilindrical
		conical
	Pistil	conical acuminate
		style
		truncatae style
<i>Infructescence</i>		
Spatial disposition	erect	
	pendent	
Infructescence color (before stylar cap fall off)	white to cream	
	greenish white	
	green	
	light yellow	
	orange	
Infructescence peduncle length		
Infructescence peduncle diameter		
Infructescence length		
Infructescence diameter		
Seeds	Seed number	
		1
		2
		3

Seed color

green
brown
gray/white
black

Seed shape

oblong
spheroidal

Seed length

Seed diameter

Table 2. Number of specimens studied per species. Excluded specimens included specimens with ambiguous identification or others that represent new taxa to be described.

Species Name	Number of specimens
<i>Monstera acacoyaguensis</i>	12
<i>Monstera acuminata</i>	147
<i>Monstera adansonii</i>	1013
<i>Monstera amargalensis</i>	3
<i>Monstera anomala</i> sp. nov.	20
<i>Monstera aureopinnata</i>	69
<i>Monstera barrieri</i>	6
<i>Monstera buseyi</i>	49
<i>Monstera cenepensis</i>	3
<i>Monstera costaricensis</i>	24
<i>Monstera deliciosa</i>	166
<i>Monstera dilacerata</i>	55
<i>Monstera dissecta</i>	82
<i>Monstera dubia</i>	279
<i>Monstera egregia</i>	12
<i>Monstera epipremnoides</i>	16
<i>Monstera filamentosa</i>	23
<i>Monstera florescanoana</i>	2
<i>Monstera glaucescens</i>	89
<i>Monstera gracilis</i>	41
<i>Monstera kessleri</i>	2
<i>Monstera lechleriana</i>	243
<i>Monstera lentii</i>	18
<i>Monstera luteynii</i>	4
<i>Monstera maderaverde</i>	2
<i>Monstera membranacea</i>	23
<i>Monstera minima</i>	3
<i>Monstera molinae</i>	19
<i>Monstera obliqua</i>	709
<i>Monstera oreophila</i>	103
<i>Monstera pinnatipartita</i>	161
<i>Monstera pittieri</i>	55
<i>Monstera planadaensis</i>	5
<i>Monstera punctulata</i>	39
<i>Monstera puntarenense</i>	2
<i>Monstera siltepecana</i>	158

<i>Monstera spruceana</i>	256
<i>Monstera standleyana</i>	44
<i>Monstera subpinnata</i>	107
<i>Monstera tenuis</i>	53
<i>Monstera tuberculata</i>	64
<i>Monstera vasquezii</i>	4
<i>Monstera xanthospatha</i>	19
<i>Monstera integrifolia</i> sp. nov.	4
Total	4208
Excluded specimens	791

Figure 1. An example of distribution maps for *Monstera* species to be included in the interactive key.




Figure 2. View of the Lucid Player tool. Four windows are available. Top left window contains the characters (features available); bottom left contain the characters chosen; top right contain the taxa remaining; and bottom left the taxa discarded.

KeyFeaturesEntitesViewWindowHelp

Features Available: 108

▼ Ecology and distribution

▶ Geographic distribution

Elevation

▶ Seeding

▶ Seeding habit

☒ SIDON-LIKE

▶ Juvenile plants

☐ with foliage leaves

▶ Adult plants

▶ Growth habit

▶ Stem

▶ Cataphylls

▶ Leaves

▶ Flowering shoots

▶ Inflorescence

Number per axile

▶ Spatial position

▶ Length in proportion to leaves

▶ Peduncle

▶ Spathe

▶ Spadix

▼ Monstera 2

▼ Ecology and distribution

▶ Geographic distribution

☒ Brazil

☒ French Guiana

▶ Seeding

▶ Seeding habit

☒ SIDON-LIKE

▶ Monstera 3

▶ Monstera aureoipinnata Croat

▶ **Monstera dubia (Kunt) Engl. & K. Krause**

▶ Monstera spruceana (Schott) Engl.

▶ Monstera 50

▶ Monstera acacovaguensis Maunda

▶ Monstera acuminata K. Koch

▶ Monstera adansonii Schott

▶ Monstera amargaliensis Croat & M. M. Mora

▶ Monstera barrieri Croat

▶ Monstera bocabotrana

▶ Monstera Duseyl Croat & Grayum

▶ Monstera ceneperis Croat

▶ Monstera chlamense

▶ Monstera codruse

▶ Monstera coloradense

▶ Monstera costaricensis (Engl. & K. Krause) Croat & Grayum

▶ Monstera cobdruense

▶ Monstera delicosa Liebm.

▶ Monstera discida (Schott) Croat & Grayum

▶ Monstera dyerii

▶ Monstera epipremnoides Engl.

▶ Monstera filamentosa Croat & Grayum

▶ Monstera filamentosana Croat, T. Koerner & Acebey

▶ Monstera glaucescens Croat & Grayum

▶ Monstera gracilis Engl.

▶ Monstera hammellii

TreesLists

TreesListsImages

Monstera dubia (Kunt) Engl. & K. Krause




◀▶▶▶

⌂

⌕

Figure 3. Examples of morpho-anatomical traits from flowers in *Monstera*. **A, B, D, & E** show different flower shapes present in *Monstera*; **A**, *M. punctulata*; **B**, *M. tuberculata*; **D**, *M. integrifolia*; **E**, *M. lentii*. **C & G** show cross section from the stylar region (**C**), and the ovary (**D**) of *M. punctulata*; distribution of vascular bundles and trichosclereids are visible. **G**, cross section of the stylar region of *M. punctulata*; druses and rhapides are visible. **H**, cross section to the stigma level of *M. pittieri* using polarized light; druses are visible.



APPENDIX 1. Zuluaga, A., K. Cameron, T. Croat & M. Medecilo. 2015. Testing the monophyly of *Spathiphyllum*, and the relationship between Asian and tropical American species. *Aroideana* 38E(1):107-115.

Testing the monophyly of *Spathiphyllum*, and the relationship between Asian and tropical American species.

Alejandro Zuluaga and Kenneth Cameron
Department of Botany
University of Wisconsin-Madison,
Madison, WI 53706

Thomas Croat
P. A. Schulze Curator of Botany
Missouri Botanical Garden
P.O. Box 299, St. Louis, MO 63166

Melanie Medecilo
Biological Sciences Department,
De La Salle University-Dasmarias
Cavite, Philippines

ABSTRACT

As part of a broader phylogenetic study of the subfamily Monsteroideae we used three plastid and one nuclear markers to test the monophyly of the genus *Spathiphyllum*. Our results support the tribe Spathiphyllae as sister of the remaining Monsteroideae, and the monophyly of *Spathiphyllum*. The monotypic genus *Holochlamys* appears nested within *Spathiphyllum* in a clade with the Asian species and *S. cannifolium*. However, support for this clade is low. The section *Spathiphyllum* shows a high support and is sister to the remaining species in the genus. As a consequence *Spathiphyllum* remains as

the only disjunct aroid genus between the Old and New World Tropics.

INTRODUCTION

With the increased application of molecular data for phylogenetic analysis, the understanding of relationships within Araceae has been improved considerably (Cabrera, 2008, Cusimano, 2011, Henríquez, 2014). Additionally, new fossils and biogeographical studies have helped to elucidate the historical and current geographic distribution patterns in the family (Nauheimer, 2012a). As a result, aroid classification has changed

dramatically; several genera have been resurrected; and new genera have been created (Cabrera, 2008, Cusimano, 2011). Nevertheless, many critical issues are still in need of more data.

Before this molecular era there were thought to be three geographically disjunct genera between the old world and new world tropics. Species in the genera *Homalomena*, *Schismatoglottis* and *Spathiphyllum* were found in both tropical America and tropical Asia, particularly within the Malay Archipelago and Melanesia (van Steenis, 1962, Grayum, 1990, Mayo et al., 1997). However, recent studies (Barabé, 2002, Gauthier, 2008, Wong et al., 2013) have shown evidence challenging the monophyly of disjunct *Schismatoglottis* and *Homalomena* (they are no longer disjunct in their revised circumscriptions), leaving *Spathiphyllum* as the only remaining aroid genus with a tropical disjunction.

The genus *Schismatoglottis*, with only three species in South America, and about 150 in tropical Asia, belongs to the tribe Schismatoglottidae that in its current circumscription comprises 11 genera. Several of these genera are small groups recognized thanks to molecular data (Wong et al., 2010a, 2010b, Low, 2014). The three Neotropical species previously recognized in *Schismatoglottis* are now in the resurrected genus *Philonotia* (Wong, 2010c) that belong to its own tribe Philonotieae. Equally, the genus *Homalomena* is considerable more diverse in Tropical Asia with about 500 species, versus only 10 species in Tropical

America (Boyce & Croat, 2013). Unlike *Schismatoglottis*, molecular evidence for the separation of Neotropical *Homalomena* is not conclusive (Barabé, 2002, Gauthier, 2008, Wong et al., 2013). The most plausible options for the assignment of the Neotropical *Homalomena* would be the resurrection of the genus *Adelomena*, or the combination of these species of *Homalomena* with *Philodendron* subgenus *Pterosmischum* in the genus *Elopium* (Wong et al., 2013). Thus, more extensive sampling and more molecular data are necessary to fully understand the relationships among *Homalomena*, and the subgenera of *Philodendron*.

Finally, we have the genus *Spathiphyllum*, which has a different distribution pattern. The genus is more diverse in Tropical America with about 50 species (Cardona, 2004), and only three species in Southeast Asia. *Spathiphyllum commutatum* Schott is widespread in Malesia and reaches Micronesia; *S. solomonense* Nicolson is restricted to the Solomon Islands; and *S. schlechteri* (Engl. & K.Krause) Nicolson is restricted to New Guinea. Neotropical species are concentrated in the northern Andes, with only a few species reaching Central America and Mexico (Cardona, 2004), and one species, *S. leave* reaches Cocos Island in the Pacific Ocean west of Costa Rica. No comprehensive phylogenetic studies, until now, have included more than three *Spathiphyllum* from Tropical America, or any of the Asian species.

Spathiphyllum together with *Holochlamys becarii* form the tribe Spathiphyllae in the subfamily Monsteroideae (Cabrera, 2008, Cusimano, 2011). The relationships of the tribe Spathiphyllae within Monsteroideae have been controversial, but recent studies strongly support Spathiphyllae as sister to the remaining Monsteroideae (Chartier, 2014, Henriquez, 2014). These two genera are very similar morphologically and ecologically, differing mainly in a few characteristics of the flowers. Whereas *Spathiphyllum* species have two to three locules and ovules with axillar placentation, *Holochlamys* has one locule and ovules with basal placentation (Bunting, 1960, Mayo et al., 1997). Nevertheless, anatomical studies suggest that the *Holochlamys* ovary could actually have two locules, and the placentation may not be basal (Eyde, 1967, Carvell, 1989).

Currently, *Spathiphyllum* is divided into four sections based the fusion of the perianth segments, the form and length of the pistils relative to the perianth, and the attachment the spathe to the peduncle (Bunting, 1960). Asian species together with *S. cannifolium* and *S. leave* form the section *Massowia*, which is recognized by having the perianth segments completely connate forming a cup around the pistil. This trait led Bunting (1960) to propose the monophyly of this group. However, Williams & Dressler (1967) and Grayum (1984, 1990) suggested this could be an artificial group with *Holochlamys* nested within it. Sections *Amomophyllum*, *Dysspathiphyllum*, and *Spathiphyllum* have free perianth segments.

As part of a broader systematic study of the subfamily Monsteroideae (Zuluaga, unpublished data) we tested the monophyly of *Spathiphyllum* and *Holochlamys*, and investigated the relationships of the Asian and American species of *Spathiphyllum*.

METHODS

Taxon Sampling and DNA sequencing. During the sampling for the study of systematics of the subfamily Monsteroideae we sequenced a total of seven accessions of the genus *Spathiphyllum* from Tropical America, *S. commutatum* and *S. solomonense* from Tropical Asia, and one accession of *Holochlamys becarii*. Additional sequences for *S. phrynifolium* were obtained from GENBANK. For this study we used representatives from all ten additional Monsteroideae genera, plus six other aroids as outgroups.

DNA was extracted from silica-dried plant material using the DNeasy Plant Mini Kit (Qiagen, Valencia, CA, USA) according to the manufacturer's specifications. Some samples were extracted using a CTAB method adapted from Doyle & Doyle (1987). We sequenced four plastid markers, *matK*, *rbcL*, *trnC-petN* spacer, and partial *ycf1*. PCR products were purified using ExoSap-It (Cleveland, OH). Sequencing reactions used the BigDye Terminator Cycle Sequencing Kit, and cleaned using Agencourt CleanSeq (Beverly, MA) magnetic beads. BigDye Terminator Cycle Sequencing Ready Reaction Kit (Applied Biosystems, Foster City, CA, USA). Samples

were sequenced using PE-Biosystems version 3.7 of Sequencing Analysis at the University of Wisconsin–Madison Biotechnology Center.

Molecular analysis. All sequences were manually edited in Geneious 6.0 (Biomatters, <http://www.geneious.com>). Sequences were aligned in Geneious 6.0 using a combination of Muscle (Edgar 2004) and MAFFT (Kato 2002). Finally, all alignments were checked manually. We performed individual and combined phylogenetic analyses. Analyses for each marker/dataset were performed under Maximum Likelihood using RaxML 7.0 (Stamatakis, 2006), and MrBayes 3.1.2 on XSEDE, as implemented in the CIPRES portal. For RaxML, we used the GTRGAMMA model in all cases. For Bayesian analysis models were calculated using Jmodeltest2 (Darriba, 2012).

RESULTS.

So far at least one of the four molecular markers has been successfully sequenced for *S. cannifolium* (two accessions), *S. cochlearispathum*, *S. commutatum*, *S. floribundum*, *S. minor*, *S. patulinervum*, *S. solomonense*, *S. wallisii*, and *Holochlamys beccarii*. Only the *trnC-petN* spacer was sequenced for the Asian *Spathiphyllum*. The combined alignment length is 4970 bp with 36% missing data. Despite using four molecular markers the identical sites between pairs of species within *Spathiphyllum* is more than 97%. This low variation is common within the genera

of the subfamily Monsteroideae (Unpublished data).

Trees derived from individual gene analyses are congruent, however the resolution is low except for a clade comprising the species of the section *Spathiphyllum*. Thus, a combined gene analysis is presented (**Figure 1**). All individual and combined analyses show the tribe Spathiphyllae as monophyletic and sister to the remaining Monsteroideae, thus supporting the topology presented by Chartier (2014) and Henríquez (2014). Section *Spathiphyllum* represented by *S. cochlearispathum*, *S. phrynifolium*, and *S. wallisii* shows strong support, and is sister to the remaining Spathiphyllae. The genus *Holochlamys* appears nested within *Spathiphyllum* in a clade with the Asian species and one accession of *S. cannifolium* (section *Massowia*). Section *Amomophyllum* (*S. minor*, *S. floribundum*, *S. patulinervum*) and one accession of *S. cannifolium* are sister to the clade containing *Holochlamys*. However support for the last two clades is low (Figure 1).

DISCUSSION.

A monophyletic tribe Spathiphyllae has been supported since the first molecular studies in Araceae, based on only a few genes, were published (Tam, 2004, Cabrera, 2008). However, the position of Spathiphyllae as sister to rest of Monsteroideae is only recovered in the most recent studies with more data included




Figure 1. Maximum likelihood tree from RaxML using four plastid markers (*matK*, *rbcL*, *trnC-petN* spacer, and partial *ycf1*). Bootstrap supports >75 and Bayesian posterior probabilities >0.85 are shown on the branches.

(Chartier, 2014, Henriquez, 2014). Our results strongly support this topology (**Figure 1**). Most of the synapomorphies defining Monsteroideae as presented in Cusimano (2011) seem to be more appropriate to be assigned to the clade containing all Monsteroideae but Spathiphyllae. These synapomorphies and the alternative characters in Spathiphyllae are: pollen zonate (ring-like) (vs. pollen multiaperturate with exine surface

striate/polyplicate); spathe soon deciduous with a distinct basal abscission (vs. Spathe marcescent), large trichosclereids not in bundles (vs. smaller trichosclereids in bundles). The only synapomorphy remaining for Monsteroideae is the presence of trichosclereids.

Within the tribe Spathiphyllae, the support for the position of *H. becarii* is not conclusive, but some markers suggest this

species may belong to the genus *Spathiphyllum*. *Holochlamys* appears consistently embedded in a clade with section *Massonia*. These two genera are very similar in habit and morphology (Bunting, 1960), stem vasculature (French & Tomlison, 1981), palynological features (Grayum, 1992), and floral anatomy (Carvell, 1989). Moreover, *Holochlamys* shares with section *Massonia* the presence of fused perianth parts forming a cup around the pistil (Bunting, 1960, Carvell, 1989). The proposed differences between these two genera - a unilocular ovary, and ovules with basal placentation - have been challenged by more detailed anatomical studies (Eyde, 1967). Carvell (1989) undertook a very detailed anatomical study of Monsteroideae and Pothoideae, and reported unilocular ovaries, and basal ovules for *H. beccarii*. However, he also reported the presence of an incomplete septum in several flowers, and the attachment of some of the ovules to that septum. Additionally, Carvell (1989) found considerable intra- and interspecific variation in locule, ovules, tepals, and stamen numbers in *Spathiphyllum*. Despite the fact that we consider there to be no strong morphological evidence to separate *Holochlamys* from *Spathiphyllum* (especially from the section *Massonia*), our analysis currently lacks adequate resolution within this clade, and so we prefer to be cautious at this point in time, necessarily gathering more data before proposing any taxonomic changes for these species.

Our sampling did not allow a complete assessment of the sectional classification in

Spathiphyllum as proposed by Bunting (1960). The only section with strong support through all our analysis is the section *Spathiphyllum*, which is also supported by the morphology of their flowers with a long and conical pistil exceeding the perianth (Bunting, 1960). Williams and Dressler (1967), and Grayum (1984, 1990), discussed hypotheses for the origin and biogeography of *Spathiphyllum* and *Holochlamys*. We are undertaking additional molecular and biogeographical analyses of the subfamily Monsteroideae, and more detailed discussion on this topic will be presented in subsequent papers.

REFERENCES

- Barabé, D., A. Bruneau, F. Forest & C. Lacroix, 2002. The correlation between development of atypical bisexual flowers and phylogeny in the Aroideae (Araceae). *Plant. Syst. Evol.* 232: 1–19.
- Boyce, P.C. & T.B. Croat. 2013. The Überlist of Araceae, totals for published and estimated number of species in aroid genera. Available at <http://www.aroid.org/genera/130307uberlist.pdf> (accessed November 2014).
- Bunting, G. S. 1960. A revision of the genus *Spathiphyllum* (Araceae). *Mem. New York Bot. Gard.* 10: 1–54.

- Cabrera, L. I., G. A. Salazar, M. W. Chase, S. J. Mayo, J. Bogner, & P. Davila. 2008. Phylogenetic relationships of aroids and duckweeds (Araceae) inferred from coding and noncoding plastid DNA. *Am. J. Botany* 95(9): 1153–1165.
- Cardona F. 2004 Synopsis of the genus *Spathiphyllum* (Araceae) in Colombia. *Ann. Missouri Bot. Gard.* 91(3): 448–456
- Carvell, W.N. 1989. *Floral anatomy of the Pothoideae and Monsteroideae (Araceae)*. PhD Dissertation. Miami University. Miami, FL, USA.
- Chartier, M., M.Gibernau & S.S.Renner. 2013. The evolution of pollinator-plant Interaction types in the Araceae. *Evolution* 68(5): 1533–1543
- Cusimano, N., J. Bogner, S.J. Mayo, P.C. Boyce, S.Y. Wong, M. Hesse, W.L.A. Hetterscheid, R. C. Keating, & J. C. French. 2011. Relationships within the Araceae: Comparison of morphological patterns with molecular phylogenies. *Am. J. Botany* 98(4): 1–15.
- Eyde, R.H., D.H. Nicolson, & P. Sherwin. 1967. A survey of floral anatomy in Araceae. *Amer. J. Bot.* 54: 478–497.
- French, J.C. & P.B. Tomlinson. 1981. Vascular patterns in stems of Araceae: subfamily Pothoideae. *Amer. J. Bot.* 68(5): 713–729.
- Gauthier, M. P.L., D.Barabe & A. Bruneau. 2008. Molecular phylogeny of the genus *Philodendron* (Araceae): delimitation and infrageneric classification. *Bot. J. Linn. Soc.* 156(1): 13–27.
- Grayum, M.H. 1990. Evolution and phylogeny of the Araceae. *Ann. Missouri Bot. Gard.* 77: 628–697.
- Grayum, M.H. 1992. Comparative external pollen ultrastructure of the Araceae and putatively related taxa. *Monogr. Syst. Bot. Mo. Bot. Gard.* 43: 1–167.
- Gonçalves, E.G. 2012. *Lorenzia* (Araceae-Spathicarpeae): a new genus from northern Brazil supported by *matK* sequence data. *Syst. Bot.* 37(1): 48–52.
- Low, S.L., S.Y. Wong & P.C. Boyce. 2014. *Schottarum* (Schismatoglottideae: Araceae) substantiated based on combined nuclear and plastid DNA sequences. *Plant. Syst. Evol.* 300(4): 607–617.
- Mayo, S.J., J. Bogner & P.C. Boyce. 1997. The Genera of Araceae. Royal Botanic Gardens, Kew, UK.

- Nauheimer, L., D. Metzler & S.S. Renner. 2012a. Global history of the ancient monocot family Araceae inferred with models accounting for past continental positions and previous ranges based on fossils. *New Phytologist* 195: 938–950.
- Nauheimer, L., P.C. Boyce & S.S. Renner. 2012b. Giant taro and its relatives: A phylogeny of the large genus *Alocasia* (Araceae) sheds light on Miocene floristic exchange in the Malesian region. *Mol. Phylogenet. Evol.* 63(1): 43–51.
- Nauheimer, L., & P.C. Boyce 2014. *Englerarum* (Araceae, Aroideae): a new genus supported by plastid and nuclear phylogenies. *Plant. Syst. Evol.* 300(4): 709–715.
- Stamatakis, A., 2006a. RAxML-VI-HPC: maximum likelihood-based phylogenetic analyses with thousands of taxa and mixed models. *Bioinformatics*, 22(21): 2688–2690.
- Stamatakis, A., 2006b. Phylogenetic models of rate heterogeneity: a high performance computing perspective. In: Proceedings of the 20th IEEE/ACM International Parallel and Distributed Processing Symposium (IPDPS2006), Rhodes, Greece.
- Stamatakis, A., P. Hoover & J. Rougemont. 2008. A rapid bootstrap algorithm for the RAxML Web servers. *Syst. Biol.* 57(5): 758–771.
- Swofford, D.L., 1991. PAUP: Phylogenetic Analysis Using Parsimony, Version 3.1 Computer program distributed by the Illinois Natural History Survey, Champaign, Illinois.
- Steenis, C.G.G.J. van. 1962. The land-bridge theory in botany with particular reference to tropical plants. *Blumea* 11: 235–372
- Tam, S., P.C. Boyce, T. Upson, D. Barabé, A. Bruneau, F. Forest & J.S. Parker. 2004. Intergeneric and Intrafamilial Phylogeny of Subfamily Monsteroideae (Araceae) Revealed by Chloroplast trnL-F Sequences. *Am. J. Botany* 91(3): 490–498.
- Wong, S.Y. & P.C. Boyce. 2010a. Studies on Schismatoglottideae (Araceae) of Borneo IX: A new genus, *Hestia*, and resurrection of *Apoballis*. *Bot. Stud. (Taipei)*, 51: 249–255
- Wong, S. Y. P.C. & Boyce, P. C. 2010b. Studies on Schismatoglottideae (Araceae) of Borneo XI: *Ooia*, a new genus, and a new generic delimitation for *Piptospatha*. *Bot. Stud. (Taipei)* 51: 543–552.

Wong, S.Y., P.C. Boyce, A.S.B. Othman & L.C. Pin. 2010c. Molecular phylogeny of tribe Schismatoglottideae (Araceae) based on two plastid markers and recognition of a new tribe, Philonotieae, from the Neotropics. *Taxon* 59(1): 117–124.

Wong, S.Y. T.P. Jean, N.K. Kiaw, A.S.B. Othman, L.H. Boon, F.B. Ahmad & P.C. Boyce. 2013. Phylogeny of Asian *Homalomena* (Araceae) based on the ITS Region Combined with Morphological and Chemical Data. *Syst. Bot.* 38(3): 589–599.