



Dimetane Extentabs advertisement.

[s.l.]: [s.n.], 1968

<https://digital.library.wisc.edu/1711.dl/AW4SVRNIK2IIF8O>

<http://rightsstatements.org/vocab/InC/1.0/>

The libraries provide public access to a wide range of material, including online exhibits, digitized collections, archival finding aids, our catalog, online articles, and a growing range of materials in many media.

When possible, we provide rights information in catalog records, finding aids, and other metadata that accompanies collections or items. However, it is always the user's obligation to evaluate copyright and rights issues in light of their own use.


To keep the dust-sensitive patient comfortable.

When patients are sensitive to dust and other inhalant allergens, Dimetane® (brompheniramine maleate) works hard. Each Extentab® helps keep patients symptom free for 10 to 12 hours and does this with little chance of drowsiness or overstimulation. Dimetane works in stubborn as well as routine allergies and in patients who can't tolerate other antihistamines.

Try Dimetane first whenever allergic symptoms appear and give it if allergens cannot be completely avoided or hyposensitization completely achieved.

Try these ideas to help keep a room dust-free and comfortable

The dust-free room need not be a room stripped bare. You can tell patients or their parents how to make such rooms comfortable, pleasant and colorful yet free of dust collectors and other allergenic articles.



GIVE DIMETANE® Extentabs® FIRST!

(BROMPHENIRAMINE MALEATE)

INDICATIONS: For the prevention and symptomatic relief of many allergic manifestations as in hay fever, conjunctivitis, angioneurotic edema, pruritus, rhinitis, atopic eczema, urticaria, dermatitis, bronchial asthma, common cold, drug reaction, rhus dermatitis, insect and spider bites.

CONTRAINDICATIONS: Hypersensitivity to antihistamines. Not recommended for use during pregnancy.

PRECAUTIONS: Until response is determined, patient should be cautioned against engaging in mechanical operations requiring alertness.

SIDE EFFECTS: Hypersensitivity reactions, including skin rashes, urticaria, hypotension, and thrombocytopenia, have been reported rarely. Occasional transitory drowsiness, lassitude, nausea, giddiness, dryness of the mouth, mydriasis, increased irritability or excitement may be encountered.

See package insert for full prescribing information.

**Works all day
or all night
protecting the
“sensitives”**

A-H-ROBINS
A. H. ROBINS COMPANY
RICHMOND, VA. 23220

