



LIBRARIES
UNIVERSITY OF WISCONSIN - MADISON

Tussionex advertisement.

[s.l.]: [s.n.], 1962


<https://digital.library.wisc.edu/1711.dl/2EDBYPLFWG27I8Z>

<http://rightsstatements.org/vocab/InC/1.0/>

The libraries provide public access to a wide range of material, including online exhibits, digitized collections, archival finding aids, our catalog, online articles, and a growing range of materials in many media.

When possible, we provide rights information in catalog records, finding aids, and other metadata that accompanies collections or items. However, it is always the user's obligation to evaluate copyright and rights issues in light of their own use.

How to crack a hard-boiled cough



(Dihydrocodeinone and Phenyltoloxamine Resins)

TUSSIONEX[®]

'STRASIONIC'
ANTITUSSIVE

A SINGLE TEASPOON STOPS PERSISTENT COUGH 8-12 HOURS

... yet does not interfere with desirable expectoration. This prolonged relief is made possible through a cation exchange ('Strasionic') formulation of dihydrocodeinone potentiated by phenyltoloxamine.¹⁻³ Prescribe Tussionex for dependable cough control with less drug, at lower cost to your patient. Just two ounces usually lasts six days and nights.

TUSSIONEX SUSPENSION

Each teaspoonful (5 cc.) contains:
Dihydrocodeinone 5 mg.
(Warning: May be habit-forming)
Phenyltoloxamine 10 mg.

TUSSIONEX TABLETS

Each tablet contains:
Dihydrocodeinone 5 mg.
(Warning: May be habit-forming)
Phenyltoloxamine 10 mg.

As cation exchange resin complexes of sulfonated polystyrene

DOSAGE: Administer q 8-12 h as follows: Adults—1 teaspoonful or tablet. Children—over 5 years, 1 teaspoonful; 1-5 years, ½ teaspoonful; under 1 year, ¼ teaspoonful.

SIDE EFFECTS: Occasional mild constipation, nausea, facial pruritus, or drowsiness. **CONTRAINDICATIONS:** Head injuries or other neurosurgical conditions; history of narcotic addiction.

CLASS B NARCOTIC. On oral Rx where state law permits.

References: 1. Cass, L. J., and Frederik, W. S.: *Ann. Int. Med.* 49:151, 1958. 2. Chan, Y. T., and Hays, E. E.: *Am. J. M. Sc.* 234:207, 1957. 3. Townsend, E. H.: *New England J. Med.* 258:63, 1958.

STRASBURGH LABORATORIES
ROCHESTER, NEW YORK, U.S.A.  **LABORATORIES**
DIV. WALLACE & TIERNAN INC.