



LIBRARIES
UNIVERSITY OF WISCONSIN - MADISON

Phenaphen with Codeine advertisement.


[s.l.]: [s.n.], 1963

<https://digital.library.wisc.edu/1711.dl/5GMU6FYRPVM2J86>

<http://rightsstatements.org/vocab/InC/1.0/>

The libraries provide public access to a wide range of material, including online exhibits, digitized collections, archival finding aids, our catalog, online articles, and a growing range of materials in many media.

When possible, we provide rights information in catalog records, finding aids, and other metadata that accompanies collections or items. However, it is always the user's obligation to evaluate copyright and rights issues in light of their own use.



when you want
 "the best out of codeine"
 with minimal risk of
 codeine side effects

With PHENAPHEN WITH CODEINE, potentiation and sedative enhancement induce analgesia on doses of codeine so low as to be ineffective when given alone. Side effects are rare: occasional mild nausea, constipation or drowsiness. Contraindicated only in patients hypersensitive to any ingredient.



PHENAPHEN[®] with CODEINE

	Phenaphen No. 2	Phenaphen No. 3	Phenaphen No. 4
Each	194.0 mg.	194.0 mg.	194.0 mg.
PHENAPHEN	162.0 mg.	162.0 mg.	162.0 mg.
WITH CODEINE	0.031 mg.	0.031 mg.	0.031 mg.
Capsule	16.2 mg.	16.2 mg.	16.2 mg.
contains:	16.2 mg.	16.2 mg.	16.2 mg.
	(Warning: May be habit-forming)		
	16.2 mg.	(1/2 gr.) 32.4 mg.	(1 gr.) 64.8 mg.
	(Warning: May be habit-forming)		

Also available:
 PHENAPHEN
 —same formula,
 without codeine.

A. H. ROBINS COMPANY, INC., Richmond 20, Virginia