



**LIBRARIES**

UNIVERSITY OF WISCONSIN-MADISON

## **A compendium of research on angling regulations for brook trout conducted at Lawrence Creek, Wisconsin. Report 54 1970**

Hunt, Robert L. (Robert Leroy) 1933-  
Madison, Wisconsin: Dept. of Natural Resources, 1970

<https://digital.library.wisc.edu/1711.dl/VH5IZFAEOESLV8A>

<http://rightsstatements.org/vocab/InC/1.0/>

For information on re-use see:

<http://digital.library.wisc.edu/1711.dl/Copyright>

The libraries provide public access to a wide range of material, including online exhibits, digitized collections, archival finding aids, our catalog, online articles, and a growing range of materials in many media.

When possible, we provide rights information in catalog records, finding aids, and other metadata that accompanies collections or items. However, it is always the user's obligation to evaluate copyright and rights issues in light of their own use.

Wis Doc  
Nat.  
3:  
R 4/  
54  
c. 5

**RESEARCH  
REPORT 54**

**LOAN COPY**

**A COMPENDIUM OF RESEARCH ON  
ANGLING REGULATIONS FOR BROOK TROUT  
CONDUCTED AT LAWRENCE CREEK, WISCONSIN**



Department

Natural  
Resources

Madison, Wis.

1970

Dept. of Natural Resources  
Technical Library  
3911 Fish Hatchery Road  
Fitchburg, WI 53711 - 5397

By  
**Robert L. Hunt**

**LOAN COPY**



## ABSTRACT

The major findings, conclusions and management implications derived during 13 years of testing and evaluating experimental angling regulations applied to the fishery for wild brook trout (Salvelinus fontinalis) in Lawrence Creek are presented. Four general types of regulations were tested: size limits, bag limits, fish refuges and restrictions on fishing methods.

The size limit, if wisely applied, is the best single regulation for preventing excessive angler harvest of brook trout populations. A good "rule of thumb" is a size limit set at the length attained by 30% of the trout population in May, or 40% in June, or 50% in September (excluding age 0). This is 7 inches for the population in Lawrence Creek.

Bag limits provide no protection until the limit is reached, and since most of the catch of wild brook trout consists of 1 or 2 trout/trip, very restrictive bag limits would be needed to substantially reduce the catch. Angling effort appeared to be influenced by restrictions on the catch.

Angling regulations that allow some harvest or fishing opportunity every year from all fishable trout waters are clearly preferable to elimination of all fishing through creation of fish refuges.

Fishermen were attracted by "fly fishing only" regulations, and enjoyed high quality fishing, but there was little difference in other aspects of the fisheries in the flies-only zone versus the any-lure zone, and no detectable responses by the trout populations due to the flies-only restriction.

## ACKNOWLEDGMENTS

During the 13 years of conducting the trout research project at Lawrence Creek, many employees of the Wisconsin Department of Natural Resources contributed their skills and energies to the project. I acknowledge and thank them for their various contributions.

Special credit is due to James McFadden, the first project leader, who developed a solid foundation for subsequent research and reported many of the basic findings in his classic 1961 monograph, "A population study of the brook trout, Salvelinus fontinalis". John Brasch was mainly responsible for initiating the research project and for administering it during the first 3 years. Lyle Christenson continued this task during the last 10 years. Mr. Christenson and my fellow biologists then in the Cold Water Group, Oscar Brynildson, Ray White, and John Mason, also deserve special recognition for their professional advice and assistance in field work, data interpretations, and report preparations and revisions. Clarence Todd served capably and faithfully as project technician throughout the study. I also acknowledge the thousands of cooperative anglers who made their vital contributions to this project. Mrs. Elaine Sellin typed the various rough drafts and tables of this report.

This project was supported in part by funds from the Federal Aid to Wildlife Restoration Act, under Dingell-Johnson project F-83-R.

The author is a Fishery Biologist with the Bureau of Research, now stationed at Waupaca.

Edited by Ruth L. Hine

CONTENTS

|   | Page |
|---|------|
| INTRODUCTION . . . . .                            | 4    |
| SIZE LIMITS . . . . .                             | 6    |
| Summary of Significant Findings . . . . .         | 6    |
| Conclusions and Management Implications . . . . . | 9    |
| BAG LIMITS . . . . .                              | 9    |
| Summary of Significant Findings . . . . .         | 10   |
| Conclusions and Management Implications . . . . . | 10   |
| FISH REFUGES . . . . .                            | 12   |
| Summary of Significant Findings . . . . .         | 12   |
| Conclusions and Management Implications . . . . . | 13   |
| RESTRICTIONS ON FISHING METHODS . . . . .         | 14   |
| Summary of Significant Findings . . . . .         | 15   |
| Fly Fishing Only . . . . .                        | 15   |
| Catch and Release Fishing . . . . .               | 20   |
| Conclusions and Management Implications . . . . . | 21   |
| APPENDIX . . . . .                                | 23   |
| Tables 1-13 . . . . .                             | 23   |
| LITERATURE CITED . . . . .                        | 36   |

## INTRODUCTION

This report summarizes the major findings, conclusions and management implications derived during 13 years of testing and evaluating experimental angling regulations applied to the fishery for wild brook trout (Salvelinus fontinalis) in Lawrence Creek. Much of this information has been previously published (McFadden, 1956 and 1961; Hunt, Brynildson and McFadden, 1962; Hunt and Brynildson, 1964; and Hunt, 1964, 1966). The intent of this report is to consolidate the most useful information in these several papers, and to augment it with previously unpublished findings derived during the last 4 years of the study.

The contents is organized mainly under major headings descriptive of the 4 general types of fishing regulations tested: size limits, bag limits, restrictions on fishing methods, and fish refuges (prohibition of any fishing). Most of the supporting data are included in the Appendix rather than the body of the report, and information has also been drawn from the published reports cited above.

Lawrence Creek and our methods of investigation there have been described in the papers cited above and by White (1967) and Hunt (1969). In brief, the stream is 3.3 miles long, has an average width of 23 feet and a surface area of approximately 10 acres. It supports a dense population of wild brook trout and has a reputation for good fishing. Stream flow is quite stable within the year and from year to year. The water is moderately hard (total alkalinity approximately 160 ppm). Gradient is less than 10 feet/mile, and the bottom type is predominantly sand with the two major spawning areas concentrated in short stretches of the upper two study sections. Lawrence Creek is located centrally in Wisconsin but at the southern end of the present distribution of brook trout streams in the state. The growth rate of brook trout in Lawrence Creek probably exceeds the average for the state because of its southerly location and relatively fertile environment.

The stream was divided into 4 study sections (A-D) with no barriers between sections to prevent free movement of trout.

The two most important methods employed were precise estimates of the trout population with electrofishing gear before and after each fishing season and a compulsory, registration-type, creel census of all anglers. Each year age 0 brook trout captured during the electrofishing inventories were permanently marked with fin-clips. Within 3 years approximately 75% of the population consisted of known-age individuals, and thereafter the necessity to collect scale samples for aging was largely eliminated. Anglers obtained free daily permits at the creel census station near the stream. They could choose any section of stream open to fishing, but permits were issued for only one section/trip. All catches were presented to the census clerks for examination when the daily permits were returned.

Each year the opening and closing dates of the fishing season at Lawrence Creek coincided with the statewide dates, but at Lawrence Creek fishing was permitted only from 6:00 a.m. to 10:00 p.m. to facilitate operation of the census station. The various regulations tested and the years in which they were in effect were as follows:

- 1955:           6-inch size limit  
                daily bag of 10
  
- 1956-57:       no size limit  
                no bag limit  
                no fishing in section A
  
- 1958-60:       9-inch size limit  
                daily bag of 5  
                no fishing in section A
  
- 1961-67:       8-inch size limit  
                daily bag of 5  
                fly-fishing-only in sections C and D

Sport fishing for wild brook trout, the backbone of the trout fishery in Wisconsin, will not be maintained without an integrated and flexible management plan aimed at protecting both the environment and the trout populations. Our research at Lawrence Creek demonstrated that even a dense population of wild brook trout can be overharvested by sport fishing at levels of fishing effort that already exist in Wisconsin. Unfortunately, in the predator-prey relationship that exists between fishermen and brook trout there appear to be no natural controls effective enough to prevent such overharvest. In fact, the kind of mortality that angling inflicts upon brook trout populations is inversely density dependent. At any given level of angling effort, the rate of angling mortality increases as the density of the trout population decreases. Or, to put it another way, depletion of a sparse population requires proportionately less fishing effort than is required to deplete a dense population. Therefore, until such time as anglers voluntarily restrict their harvest to safe levels, it is essential that laws, in the form of angling regulations, be imposed to prevent overfishing. Such angling regulations might also be used to perpetuate and encourage the cultural qualities of sportsmanship traditionally associated with trout fishing, but their most basic prerequisite is the biological necessity of protecting an adequate number of spawners.

All of the commonly applied angling regulations tend to reduce the number of trout harvested, but the mere presence of fishing restrictions on the books does not assure they are doing the job intended. Excessive harvests occurred at Lawrence Creek two seasons and almost occurred another season even though there were regulations on the length of the fishing season, fishing methods and daily fishing hours, and on the size and number of fish that could be kept



during one of the 3 seasons. However, our research at Lawrence Creek also provided some useful facts for improving the application of trout fishing regulations to prevent overfishing, and as more information is obtained on the total trout resource of the state, these facts should become increasingly applicable.

#### SIZE LIMITS

As Allen and Cunningham (1957) stated in their discussion of trout fishing regulations in New Zealand, the direct effect of size limits is to cause anglers to return to the water trout that they might keep if it were legal to do so. Presumably, enough of these released undersized trout will survive long enough to contribute to the fishery by being caught later at a larger size or by adding to subsequent generations by spawning.

Three size limits were tested at Lawrence Creek (6, 8 and 9 inches) and during 2 successive seasons there was no minimum size limit. The 6-inch limit was the first one tested because it was the normal state-wide size limit. Our evaluation of this size limit provided a baseline for later reference to results obtained from testing other size limits that applied only at Lawrence Creek.

#### Summary of Significant Findings

(1) During the 2 seasons (1956-57) when there was no size limit in effect, few anglers kept trout less than 6 inches long. Consequently, the size distribution of the catch these 2 seasons was similar to that recorded when a 6-inch limit was in effect in 1955 (Fig. 1).

(2) During seasons when size limits of 8 or 9 inches were in effect, only 25% of the trout creeled were an inch or more larger than the prevailing size limit (Fig. 2).

(3) No age 0 (young-of-the-year) brook trout were harvested, even during the 2 seasons when no size limit applied (Table 1).

(4) Because most of the age I+ trout, which constituted the harvested portion of the population, were included in the size range from 6 to 10 inches, a change in the size limit of an inch or two greatly increased or decreased the percentage of trout that were legal-sized. For example, when the 1967 fishing season began, approximately 44% of the trout population consisted of 6- to 10-inch fish, and only 2% of the trout exceeded 10 inches. If the size limit had been set at 6, 7, 8 or 9 inches, the proportion of legal-sized trout would have been 46%, 30%, 16% or 5%, respectively.

(5) When the 6-inch limit prevailed, at least 1 legal trout was creeled on 47% of the angling trips (Table 2). When the size limit was increased to 8 inches, only 34% of the angling trips were "successful", and the proportion of successful trips dropped to 17%




FIGURE 1. Length frequency distributions of the catches of brook trout during the 1955-57 fishing seasons at Lawrence Creek.




FIGURE 2. Length frequency distributions of the catches of brook trout during the 1958-60 and 1961-67 fishing seasons at Lawrence Creek.

when the size limit was increased only 1 more inch (to 9 inches). When there was no size limit, the proportion of successful trips was similar to that recorded when the 6-inch limit applied. Catch/hour under the 6, 8, and 9-inch limits averaged 0.69, 0.34, and 0.13, respectively.

(6) During the two seasons (1956-57) when there was no size limit, the number of trout harvested represented 59% and 65% of the preseason population (Table 3). These rates of exploitation greatly exceeded the long-term equilibrium level that this brook trout population could sustain.

(7) During the seasons when there was an 8-inch limit, exploitation ranged from 6% to 12% and averaged 9%. When the 9-inch limit prevailed, exploitation was only 1% to 8%/season and averaged 4% (Table 3).

## Conclusions and Management Implications

(1) The size limit, if wisely applied, is the best single regulation for preventing excessive angler harvest of brook trout populations. The size limit applies to every trout caught, and it can be related to a rather stable biological parameter, growth rates of the trout populations.

(2) The present statewide size limit of 6 inches is too low for brook trout populations exhibiting growth rates similar to that of the Lawrence Creek population. Although the angler harvest was not excessive in Lawrence Creek for the season when a 6-inch limit was tested, excessive harvest could have occurred if more anglers had fished or if the trout population that year had not been as large as it was. Since few anglers chose to keep brook trout less than 6 inches long, the 6-inch limit was only a little better than no size limit at all.

(3) A "rule of thumb" that provides an approximation of a good size limit for brook trout is this: The size limit should be set at the length attained by 30% of the trout population in May, or 40% in June, or 50% in September (excluding age 0). For the brook trout population in Lawrence Creek, a 7-inch limit would be dictated by this rule of selection. However, before biologically sound regional size limits can be applied in Wisconsin, it will be necessary to substantially augment existing information on growth rates of brook trout populations. A concerted effort should be made soon to obtain such information from a statewide representation of streams.

### BAG LIMITS

In theory, daily bag limits set arbitrary upper limits on the catches of individual anglers regardless of their skill or angling effort. Presumably by limiting the catches of individuals, a cumulative effect of limiting the total harvest is also realized and more trout will survive to spawn or be caught at larger sizes than would be true if no bag limits existed.

Bag limits have also been proposed for such nonbiological reasons as distributing the total catch more evenly among anglers, as arbitrary goals for the anglers to attain ("I got my limit!"), and as a means to prevent fish-hogging and waste by encouraging anglers to fish for sport not numbers of fish.

At Lawrence Creek, bag limits of 5 to 10 per day were tested, and during two seasons (1956-57) there was no bag limit. These were the two seasons when there was also no minimum size limit. Consequently, anglers could keep all the trout they caught if they desired. The daily limit of 10, in effect in 1955, represented the normal statewide bag limit for trout.

## Summary of Significant Findings

(1) A bag of 10 trout was attained on only 50 (3%) of the angling trips in 1955, the season when this bag limit applied (Table 4).

(2) When there was no bag limit, 10 or more trout were creeled on 4% of the angling trips. Catches in excess of 10/trip increased the seasonal harvest by 5% in 1956 and 11% in 1957 (Table 4).

(3) Within the ranges of angling pressure and trout density observed at Lawrence Creek, theoretical reductions in the bag limit appeared to provide about the same proportional protection for sparse trout populations as for dense ones (Fig. 3). This type of relationship is not biologically desirable. Ideally, a bag limit regulation should provide proportionately more protection as the number of trout decreases.

(4) The daily bag of 5/day was tested in conjunction with rather high size limits (8 and 9 inches) for brook trout. Limit catches of trout over 9 inches were made on less than 1% of the angling trips and catches of 5 trout over 8 inches long were made on only 6% of the angling trips (Table 4).

(5) During all seasons and regardless of the bag limit allowed, most of the harvest was accounted for by catches of 1-3 trout/trip (Table 4).

(6) During all seasons and regardless of the liberality of the bag and size limits, more than 50% of the anglers failed to catch a single wild brook trout (Table 4).

(7) During all seasons most of the limit catches were taken by a few skillful anglers who fished several times each season at Lawrence Creek. Their catches/hour of effort were usually several times greater than those recorded by anglers who caught only 1 or 2 trout/trip. The difference between anglers who consistently caught several trout/trip and those who caught few or no trout/trip, was largely one of angling skill, not persistence, and angling skill became an increasingly important factor as the fishing regulations were made more restrictive (Table 5).

(8) Among the skillful anglers, the effect of a bag limit which they could reasonably attain may have been a stimulus to keep fishing until the limit was reached (Hobbs, 1948). More limit catches of 5 or 10 were made than catches of 1 or 2 less than the allowable limit (Table 4).

## Conclusions and Management Implications

Bag limits provide no protection until the limit is reached. In contrast to the size limit, the bag limit does not apply to every trout caught. Since most of the catch of wild brook trout is probably accounted for by anglers creeling 1 or 2 trout/trip, very




FIGURE 3. Theoretical effect of daily bag limits of less than 10 in reducing the catches of brook trout made during the 1955-57 fishing seasons at Lawrence Creek. Preseason trout population densities are listed in the legend.

restrictive bag limits would be needed to effect substantial reductions in the catch, and especially where bag limits are applied to broad geographical regions (statewide). For example, if angling effort remained unchanged, cutting the bag limit from 10 to 5 would probably reduce the total catch by only 20%, not 50%.

The experiments at Lawrence Creek were not designed to evaluate the psychological reactions of anglers to various bag limits, but circumstantial evidence suggested that angling effort was influenced by the restrictiveness of the bag limit. Even though few anglers were able to catch a limit of 10 trout, the opportunity to catch only 5 rather than 10/trip may have caused part of the decline in fishing pressure that occurred after the bag limit of 5 was adopted. If this hypothesis is valid, application of bag limits that differ over the course of a season, or that differ regionally within the state should be tested as a means to control one of the most important factors determining the size of the harvest -- the number of anglers fishing a given stream or region of streams.

### FISH REFUGES

Fish refuges have been proposed as a means to insure the survival of some spawning trout which, hopefully, will produce a surplus of progeny that will immigrate to the adjacent fishing water and bolster the depleted fish population there. During a 5-year period (1956-1960), the upper mile of Lawrence Creek was designated a fish refuge and an intensive effort was made to mark trout in this refuge zone and in the zone of fishing water below. All trout creeled were examined for markings, and information on the movement of trout into and out of the refuge was also obtained during each electrofishing census of the trout population. Information to evaluate the success of the refuge experiment was also available from the creel census the year prior to establishment of the refuge.

#### Summary of Significant Findings

(1) The brook trout population in the refuge increased substantially during the first 2 years of the experiment, but then it declined during the last 3 years to a level comparable to that present when the refuge was established (Fig. 4).

(2) Twice as many catchable-sized trout moved into the refuge as moved out (2,789 vs. 1,122). More young-of-the-year moved out than moved in, but the emigrants from the refuge increased the total number of trout in the fishing water by only 5% or less.

(3) The buildup and decline of the trout population in the refuge was largely due to changes in the survival rates of the resident population, not to gains or losses of trout into or out of the refuge (Table 6).




FIGURE 4. Standing crops of brook trout in the fish refuge of Lawrence Creek compared to those in the fishing water.

(4) Only 1% of the catch (45 of 4,695) consisted of trout that had emigrated from the refuge. During the fishing season when the refuge zone had been open to fishing, 21% of the total catch was made there. Most of the trout born in the refuge stayed there throughout their life.

Conclusions and Management Implications

As a means of providing better trout fishing, the mile-long head-water refuge was a failure. Many trout that could have been harvested or fished over were lost to natural mortality because they did not leave the refuge. The experiment did show that angling had been a critical factor limiting the density of the trout population in the refuge zone. However, despite the stockpiling response to elimination of all angling mortality, large-scale emigration did not occur.



Angling regulations that allow some harvest or fishing opportunity every year from all fishable trout waters are clearly preferable to elimination of all fishing through creation of fish refuges.

The possible use of fish refuges for periods of a year or 2 has not been explored in Wisconsin. In such instances, however, the emphasis would not be on "seeding" the adjacent fishing water but maximizing the opportunity of the resident trout population to take advantage of the growth potential of the state's more productive streams. This technique would probably have a rather restricted use, but it appears to be a procedure worth trying as a means of producing high quality fisheries and more efficient utilization of the inherent growth potential of trout populations. Better use of the growth potential in Wisconsin's more fertile trout streams could also be realized by alternately opening and closing upper and lower portions of streams on July 1st each year rather than allowing fishing all summer long. The bulk of the wild trout caught each season are yearlings or 2-year-olds. If such fish were periodically protected for the latter half of 1 summer and the first half of the following summer many more pounds of larger-than-average trout would be produced and eventually cropped.

#### RESTRICTIONS ON FISHING METHODS

Nearly all sport fishing in public waters is limited to some modification of hook and line gear. This in itself represents an often overlooked but radical limitation on the efficiency of capture as compared to the improvement that could be attained by employing electrofishing gear, chemical poisons or anesthetics, nets, seines, or explosives -- all of which are used legally by commercial fishermen or scientists. Any further restriction on the various methods of hook and line fishing, imposed either legally or voluntarily, will further tend to reduce the total catch of trout from a body of water regardless of what methods are eliminated. This is because over the course of a fishing season, the variable conditions of weather and water make each method more efficient on some days than all other methods (Allen and Cunningham, 1957). Moreover, most anglers are more proficient at 1 method of fishing than several methods. If their favored method is prohibited, they must then choose to fish elsewhere or fish less proficiently.

Recently, as the number of trout fishermen has tended to increase and fishing quality has tended to decrease, 2 fishing restrictions are receiving increasing emphasis. One involves the concept of releasing all or nearly all of the trout caught. This type of trout fishing has been labelled "fishing for fun" by many anglers and authors, but as Stroud (1964) has convincingly demonstrated, "catch and release fishing" is a much more appropriate designation because most fishing is done for recreation not economic gain. The second concept receiving increasing emphasis is prohibition of bait fishing and the attendant high mortality it inflicts on trout that are hooked and released. However, even the

high mortality rate inflicted on undersize trout hooked and released by bait fishermen could be substantially reduced if such anglers could be educated to cut their lines and release undersize, deeply hooked trout with the hooks still in them. As Mason and Hunt (1967) demonstrated, approximately 2/3 of such released trout would probably survive and continue to grow.

Some trout water has now been restricted to "fly fishing only" or "artificial lures only" in at least 25 states, but such restrictions are not widespread in any state (Wallis, 1963). Wherever the emphasis is on "catch and release fishing", bait fishing is also prohibited, but in many "fly fishing only" waters substantial harvests are permitted.

At Lawrence Creek fly fishing was the only method permitted in the lower 2 (the "flies-only zone") of the 4 study sections during 7 successive seasons (1961-67). In the other 2 sections (the "any-lure zone"), all of the commonly accepted methods of trout fishing, including fly fishing, were allowed. Thus, fly fishermen could choose to fish in any of the lower sections, but anglers using spinning lures or fishing with bait could choose only 1 of the upper 2 sections. As in other years, all anglers were required to pick up a daily permit at the creel census station near the stream, and only 1 section could be fished/trip.

"Catch and release fishing" was not specifically tested, but during the 3 seasons when a 9-inch size limit was being tested, the fishing approached a catch and release condition. Only 6% of the trout caught were legal-sized.

### Summary of Significant Findings

#### Fly Fishing Only:

(1) Under the conditions of fishing pressure, catch, and trout densities that prevailed at Lawrence Creek, fly fishing had no uniquely beneficial biological effects that could be detected. Changes in standing crops, survival rates, reproduction, and growth of the trout populations in the 2 fishing zones appeared to be independent of the methods of angler harvest (Tables 7-9).

(2) In both zones hooking mortality inflicted on released trout was apparently compensated for by a decrease in mortality due to other causes. Summer mortality unaccounted for by trout harvested was not consistently higher in the any-lure zone under the prevailing conditions of hooking and releasing up to 50% of the trout population (Tables 8-10).

(3) Angling exploitation was similar in both fishing zones, averaging about 10% of the preseason population and ranging from 7-11% in the any-lure zone and 5-12% in the flies-only zone (Table 10).

(4) In both fishing zones, cropping tended to exceed recruitment during the first half of the season, but during the second half recruitment predominated. By the end of the fishing season, there tended to be more legal trout remaining than were harvested and more than were present at the beginning of the season (Table 11).

(5) In both fishing zones, the number of legal and sublegal (6-8 inches) trout tended to increase during the 7-year period, but the ratio of sublegal trout to legal trout in the flies-only zone versus the any-lure zone did not increase. The assumed lower hooking mortality of released trout in the flies-only zone did not detectably benefit the number of sublegal trout escaping all causes of mortality (Table 11 and Figs. 5 and 6).

(6) Improved standing crops of trout in the 2 fishing zones reflected better-than-average rates of overwinter survival and relatively low angler exploitation rather than differences in the methods of fishing employed (Tables 10 and 12).

(7) Fly fishermen were definitely attracted to Lawrence Creek by the fly-fishing-only restriction. More fly fishermen were registered in 1967 than in any previous year despite the fact that the total number of anglers in 1967 was 30% lower than the record high total registered in 1955. During 6 fishing seasons prior to the adoption of this regulation, fly fishermen accounted for 16-28% of the angling trips/season and 14-22% of the seasonal catches. After its adoption, they accounted for 39-55% of the trips/season and 40-66% of the seasonal catches (Table 13).

(8) Nearly all of the fly fishermen chose to fish in the flies-only zone. However, despite the increase in fly fishermen and their concentration in the lower half of the stream, this portion received less fishing pressure than it did when all types of angling were permitted there.

(9) Angling effort was always greater in the any-lure zone and the number of trout creelcd there was greater during 6 of the 7 seasons. However, the catch/hour of legal trout was always higher in the flies-only zone and the catch/hour of trout released was also higher in the flies-only zone during 5 of the 7 seasons. Catches of legal trout averaged 0.29 and 0.41/hour in the any-lure zone and flies-only zone, respectively, and catches of trout that were released averaged 1.43 and 1.66/hour in the two zones over the 7-year period (Table 10 and Figures 7-8).

(10) The average length of trout creelcd/season in the flies-only zone was always slightly larger and averaged 8.8 inches versus 8.6 inches in the any-lure zone for the 7-year period. Approximately 4.9 trout were released/trout creelcd in the any-lure zone. In the flies-only zone, the ratio was 4.0 to 1 (Table 10).




FIGURE 5. Number of legal and sublegal (6- to 8-inch) brook trout in the any-lure and flies-only zones of Lawrence Creek at the beginning of the 1961-67 fishing seasons.




FIGURE 6. Number of legal and sublegal (6- to 8-inch) brook trout in the any-lure and flies-only zones of Lawrence Creek at the end of the 1961-67 fishing seasons.




FIGURE 7. Angling effort and catch in the any-lure and flies-only zones of Lawrence Creek during the 1961-67 fishing seasons.




FIGURE 8. Catch/hour of brook trout creeled or released in the any-lure and flies-only zones of Lawrence Creek during the 1961-67 fishing seasons.

(11) Less than 1% of the fly fishermen reported releasing all legal-sized trout caught.

Catch and Release Fishing:

(12) During the 1958-60 fishing seasons, when a high size limit was in effect, the angler harvests were equivalent to only 4%, 8%, and 1% of the preseason trout populations. These compare to exploitation rates of 32%, 59%, and 65% during the preceding 3 seasons when regulations were very liberal (Table 3).

(13) During the 1958-60 seasons, an average of only 6% of the trout caught were legal-sized. Anglers released approximately 15 trout for every trout creeled.

(14) Despite an average catch-rate of 1-3 sublegal trout/hour on an easily fished, aesthetic stream, the reaction of anglers was generally negative during the 1958-60 seasons. Angling pressure declined from the 400-500 hours/acre characteristic of the 1955-57 seasons to only 250, 215, and 137 hours/acre during 1958-60, respectively. Catches/hour of legal trout also declined radically, from the averages of 0.69, 0.47, and 0.89 trout/hour during 1955-57 to 0.13, 0.16, and 0.09 trout/hour during the 1958-60 seasons (Table 2).

(15) During the entire 1960 season only 85 brook trout were cropped as compared to a catch of 512 on the opening day of the 1955 season when the normal statewide size and bag limit regulations prevailed. The total catch that year was 3,040.

### Conclusions and Management Implications

Designation of part of Lawrence Creek for "fly fishing only" proved to be popular among fly fishermen. They were attracted by the regulation, and they enjoyed high quality fishing for wild brook trout without abusing the trout population. However, except for the fishing methods, there was little difference in other aspects of the fisheries in the flies-only zone versus the any-lure zone, and there were no detectable responses by the trout populations that could be attributed to the presence or absence of the flies-only restriction. In both zones the amount of fishing pressure and the restrictive impact of the 8-inch size limit and bag limit of 5 overshadowed effects on the harvests and residual populations related to fishing methods. The only difference associated with fishing methods that may have had some impact on the fisheries in the two fishing zones was the consistently higher catch/hour by fly fishermen versus all anglers using the any-lure zone. The greater success rate in the flies-only zone may have been only a reflection of the slightly greater number of legal trout normally present there or the possible attraction of better-than-average fly fishermen to Lawrence Creek in response to the flies-only regulation.

If such a regulation does attract expert fly fishermen, and if nearly all of them keep the legal trout they catch, as they did at Lawrence Creek, the imposition of a flies-only regulation may not, of itself, prevent overexploitation.

At the present time in Wisconsin, this regulation appears to be more practical for managing trout fishermen than trout populations. Its application would appear to be biologically valid only in conjunction with an intensive catch and release fishery, a fishery where nearly every trout is likely to be caught one or more times. Where the emphasis is on a substantial harvest, there is little biological justification at present for limiting this harvest to fly fishermen only.

The general reaction of anglers to catching and releasing large numbers of 6-8 inch brook trout was definitely negative. Apparently, if such recreation is to be favorably accepted in Wisconsin, the



average size of trout released must be considerably larger. In Wisconsin, this type of fishery would, therefore, be limited to a few of the most productive lakes and streams containing fast-growing brown or rainbow trout. If an emphasis is placed on "catch and release fishing" on a season-long basis, it must be recognized that very little of the trout flesh produced will be cropped. Since natural mortality and angling mortality tend to compete, very few of the trout hooked and released unharmed will live long enough to reach trophy size. A large part of the harvestable surplus of trout flesh produced will be lost. This voluntary loss must be balanced against the additional sport provided by the greater number of larger-than-average trout such specially designated waters can maintain for a longer period each fishing season.

Even though many states now have some form of special regulations designed to emphasize the unharmed release of all or nearly all of the trout captured, there has been little biological evaluation of the validity of continuing to apply such restrictions (personal communication with 18 fish and game agencies). Moreover, among the studies that have been conducted, it is difficult to find consistent conclusions. Unfortunately, in many instances where fly-fishing-only rules have been adopted to protect released under-sized trout, higher size limits and lower bag limits have also been instituted simultaneously. In such situations it is impossible to relate any beneficial responses by trout populations solely to the flies-only rules. The most notable exception to this type of multiple change of fishing regulations is the excellent study reported by Shetter and Alexander (1962). In this study the normal statewide regulations of a 7-inch size limit and daily bag of 10 applied to all 4 study sections of Hunt Creek, but in 2 of the sections only fly fishing was permitted during a 5-year period. In Hunt Creek, as in Lawrence Creek, the flies-only rule influenced angling pressure and choice of fishing zones, but it caused no biologically important responses by the trout populations.

Perhaps the most valid judgement at this stage is that each proposed application of such special regulations must be carefully scrutinized on its own merits, taking into account both its biological and social implications. Furthermore, because of the varied success that has resulted to date, additional application of such special regulations should not be initiated without a definite plan to study and evaluate the consequences.

## APPENDIX

TABLE 1

Percentage of the Total Number and Total Weight of the Yield  
of Brook Trout Accounted for by Various Age Groups During  
the 1955-67 Fishing Seasons at Lawrence Creek

| Year | Age Group |        |        |        |        |        |        |        | Size<br>Limit |
|------|-----------|--------|--------|--------|--------|--------|--------|--------|---------------|
|      | I         |        | II     |        | III    |        | IV+    |        |               |
|      | Number    | Weight | Number | Weight | Number | Weight | Number | Weight |               |
| 1955 | 62.5      | 44.4   | 35.5   | 49.0   | 1.9    | 5.9    | 0.1    | 0.7    | 6 inches      |
| 1956 | 47.9      | 33.5   | 50.0   | 59.9   | 1.9    | 5.2    | 0.2    | 1.4    |               |
| 1957 | 89.9      | 82.6   | 8.2    | 13.3   | 1.3    | 3.6    | 0.1    | 0.5    | None          |
| 1958 | 7.2       | 6.2    | 82.0   | 76.7   | 9.9    | 13.7   | 0.9    | 3.4    |               |
| 1959 | 4.5       | 3.8    | 79.8   | 81.1   | 15.2   | 14.6   | 0.5    | 0.5    | 9 inches      |
| 1960 | 8.2       | 7.0    | 27.0   | 23.7   | 64.8   | 69.3   | 0.0    | 0.0    |               |
| 1961 | 24.2      | 19.7   | 74.4   | 77.7   | 0.7    | 1.2    | 0.7    | 1.4    | 8 inches      |
| 1962 | 14.8      | 13.0   | 74.8   | 71.0   | 10.4   | 16.0   | --     | --     |               |
| 1963 | 3.5       | 2.9    | 86.3   | 85.3   | 8.8    | 9.3    | 1.4    | 2.5    |               |
| 1964 | 7.5       | 5.9    | 66.1   | 62.1   | 24.8   | 28.5   | 1.6    | 3.5    |               |
| 1965 | 7.2       | 5.9    | 71.8   | 69.1   | 15.6   | 16.9   | 5.4    | 8.1    |               |
| 1966 | 8.8       | 7.6    | 70.5   | 68.2   | 18.9   | 21.3   | 1.7    | 2.9    |               |
| 1967 | 2.2       | 2.1    | 76.8   | 74.2   | 19.2   | 21.1   | 1.8    | 2.6    |               |

TABLE 2

## Indices of Angling Quality for the 1955-67 Fishing Seasons at Lawrence Creek

| Year         | Size Limit<br>(Inches) | Yield  |        | Angling<br>Effort |       | No. of<br>Trout<br>Released | Catch/Hour |          | Percent<br>"Successful"<br>Trips | Trout Released/<br>Trout Creeled | Size<br>Limit |
|--------------|------------------------|--------|--------|-------------------|-------|-----------------------------|------------|----------|----------------------------------|----------------------------------|---------------|
|              |                        | Number | Pounds | Trips             | Hours |                             | Creeled    | Released |                                  |                                  |               |
| 1955         | 6                      | 3,040  | 537    | 1,712             | 4,653 | 2,435                       | 0.69       | 0.52     | 47                               | 0.8                              | 6 inches      |
| 1956         | 0                      | 1,360  | 220    | 1,313             | 3,051 | 376                         | 0.47       | 0.13     | 34                               | 0.3                              | None          |
| 1957         | 0                      | 2,778  | 431    | 1,249             | 3,168 | 685                         | 0.89       | 0.22     | 44                               | 0.2                              |               |
| 1958         | 9                      | 223    | 68     | 758               | 1,847 | 3,704                       | 0.13       | 2.00     | 19                               | 16.6                             | 9 inches      |
| 1959         | 9                      | 243    | 77     | 728               | 1,560 | 1,846                       | 0.16       | 1.18     | 19                               | 7.6                              |               |
| 1960         | 9                      | 85     | 27     | 434               | 1,007 | 2,572                       | 0.09       | 2.55     | 12                               | 30.2                             |               |
| 1961         | 8                      | 442    | 107    | 592               | 1,279 | 1,593                       | 0.35       | 1.25     | 32                               | 3.6                              | 8 inches      |
| 1962         | 8                      | 540    | 131    | 896               | 2,173 | 4,143                       | 0.25       | 1.91     | 27                               | 7.7                              |               |
| 1963         | 8                      | 752    | 169    | 874               | 2,324 | 2,551                       | 0.33       | 1.10     | 36                               | 3.4                              |               |
| 1964         | 8                      | 625    | 149    | 881               | 2,141 | 4,097                       | 0.30       | 1.91     | 30                               | 6.6                              |               |
| 1965         | 8                      | 671    | 157    | 858               | 2,122 | 3,243                       | 0.32       | 1.53     | 33                               | 4.8                              |               |
| 1966         | 8                      | 862    | 203    | 826               | 2,087 | 2,950                       | 0.41       | 1.41     | 41                               | 3.4                              |               |
| 1967         | 8                      | 1,111  | 266    | 1,178             | 2,784 | 3,565                       | 0.40       | 1.28     | 37                               | 3.2                              |               |
| Avg. 1956-57 |                        | 2,069  | 326    | 1,281             | 3,110 | 530                         | 0.68       | 0.18     | 39                               | 0.3                              | None          |
|              | 1958-60                | 184    | 57     | 640               | 1,471 | 2,707                       | 0.13       | 1.06     | 17                               | 18.1                             | 9 inches      |
|              | 1961-67                | 715    | 169    | 872               | 2,130 | 3,163                       | 0.34       | 1.48     | 34                               | 4.7                              | 8 inches      |

TABLE 3

Percentage of the Preseason Population of Brook Trout  
Harvested According to Age Group During  
the 1955-67 Fishing Seasons at Lawrence Creek

| Year         | Age Group |       |      | Stream Total<br>for All Adult<br>Trout (I+) | Size<br>Limit |
|--------------|-----------|-------|------|---|---------------|
|              | I         | II    | III+ |   |               |
| 1955         | 24.4      | 71.7  | 42.2 | 32.2  | 6 inches      |
| 1956         | 45.6      | 85.2  | 35.4 | 59.0  | None          |
| 1957         | 62.0      | 100.0 | 72.2 | 64.6  |               |
| 1958         | 0.3       | 19.8  | 42.8 | 3.9   | 9 inches      |
| 1959         | 1.0       | 11.2  | 30.6 | 8.2   |               |
| 1960         | 0.1       | 12.8  | 29.1 | 1.2   |               |
| 1961         | 3.0       | 39.8  | 40.0 | 9.9   | 8 inches      |
| 1962         | 0.9       | 36.3  | 51.9 | 5.5   |               |
| 1963         | 0.6       | 26.9  | 58.8 | 10.5  |               |
| 1964         | 0.6       | 32.5  | 41.2 | 6.8   |               |
| 1965         | 1.0       | 25.5  | 57.6 | 9.4   |               |
| 1966         | 1.1       | 34.5  | 46.4 | 9.6   |               |
| 1967         | 0.5       | 28.8  | 52.1 | 11.9  |               |
| Avg. 1956-57 | 57.7      | 91.7  | 50.0 | 62.7  | None          |
| 1958-60      | 0.6       | 14.1  | 31.7 | 3.5   | 9 inches      |
| 1961-67      | 1.1       | 32.0  | 49.7 | 9.1   | 8 inches      |

TABLE 4

Frequency of Various Catches of Brook Trout Per Trip  
During the 1955-67 Fishing Seasons at Lawrence Creek \*

| Catch/<br>Trip | Fishing Season          |      |      |      |      |      |      |      |      |      |      |      |      |  |
|----------------|-------------------------|------|------|------|------|------|------|------|------|------|------|------|------|--|
|                | 1955                    | 1956 | 1957 | 1958 | 1959 | 1960 | 1961 | 1962 | 1963 | 1964 | 1965 | 1966 | 1967 |  |
|                | Number of Angling Trips |      |      |      |      |      |      |      |      |      |      |      |      |  |
| 0              | 912                     | 882  | 683  | 614  | 588  | 383  | 401  | 646  | 566  | 617  | 578  | 484  | 717  |  |
| 1              | 188                     | 153  | 136  | 88   | 80   | 32   | 78   | 114  | 130  | 103  | 110  | 131  | 168  |  |
| 2              | 156                     | 96   | 91   | 31   | 34   | 7    | 45   | 50   | 61   | 63   | 64   | 65   | 90   |  |
| 3              | 103                     | 55   | 58   | 12   | 9    | 6    | 25   | 36   | 43   | 35   | 35   | 49   | 52   |  |
| 4              | 94                      | 37   | 50   | 3    | 10   | 3    | 16   | 16   | 19   | 23   | 27   | 31   | 68   |  |
| 5              | 63                      | 32   | 44   | 5    | 7    | 3    | 27   | 34   | 55   | 40   | 44   | 66   | 83   |  |
| 6              | 58                      | 15   | 28   |      |      |      |      |      |      |      |      |      |      |  |
| 7              | 33                      | 8    | 31   |      |      |      |      |      |      |      |      |      |      |  |
| 8              | 24                      | 5    | 22   |      |      |      |      |      |      |      |      |      |      |  |
| 9              | 31                      | 8    | 18   |      |      |      |      |      |      |      |      |      |      |  |
| 10             | 50                      | 8    | 16   |      |      |      |      |      |      |      |      |      |      |  |
| 11+            |                         | 14   | 72   |      |      |      |      |      |      |      |      |      |      |  |
| Total<br>Trips | 1712                    | 1313 | 1249 | 758  | 728  | 434  | 592  | 896  | 874  | 881  | 858  | 826  | 1178 |  |

\*1955: Bag limit of 10  
1956-57: No bag limit  
1958-67: Bag limit of 5

TABLE 5

Distribution of Trips and Catch Among Anglers  
Making Various Numbers of Trips During  
the 1955-64 Fishing Seasons at Lawrence Creek

| Year | Trips/<br>Season | Number of<br>Anglers | Percentage<br>of Anglers | Percentage<br>of Total<br>Trips | Percentage<br>of Total<br>Catch | Catch/<br>Hour | Bag<br>Limit |
|------|------------------|----------------------|--------------------------|---------------------------------|---------------------------------|----------------|--------------|
| 1955 | 1-2              | 881                  | 87.8                     | 58.8                            | 31.4                            | 0.34           | 10           |
|      | 3-4              | 73                   | 7.3                      | 14.5                            | 18.9                            | 0.77           |              |
|      | 5-9              | 32                   | 3.2                      | 12.2                            | 20.5                            | 1.16           |              |
|      | 10+              | 17                   | 1.7                      | 14.5                            | 29.2                            | 1.36           |              |
| 1956 | 1-2              | 936                  | 93.9                     | 78.9                            | 46.9                            | 0.27           | None         |
|      | 3-4              | 45                   | 4.5                      | 11.1                            | 21.9                            | 0.80           |              |
|      | 5-9              | 12                   | 1.2                      | 5.8                             | 18.8                            | 1.54           |              |
|      | 10+              | 4                    | 0.4                      | 4.2                             | 12.4                            | 1.35           |              |
| 1957 | 1-2              | 711                  | 91.0                     | 65.6                            | 36.9                            | 0.55           |              |
|      | 3-4              | 45                   | 5.8                      | 12.3                            | 16.3                            | 1.06           |              |
|      | 5-9              | 14                   | 1.8                      | 6.5                             | 15.0                            | 2.15           |              |
|      | 10+              | 11                   | 1.4                      | 15.6                            | 31.8                            | 1.86           |              |
| 1958 | 1-2              | 368                  | 85.8                     | 57.8                            | 33.7                            | 0.08           |              |
|      | 3-4              | 38                   | 8.8                      | 16.6                            | 15.2                            | 0.10           |              |
|      | 5-9              | 17                   | 4.0                      | 13.7                            | 16.6                            | 0.16           |              |
|      | 10+              | 6                    | 1.4                      | 11.9                            | 34.5                            | 0.24           |              |
| 1959 | 1-2              | 355                  | 87.9                     | 56.8                            | 24.3                            | 0.06           |              |
|      | 3-4              | 26                   | 6.4                      | 12.2                            | 7.7                             | 0.10           |              |
|      | 5-9              | 16                   | 4.0                      | 15.2                            | 26.3                            | 0.26           |              |
|      | 10+              | 7                    | 1.7                      | 15.8                            | 41.7                            | 0.38           |              |
| 1960 | 1-2              | 297                  | 93.4                     | 77.0                            | 42.9                            | 0.04           |              |
|      | 3-4              | 12                   | 3.8                      | 9.0                             | 6.6                             | 0.06           |              |
|      | 5-9              | 8                    | 2.5                      | 11.5                            | 31.9                            | 0.23           |              |
|      | 10+              | 1                    | 0.3                      | 2.5                             | 18.7                            | 0.59           |              |
| 1961 | 1-2              | 306                  | 89.7                     | 59.1                            | 20.5                            | 0.11           | 5            |
|      | 3-4              | 22                   | 6.5                      | 3.4                             | 15.1                            | 0.36           |              |
|      | 5-9              | 6                    | 1.8                      | 7.9                             | 13.1                            | 0.86           |              |
|      | 10+              | 7                    | 2.0                      | 21.0                            | 51.3                            | 0.91           |              |
| 1962 | 1-2              | 464                  | 89.0                     | 62.3                            | 28.6                            | 0.11           |              |
|      | 3-4              | 37                   | 7.1                      | 14.0                            | 16.9                            | 0.26           |              |
|      | 5-9              | 12                   | 2.3                      | 9.7                             | 17.6                            | 0.54           |              |
|      | 10+              | 8                    | 1.6                      | 14.0                            | 36.9                            | 0.72           |              |
| 1963 | 1-2              | 456                  | 89.6                     | 65.2                            | 40.0                            | 0.21           |              |
|      | 3-4              | 33                   | 6.5                      | 12.7                            | 9.7                             | 0.21           |              |
|      | 5-9              | 12                   | 2.3                      | 8.8                             | 13.1                            | 0.47           |              |
|      | 10+              | 8                    | 1.6                      | 13.3                            | 37.2                            | 0.89           |              |
| 1964 | 1-2              | 433                  | 86.9                     | 60.2                            | 39.5                            | 0.20           |              |
|      | 3-4              | 40                   | 8.0                      | 15.4                            | 15.5                            | 0.29           |              |
|      | 5-9              | 18                   | 3.6                      | 13.7                            | 20.7                            | 0.42           |              |
|      | 10+              | 7                    | 1.5                      | 10.6                            | 24.3                            | 0.68           |              |

TABLE 6

Percentage Mortality and Survival of Age I+ Brook Trout  
in the Refuge Zone Versus the Fishing Water Zone of Lawrence Creek

| Year<br>(April-Sept.) | "Refuge" Zone        |                      |          | "Fishing Water" Zone |                      |          |
|-----------------------|----------------------|----------------------|----------|----------------------|----------------------|----------|
|                       | Natural<br>Mortality | Angling<br>Mortality | Survival | Natural<br>Mortality | Angling<br>Mortality | Survival |
| 1955*                 | 32                   | 12                   | 56       | 27                   | 39                   | 34       |
| 1956                  | 39                   |                      | 61       | 24                   | 59                   | 17       |
| 1957                  | 33                   |                      | 67       | 2                    | 64                   | 34       |
| 1958                  | 48                   |                      | 52       | 47                   | 4                    | 49       |
| 1959                  | 70                   |                      | 30       | 50                   | 8                    | 42       |
| 1960                  | 79                   |                      | 21       | 57                   | 1                    | 42       |

\* The refuge was established after the 1955 fishing season ended. Angling mortality, natural mortality and survival rates for 1955 are included for comparative purposes when both zones were open to fishing.

TABLE 7

Standing Crops of Brook Trout in the Any-Lure Zone  
and Flies-Only Zone of Lawrence Creek During 1961-67\*

|      | Number of Trout in April |       |     |       |                 |       |     |       |
|------|--------------------------|-------|-----|-------|-----------------|-------|-----|-------|
|      | Any-Lure Zone            |       |     |       | Flies-Only Zone |       |     |       |
|      | I                        | II    | III | Σ I+  | I               | II    | III | Σ I+  |
| 1961 | 2,026                    | 220   | 4   | 2,250 | 1,576           | 607   | 9   | 2,232 |
| 1962 | 4,073                    | 467   | 22  | 4,562 | 4,494           | 646   | 86  | 5,226 |
| 1963 | 2,657                    | 893   | 34  | 3,584 | 1,987           | 1,516 | 97  | 3,600 |
| 1964 | 4,330                    | 610   | 128 | 5,068 | 3,195           | 659   | 272 | 4,090 |
| 1965 | 3,380                    | 1,110 | 94  | 4,584 | 1,623           | 779   | 151 | 2,553 |
| 1966 | 4,548                    | 985   | 183 | 5,721 | 2,367           | 779   | 172 | 3,342 |
| 1967 | 3,660                    | 1,588 | 180 | 5,445 | 2,272           | 1,372 | 227 | 3,889 |

Number of Trout in September

|      | 0     | I     | II  | III | Σ 0+   | 0     | I     | II  | III | Σ 0+  |
|------|-------|-------|-----|-----|--------|-------|-------|-----|-----|-------|
| 1961 | 9,375 | 1,421 | 93  | 2   | 10,891 | 4,938 | 939   | 168 | 1   | 6,046 |
| 1962 | 4,382 | 2,186 | 97  | 5   | 6,670  | 3,229 | 2,337 | 106 | 17  | 5,689 |
| 1963 | 5,753 | 1,256 | 278 | 11  | 7,298  | 4,614 | 1,132 | 472 | 26  | 6,249 |
| 1964 | 6,974 | 2,576 | 198 | 42  | 9,790  | 2,706 | 1,806 | 247 | 82  | 4,848 |
| 1965 | 5,779 | 1,683 | 248 | 29  | 7,741  | 2,673 | 1,455 | 239 | 37  | 4,408 |
| 1966 | 5,910 | 2,614 | 343 | 33  | 8,901  | 2,282 | 1,501 | 357 | 56  | 4,202 |
| 1967 | 6,158 | 1,642 | 424 | 30  | 8,256  | 1,737 | 1,520 | 442 | 98  | 3,810 |

\* Any-lure zone = 1.86 miles and flies-only zone = 1.44 miles



TABLE 8

Percentage Mortality and Survival of Age I and II Brook Trout  
in the Any-Lure Zone and Flies-Only Zone of Lawrence Creek During 1961-67

|        |         | Any-Lure Zone |           |          | Flies-Only Zone |           |          |
|--------|---------|---------------|-----------|----------|-----------------|-----------|----------|
|        |         | Angling*      | Natural   | Survival | Angling*        | Natural   | Survival |
|        |         | Mortality     | Mortality |          | Mortality       | Mortality |          |
| Age I  | 1961    | 3             | 27        | 70       | 2               | 37        | 61       |
|        | 1962    | 1             | 45        | 54       | 1               | 47        | 52       |
|        | 1963    | 1             | 45        | 54       | 1               | 43        | 56       |
|        | 1964    | 1             | 40        | 59       | 1               | 42        | 57       |
|        | 1965    | --            | --        | --       | --              | --        | --       |
|        | 1966    | 1             | 42        | 56       | 2               | 35        | 63       |
|        | 1967    | 1             | 55        | 44       | 1               | 33        | 66       |
| Avg.   | 1961-67 | 1             | 42        | 57       | 1               | 40        | 59       |
| Age II | 1961    | 54            | 4         | 42       | 35              | 37        | 28       |
|        | 1962    | 51            | 28        | 21       | 25              | 59        | 16       |
|        | 1963    | 37            | 29        | 34       | 23              | 29        | 48       |
|        | 1964    | 43            | 25        | 32       | 23              | 40        | 37       |
|        | 1965    | 27            | 51        | 22       | 23              | 46        | 31       |
|        | 1966    | 38            | 27        | 35       | 30              | 24        | 46       |
|        | 1967    | 30            | 44        | 26       | 20              | 40        | 32       |
| Avg.   | 1961-67 | 40            | 30        | 30       | 27              | 39        | 34       |

\* Mortality accounted for by the angler harvest only

TABLE 9

Average Lengths in September of Ages 0-II Brook Trout  
in the Any-Lure Zone and Flies-Only Zone  
of Lawrence Creek During 1958-67

| Year *       | Average Total Length in Inches |     |     |                 |     |     |
|--------------|--------------------------------|-----|-----|-----------------|-----|-----|
|              | Any-Lure Zone                  |     |     | Flies-Only Zone |     |     |
|              | 0                              | I   | II  | 0               | I   | II  |
| 1958         | 3.6                            | 6.7 | 8.3 | 3.7             | 7.0 | 9.5 |
| 1959         | 3.7                            | 7.1 | 8.5 | 3.8             | 6.8 | 7.7 |
| 1960         | 3.9                            | 7.1 | 8.7 | 3.8             | 7.2 | 9.3 |
| 1961         | 4.0                            | 7.5 | 9.1 | 3.9             | 7.4 | 9.2 |
| 1962         | 3.9                            | 6.9 | 8.4 | 3.7             | 6.9 | 8.7 |
| 1963         | 4.0                            | 7.0 | 8.6 | 4.1             | 6.9 | 8.3 |
| 1964         | 3.7                            | 6.7 | 8.3 | 3.8             | 6.8 | 8.2 |
| 1965         | 4.0                            | 7.1 | 8.2 | 3.9             | 6.9 | 8.5 |
| 1966         | 4.1                            | 6.9 | 8.2 | 4.0             | 7.2 | 8.6 |
| 1967         | 4.0                            | 6.7 | 8.2 | 3.9             | 6.7 | 8.5 |
| Avg. 1958-60 | 3.7                            | 7.0 | 8.5 | 3.8             | 7.0 | 8.8 |
| 1961-67      | 3.9                            | 7.0 | 8.4 | 3.9             | 7.0 | 8.7 |

\* The flies-only restriction was in effect during the 1961-67 fishing seasons. Age-specific growth data for 1958-60 are presented for comparative purposes.

TABLE 10

Creel Census Statistics for the Any-Lure and Flies-Only Zones  
of Lawrence Creek for the 1961-67 Fishing Seasons\*

| Item                     | 1961 | 1962  | 1963  | 1964  | 1965  | 1966  | 1967  | Avg.  |
|--------------------------|------|-------|-------|-------|-------|-------|-------|-------|
| Angling Trips:           |      |       |       |       |       |       |       |       |
| Any-Lure Zone            | 310  | 499   | 481   | 560   | 569   | 551   | 771   | 534   |
| Flies-Only Zone          | 282  | 397   | 393   | 321   | 289   | 275   | 407   | 338   |
| Angling Hours:           |      |       |       |       |       |       |       |       |
| Any-Lure Zone            | 698  | 1,262 | 1,305 | 1,390 | 1,426 | 1,455 | 1,868 | 1,343 |
| Flies-Only Zone          | 581  | 911   | 1,019 | 751   | 696   | 633   | 916   | 787   |
| Trout Creeled - Number:  |      |       |       |       |       |       |       |       |
| Any-Lure Zone            | 187  | 300   | 340   | 364   | 420   | 512   | 614   | 392   |
| Flies-Only Zone          | 255  | 240   | 405   | 262   | 251   | 350   | 497   | 323   |
| Trout Creeled - Pounds:  |      |       |       |       |       |       |       |       |
| Any-Lure Zone            | 19   | 68    | 76    | 84    | 94    | 117   | 139   | 85    |
| Flies-Only Zone          | 28   | 63    | 94    | 65    | 63    | 86    | 127   | 75    |
| Trout Released - Number: |      |       |       |       |       |       |       |       |
| Any-Lure Zone            | 658  | 2,022 | 1,326 | 2,461 | 2,096 | 1,959 | 2,466 | 1,855 |
| Flies-Only Zone          | 933  | 2,121 | 1,225 | 1,636 | 1,147 | 991   | 1,099 | 1,307 |
| Catch/Hour - Legal:      |      |       |       |       |       |       |       |       |
| Any-Lure Zone            | .27  | .24   | .27   | .26   | .29   | .35   | .33   | .29   |
| Flies-Only Zone          | .44  | .26   | .40   | .35   | .36   | .55   | .54   | .41   |
| Catch/Hour - Released:   |      |       |       |       |       |       |       |       |
| Any-Lure Zone            | .94  | 1.60  | 1.38  | 1.77  | 1.47  | 1.35  | 1.32  | 1.43  |
| Flies-Only Zone          | 1.60 | 2.32  | 1.20  | 2.18  | 1.65  | 1.57  | 1.20  | 1.66  |
| Percent of Stock         |      |       |       |       |       |       |       |       |
| Cropped:                 |      |       |       |       |       |       |       |       |
| Any-Lure Zone            | 8    | 7     | 11    | 7     | 9     | 9     | 11    | 9     |
| Flies-Only Zone          | 12   | 5     | 11    | 6     | 10    | 11    | 13    | 10    |
| Released:                |      |       |       |       |       |       |       |       |
| Any-Lure Zone            | 29   | 44    | 37    | 49    | 46    | 34    | 45    | 41    |
| Flies-Only Zone          | 42   | 41    | 34    | 40    | 45    | 30    | 28    | 37    |
| Ratio - Trout Released   |      |       |       |       |       |       |       |       |
| to Trout Creeled:        |      |       |       |       |       |       |       |       |
| Any-Lure Zone            | 3.5  | 6.7   | 5.2   | 6.8   | 5.0   | 3.8   | 4.0   | 4.9   |
| Flies-Only Zone          | 3.7  | 8.8   | 3.0   | 6.2   | 4.6   | 2.8   | 2.2   | 4.0   |
| Percent "Successful"     |      |       |       |       |       |       |       |       |
| Trips:                   |      |       |       |       |       |       |       |       |
| Any-Lure Zone            | 26   | 27    | 31    | 27    | 32    | 39    | 33    | 31    |
| Flies-Only Zone          | 39   | 27    | 41    | 35    | 35    | 47    | 45    | 38    |
| Avg. Length of Trout     |      |       |       |       |       |       |       |       |
| Creeled - Inches:        |      |       |       |       |       |       |       |       |
| Any-Lure Zone            | 8.6  | 8.7   | 8.6   | 8.5   | 8.5   | 8.6   | 8.5   | 8.6   |
| Flies-Only Zone          | 8.8  | 8.9   | 8.7   | 8.9   | 8.7   | 8.8   | 8.8   | 8.8   |

\* Any-lure zone = 1.86 miles and flies-only zone = 1.44 miles

TABLE 11

Number and Ratios of Legal Sized and Sublegal (6-8 Inch) Brook Trout  
in the Any-Lure Zone and Flies-Only Zone of Lawrence Creek During 1961-67

| Item                           | 1961   | 1962   | 1963  | 1964  | 1965  | 1966  | 1967  | Avg.  |
|--------------------------------|--------|--------|-------|-------|-------|-------|-------|-------|
| <b>Preseason No. of Legal</b>  |        |        |       |       |       |       |       |       |
| Trout:                         |        |        |       |       |       |       |       |       |
| Any-Lure Zone                  | 81     | 284    | 300   | 411   | 337   | 558   | 561   |       |
| Flies-Only Zone                | 298    | 468    | 719   | 550   | 412   | 524   | 903   |       |
| <b>Midseason No. of Legal</b>  |        |        |       |       |       |       |       |       |
| Trout:                         |        |        |       |       |       |       |       |       |
| Any-Lure Zone                  | 118    | 125    | 313   | 289   | 238   | 433   | 380   |       |
| Flies-Only Zone                | 260    | 209    | 479   | 344   | 215   | 493   | 516   |       |
| <b>Postseason No. of Legal</b> |        |        |       |       |       |       |       |       |
| Trout:                         |        |        |       |       |       |       |       |       |
| Any-Lure Zone                  | 496    | 188    | 349   | 270   | 318   | 424   | 332   |       |
| Flies-Only Zone                | 386    | 338    | 815   | 341   | 358   | 589   | 563   |       |
| <b>Preseason No. of Sub-</b>   |        |        |       |       |       |       |       |       |
| <b>legal Trout:</b>            |        |        |       |       |       |       |       |       |
| Any-Lure Zone                  | 458    | 1,024  | 943   | 1,395 | 1,191 | 1,282 | 1,754 | 1,150 |
| Flies-Only Zone                | 563    | 1,410  | 1,096 | 1,283 | 694   | 1,038 | 946   | 1,004 |
| <b>Midseason No. of Sub-</b>   |        |        |       |       |       |       |       |       |
| <b>legal Trout:</b>            |        |        |       |       |       |       |       |       |
| Any-Lure Zone                  | 1,239  | 1,662  | 1,309 | 2,157 | 1,601 | 2,138 | 1,522 | 1,661 |
| Flies-Only Zone                | 1,024  | 2,079  | 1,372 | 1,711 | 1,026 | 1,502 | 1,088 | 1,400 |
| <b>Postseason No. of Sub-</b>  |        |        |       |       |       |       |       |       |
| <b>legal Trout:</b>            |        |        |       |       |       |       |       |       |
| Any-Lure Zone                  | 931    | 1,578  | 1,001 | 1,927 | 1,346 | 2,069 | 1,338 | 1,456 |
| Flies-Only Zone                | 603    | 1,571  | 935   | 1,364 | 1,047 | 1,171 | 1,125 | 1,117 |
| <b>Legal: Sublegal Pre-</b>    |        |        |       |       |       |       |       |       |
| <b>season Ratio:</b>           |        |        |       |       |       |       |       |       |
| Any-Lure Zone                  | 1:5.6  | 1:3.6  | 1:3.1 | 1:3.4 | 1:3.5 | 1:2.3 | 1:3.1 |       |
| Flies-Only Zone                | 1:1.9  | 1:3.0  | 1:1.5 | 1:2.3 | 1:1.7 | 1:2.0 | 1:1.1 |       |
| <b>Legal: Sublegal Mid-</b>    |        |        |       |       |       |       |       |       |
| <b>season Ratio:</b>           |        |        |       |       |       |       |       |       |
| Any-Lure Zone                  | 1:10.5 | 1:13.3 | 1:4.2 | 1:7.5 | 1:6.7 | 1:4.9 | 1:4.0 |       |
| Flies-Only Zone                | 1:3.9  | 1:9.9  | 1:2.9 | 1:5.0 | 1:4.8 | 1:3.0 | 1:2.1 |       |
| <b>Legal: Sublegal Post-</b>   |        |        |       |       |       |       |       |       |
| <b>season Ratio:</b>           |        |        |       |       |       |       |       |       |
| Any-Lure Zone                  | 1:1.9  | 1:8.4  | 1:2.9 | 1:7.1 | 1:4.3 | 1:4.9 | 1:4.0 |       |
| Flies-Only Zone                | 1:1.6  | 1:4.6  | 1:1.5 | 1:4.0 | 1:2.9 | 1:2.0 | 1:2.0 |       |

TABLE 12

Overwinter Survival\* of Brook Trout in the Any-Lure Zone  
and Flies-Only Zone of Lawrence Creek During 1960-67  
as Compared to Survival During the Winters of 1955-60

| Winter       | Percent Survival |                 |               |
|--------------|------------------|-----------------|---------------|
|              | Any-Lure Zone    | Flies-Only Zone | Entire Stream |
| 1960-61      | 33               | 43              | 37            |
| 1961-62      | 42               | 86              | 58            |
| 1962-63      | 54               | 63              | 59            |
| 1963-64      | 69               | 65              | 69            |
| 1964-65      | 47               | 53              | 49            |
| 1965-66      | 74               | 76              | 75            |
| 1966-67      | 61               | 93              | 71            |
| Avg. 1960-67 | 54               | 68              | 60            |
| 1955-60      | 49               | 56              | 51            |

\*. Since there are no barriers between zones, overwinter changes in the trout populations in the two zones reflect movement as well as survival. Only the values for the entire stream reflect true survival.

TABLE 13

Distribution of Seasonal Angling Trips and Catch  
for the 1955-1967 Fishing Seasons According to the Methods  
of Fishing Employed at Lawrence Creek

| Angling Season | Angling Trips |                          |                           | Brook Trout Creeled |                          |                           |
|----------------|---------------|--------------------------|---------------------------|---------------------|--------------------------|---------------------------|
|                | No. of Trips  | Percent by Fly Fishermen | Percent by Bait Fishermen | No. of Trout        | Percent by Fly Fishermen | Percent by Bait Fishermen |
| 1955           | 1,712         | 25                       | 61                        | 3,040               | 22                       | 68                        |
| 1956           | 1,313         | 16                       | 71                        | 1,360               | 14                       | 77                        |
| 1957           | 1,249         | 19                       | 65                        | 2,778               | 16                       | 74                        |
| 1958           | 758           | 24                       | 53                        | 223                 | 19                       | 62                        |
| 1959           | 728           | 21                       | 53                        | 243                 | 14                       | 60                        |
| 1960           | 434           | 28                       | 52                        | 85                  | 14                       | 45                        |
| 1961           | 592           | 55                       | 38                        | 442                 | 66                       | 28                        |
| 1962           | 896           | 49                       | 40                        | 540                 | 47                       | 49                        |
| 1963           | 874           | 49                       | 39                        | 752                 | 57                       | 36                        |
| 1964           | 881           | 42                       | 46                        | 625                 | 46                       | 43                        |
| 1965           | 858           | 41                       | 40                        | 671                 | 40                       | 45                        |
| 1966           | 826           | 39                       | 46                        | 862                 | 43                       | 42                        |
| 1967           | 1,178         | 40                       | 50                        | 1,111               | 46                       | 43                        |
| Avg. 1955-60   | 1,032         | 22                       | 59                        | 1,288               | 17                       | 64                        |
| 1961-67        | 872           | 45                       | 43                        | 715                 | 49                       | 41                        |

LITERATURE CITED

- Allen, K. R. and B. T. Cunningham  
1957. New Zealand angling. 1947-52. Results of the dairy scheme. New Zealand Marine Dept. Fish. Bull. No. 12: 153 pp.
- Hobbs, D. F.  
1948. Trout fisheries in New Zealand; their development and management. Fish. Bull. Wellington. No. 3., No. 9.
- Hunt, R. L., O. M. Brynildson and J. T. McFadden  
1962. Effects of angling regulations on a wild brook trout fishery. Tech. Bull. No. 26. Wis. Conserv. Dept., Madison, Wis.: 58 pp.
- Hunt, R. L. and O. M. Brynildson  
1964. A five-year study of a headwaters trout refuge. Trans. Amer. Fish. Soc. 93(2): 194-197.
- Hunt, R. L.  
1964. Evaluation of fly-fishing-only at Lawrence Creek (a three-year progress report). Misc. Res. Rept. No. 10 (Fisheries). Wis. Conserv. Dept., Madison, Wis.: 15 pp.  
1966. Production and angler harvest of wild brook trout in Lawrence Creek, Wisconsin. Wis. Conserv. Dept., Madison, Wis.: 52 pp.  
1969. Effects of habitat alteration on production, standing crops and yield of brook trout in Lawrence Creek, Wisconsin. Symposium on salmon and trout. The Univ. of British Columbia, 1969: 388 pp.
- Mason, J. W. and R. L. Hunt  
1967. Mortality rates of deeply hooked rainbow trout. The Prog. Fish Cult., 29(1): 87-91.
- McFadden, J. T.  
1956. Characteristics of trout angling at Lawrence Creek, Wisconsin. Trans. Wis. Acad. Sci., Arts, and Lett. 45:21-29.  
1961. A population study of the brook trout, Salvelinus fontinalis. Wildl. Monog. No. 7: 73 pp.
- Shetter, D. S. and G. R. Alexander  
1962. Effects of a flies-only restriction on angling and on fall trout populations in Hunt Creek, Montmorency County, Michigan. Trans. Amer. Fish. Soc. 91(3):295-302.
- Stroud, R. H.  
1964. Most fishing is for fun - what else? Sport Fishing Institute Bull. No. 150:1.

Wallis, O. L.

1963. Application of fishing-for-fun programs. Mimeog. Rept. of paper presented at 93rd An. meeting of Amer. Fish. Soc.: 25 pp.

White, D. A.

1967. Trophic dynamics of a wild brook trout stream. Ph.D. thesis. The Univ. of Wis., Madison, Wis.: 183 pp.











