

Winstrol advertisement.

[s.l.]: [s.n.], 1962

<https://digital.library.wisc.edu/1711.dl/LWYX2FU2ZKJRI8Q>

<http://rightsstatements.org/vocab/InC/1.0/>

The libraries provide public access to a wide range of material, including online exhibits, digitized collections, archival finding aids, our catalog, online articles, and a growing range of materials in many media.

When possible, we provide rights information in catalog records, finding aids, and other metadata that accompanies collections or items. However, it is always the user's obligation to evaluate copyright and rights issues in light of their own use.

BUILDS *body tissue...*

BUILDS *sense of well-being
in the weak and debilitated*

WINSTROL®

BRAND OF STANZOLOL

new physiotonic

NOW—the **highest** anabolic plus the **lowest** androgenic activity* with well-tolerated WINSTROL therapy

...for elderly patients with anorexia, asthenia and general debility—

MARKED IMPROVEMENT IN APPETITE, STRENGTH AND SENSE OF WELL-BEING

Fourteen patients, age 66 to 77, treated with Winstrol, usually in a dosage of 6 mg. daily, for various periods in order to correct underweight, weakness and chronic fatigue. Marked improvement occurred in appetite, sense of well-being and strength; almost all patients gained weight.

...for patients with osteoporosis and arthritis—

RELIEF OF PAIN, IMPROVEMENT IN MOBILITY

Twenty-one patients with arthritis treated with Winstrol for pain and limited mobility due to osteoporosis. With few exceptions, dosage was 6 mg. daily; duration of treatment varied from a few weeks to 6 months. In 8 patients relief of symptoms was excellent and in 6 moderate. Of the 7 persons in whom no relief was obtained, 5 had received treatment for less than one month and some had been given doses below 6 mg.

...for patients with malignant disease—

NOTABLE WEIGHT GAINS, INCREASED APPETITE AND SENSE OF WELL-BEING

Twenty-six patients, mostly women, weak, emaciated and seriously ill, were administered Winstrol in dosage of 6 mg. daily for periods extending up to 14 months (average 6.7 months). Notable weight gains occurred. Patients showed increased appetite, alertness and confidence, better appearance, increased mobility and tolerance to pain.

...for patients with chronic, non-malignant disorders—

IMPROVEMENT IN WEIGHT AND GENERAL ACTIVITY, INCREASED SENSE OF WELL-BEING

Eight patients with advanced tuberculosis, bronchopulmonary disease, nephritis and ulcerative colitis treated with 6 mg. of Winstrol daily for from 3 to 4 months. Gains in weight varied from 6 to 27 pounds with increased sense of well-being and improvement in general activity.

...for undernourished, underweight children and adolescents—

NOTABLE IMPROVEMENT IN APPETITE AND OUTLOOK, MARKED INCREASE IN WEIGHT AND HEIGHT

One hundred and twenty children, age 1 to 11 years, underweight and in poor health, were given Winstrol for several months. Majority received daily dosage of from 2 to 4 mg. In nearly all, appetite was improved. Over 70 per cent showed significant gains in weight of from 5 to 17 pounds.

DOSAGE: Usual adult dose, one 2 mg. tablet t.i.d. just before or with meals; children from 6 to 12 years, up to 1 tablet t.i.d.; children under 6 years, ½ tablet b.i.d. Available in bottles of 100 tablets.

Complete bibliography and literature available on request. Before prescribing, consult literature for additional dosage information, possible side effects and contraindications.

*animal data

**WITH NEW
WINSTROL**

patients look better...feel stronger—because they are stronger

Winthrop LABORATORIES
New York 18, N. Y.