



LIBRARIES
UNIVERSITY OF WISCONSIN - MADISON

Hycomine Pediatric Syrup advertisement.

[s.l.]: [s.n.], 1973

<https://digital.library.wisc.edu/1711.dl/Z7E2V7YLCIVOS8B>

<http://rightsstatements.org/vocab/InC/1.0/>

The libraries provide public access to a wide range of material, including online exhibits, digitized collections, archival finding aids, our catalog, online articles, and a growing range of materials in many media.

When possible, we provide rights information in catalog records, finding aids, and other metadata that accompanies collections or items. However, it is always the user's obligation to evaluate copyright and rights issues in light of their own use.

HYCOMINE[®] PEDIATRIC SYRUP[Ⓜ]

Each teaspoonful (5 ml.) contains: hydrocodone bitartrate (Warning: may be habit forming) 2.5 mg., homatropine methylbromide 0.75 mg., pyrilamine maleate 6.25 mg., phenylephrine hydrochloride 5 mg., ammonium chloride 30 mg.

Contraindications: Hypersensitivity to any component of the drug, and in patients with glaucoma, pyloric obstruction, or prostatic hypertrophy.

Precautions: Use with caution in diabetes, hyperthyroidism, hypertension, and cardiovascular disease. Since blurred vision, drowsiness and dizziness may occur, patients should be cautioned about activities requiring alertness.

Before prescribing antitussive medication to suppress or modify cough, it is important to ascertain that the underlying cause of the cough is identified, that modification of the cough does not increase the risk of clinical or physiologic complications, and that appropriate therapy for the primary disease is provided.

Adverse Reactions: HYCOMINE Syrup is generally well tolerated. Occasional drowsiness, blurred vision, cardiac palpitation, dizziness, nervousness, dryness of the mouth, or gastrointestinal upset may occur.

Usual Dosage: Children 6 months to 1 year, 10 drops; 1 to 3 years, $\frac{1}{4}$ teaspoonful (20 drops); 3 to 6 years, $\frac{1}{2}$ teaspoonful; 6 to 12 years, 1 teaspoonful; over 12 years, 2 teaspoonfuls. Administer after meals and at bedtime, not less than 4 hours apart.

Oral prescription where permitted by state law.

Endo Laboratories, Inc.

Subsidiary of E.I. du Pont de Nemours & Co. (Inc.)
Garden City, New York 11530



Captain Kidd has a bad cold



He's miserable! His nose is stuffed, he's sneezing, he's coughing, he has difficulty sleeping and worst of all he has no friends to play with. It's enough to drive his mother "up the wall."

When she comes to you for help, remember that HYCOMINE® Pediatric Syrup can treat the Captain's cold symptoms. It stops the cough, decongests mucous passages, and dries secretions. Especially formulated for children, HYCOMINE® Pediatric Syrup can make Captain Kidd comfortable while his cold runs its course. His mother will like that as well. ORAL PRESCRIPTIONS WHERE PERMITTED BY STATE LAW. Rx's MAY BE REFILLED 5 TIMES IN 6 MONTHS.

Gentlemen: Please send me a clinical supply (1 x 4 oz.) of HYCOMINE® Pediatric Syrup*

_____, M.D. BNDD # _____
(Please sign and return)

Street _____

City _____ State _____ Zip _____

Name, address and BNDD Number must be legible and complete for us to honor your request.

*Where permitted by state laws and regulations.

Mail to: Professional Request Dept.
Endo Laboratories, Inc.,
1000 Stewart Ave., Garden City, N.Y. 11530

HYCOMINE® PEDIATRIC SYRUP III

Each teaspoonful (5 ml) contains: hydrocodone bitartrate, 2.5 mg (Warning: May be habit forming); homatropine methylbromide, 0.75 mg; pyrilamine maleate, 6.25; phenylephrine hydrochloride, 5 mg; ammonium chloride, 30 mg.

Cold comfort for the Captain

See facing page for Brief Summary of Prescribing Information.