



Obetrol advertisement.

[s.l.]: [s.n.], 1964

<https://digital.library.wisc.edu/1711.dl/WO34PQFEU4N2X86>

<http://rightsstatements.org/vocab/InC/1.0/>

The libraries provide public access to a wide range of material, including online exhibits, digitized collections, archival finding aids, our catalog, online articles, and a growing range of materials in many media.

When possible, we provide rights information in catalog records, finding aids, and other metadata that accompanies collections or items. However, it is always the user's obligation to evaluate copyright and rights issues in light of their own use.

**...but doctor,
I eat like a bird!**



IN WEIGHT CONTROL

OBETROL®

Each OBETROL-10 tablet contains: Methamphetamine Saccharate; 2.5 mg. Methamphetamine Hydrochloride; 2.5 mg. Amphetamine Sulfate; 2.5 mg. Dextro-amphetamine Sulfate; 2.5 mg. (OBETROL-20 tablets contain twice this potency) Pat. #2748052.

This combination of amphetamines may be useful as an adjunct in the management of certain forms of obesity where an appetite depressant is indicated.

CONTRAINDICATIONS: Hypertension, advanced arteriosclerosis, coronary artery disease, cardiac arrhythmias, peripheral vascular disease, states of undue restlessness, anxiety, excitement, agitated depression, hyperthyroidism, idiosyncrasy to amphetamine, concomitant administration of a monoamine oxidase inhibitor.

PRECAUTIONS: Use with caution in individuals with anorexia, insomnia, vasomotor instability, asthenia, psychopathic personality, a history of homicidal or suicidal tendencies, and individuals who are known to be hyper-reactive to sympathomimetic agents, or emotionally unstable individuals who are known to be susceptible to drug abuse. Certain monoamine oxidase inhibitors may potentiate the action of Obetrol. **SIDE EFFECTS:** The most common side effects attended with the use of amphetamines include nervousness, excitability, euphoria, insomnia, dryness of mouth, nausea, vertigo, constipation, and headache. **DOSAGE AND ADMINISTRATION:** Initial adult dose is one-half to one 'Obetrol-10' tablet daily, preferably one-half to one hour before meals. This may be gradually increased to one 'Obetrol-10' or 'Obetrol-20' tablet one to three times daily as indicated.

SUPPLIED: Tablets scored, in bottles of 100, 500, and 1000.

REQUEST SAMPLES AND LITERATURE

OBETROL PHARMACEUTICALS • BROOKLYN, N.Y. 11207

Dr. _____

Address _____

City _____ State _____

OBETROL PHARMACEUTICALS

Div. of Rexar Pharmacal Corp.
Brooklyn, N.Y. 11207