



Donnagel with Neomycin advertisement.

[s.l.]: [s.n.], 1956

<https://digital.library.wisc.edu/1711.dl/UDBYY6YMY2ROG8H>

<http://rightsstatements.org/vocab/InC/1.0/>

The libraries provide public access to a wide range of material, including online exhibits, digitized collections, archival finding aids, our catalog, online articles, and a growing range of materials in many media.

When possible, we provide rights information in catalog records, finding aids, and other metadata that accompanies collections or items. However, it is always the user's obligation to evaluate copyright and rights issues in light of their own use.

- the
preparation
you've
asked
for

ANNOUNCING: a NEW antidiarrheal for

more certain control of virtually
all diarrheas

DONNAGEL® WITH NEOMYCIN

ANTIBIOTIC - ADSORBENT - DEMULCENT - ANTISPASMODIC



Addition of neomycin to the effective DONNAGEL formula assures even more certain control of most of the common forms of diarrhea. Neomycin is an ideal antibiotic for enteric use: it is effectively bacteriostatic against neomycin-susceptible pathogens; and it is relatively non-absorbable.

The secret of DONNAGEL WITH NEOMYCIN's clinical dependability lies in the comprehensive approach of its rational formula:

COMPONENT in each 30 cc. (1 fl. oz.)	ACTION	BENEFIT
Neomycin base, 210.0 mg. (as neomycin sulfate, 300 mg.)	antibiotic	Affords effective intestinal bacteriostasis.
Kaolin (6.0 Gm.)	adsorbent, demulcent	Binds toxic and irritating substances. Provides protective coating for irritated intestinal mucosa.
Pectin (142.8 mg.)	protective, demulcent	Supplements action of kaolin as an intestinal detoxifying and demulcent agent.
Dihydroxyaluminum aminoacetate (0.25 Gm.)	antacid, demulcent	Enhances demulcent and detoxifying action of the kaolin-pectin suspension.
Natural belladonna alkaloids: hyoscyamine sulfate (0.1037 mg.) atropine sulfate (0.0194 mg.) hyoscine hydrobromide (0.0065 mg.)	anti- spasmodic	Relieves intestinal hypermotility and hypertonicity.
Phenobarbital (1/4 gr.)	sedative	Diminishes nervousness, stress and apprehension.

Robins

Informational
literature
available
upon request.

INDICATIONS: DONNAGEL WITH NEOMYCIN is specifically indicated in diarrheas or dysentery caused by neomycin-susceptible organisms; in diarrheas not yet proven to be of bacterial origin, prior to definitive diagnosis. Also useful in enteritis, even though diarrhea may not be present. **SUPPLIED:** Bottles of 6 fl. oz. At all prescription pharmacies.

DOSAGE: Adults: 1 to 2 tablespoonsfuls (15 to 30 cc.) every 4 hours. Children over 1 year: 1 to 2 teaspoonsfuls every 4 hours. Children under 1 year: 1/2 to 1 teaspoonful every 4 hours.

ALSO AVAILABLE: DONNAGEL, the original formula, for use when an antibiotic is not indicated.