



LIBRARIES
UNIVERSITY OF WISCONSIN - MADISON

Trilisate advertisement.

[s.l.]: [s.n.], 1978

<https://digital.library.wisc.edu/1711.dl/MYWV7ATNULG4M8H>

<http://rightsstatements.org/vocab/InC/1.0/>

The libraries provide public access to a wide range of material, including online exhibits, digitized collections, archival finding aids, our catalog, online articles, and a growing range of materials in many media.


When possible, we provide rights information in catalog records, finding aids, and other metadata that accompanies collections or items. However, it is always the user's obligation to evaluate copyright and rights issues in light of their own use.

ONCE-A-DAY DOSAGE

WITH PREDICTABLE POWER FOR
24-HOUR ARTHRITIS CONTROL

ONCE-A-DAY

Trilisate[®] TABLETS/LIQUID
(choline magnesium trisalicylate)




PROVEN THERAPEUTIC RESPONSE WITH ONCE-A-DAY DOSAGE

Once-a-day TRILISATE is proven effective for relief of arthritis symptoms as measured against pretreatment levels of these major parameters: duration of morning stiffness, number of painful joints, number of swollen joints, articular index, and swelling index.^{1,2}

PREDICTABLE THERAPEUTIC SERUM LEVELS

Once-a-day TRILISATE achieves rapid, sustained therapeutic serum levels³—for a full 24 hours of arthritis control. In only three to four days, once-a-day TRILISATE provides antiarthritic effectiveness all day and all night.



WELL TOLERATED

TRILISATE is proven well tolerated in most patients, even at the higher dosages recommended for 24-hour arthritis control.¹⁻³

CONVENIENT

Your patients will especially appreciate the convenience of once-a-day dosage—proven to aid patient compliance.⁴ A single dose (preferably at bedtime) provides a full 24 hours of arthritis control.

ONCE-A-DAY
Trilisate® TABLETS/LIQUID
(choline magnesium trisalicylate)

Once-a-day dosage for all-day, all-night arthritis control

References:

1. Zeiders, R. S.: Once daily salicylate therapy in rheumatoid arthritis, presented at the 15th International Congress of Rheumatology, Paris, June 21-27, 1981.
 2. Zeiders, R. S., Seebach, L. M., Boyer, J. T., Miller, S.: Multicenter study, data on file, Medical Department, The Purdue Frederick Company, Norwalk, CT, 1981.
 3. Rothwell, K. G.: Cross-over comparative study of TRILISATE tablets once daily and twice daily in normal subjects, data on file, Medical Department, The Purdue Frederick Company, Norwalk, CT, 1981.
 4. Gately, M. S.: To be taken as directed, *J. Roy. Coll. Gen. Practit.* 16:39-44, 1968.
- U. S. Patent Numbers 3759980 and 4067974.

Purdue Frederick

© Copyright 1981, The Purdue Frederick Company/Norwalk, CT 06856

B576

PFQ-874/81

Please see following page for prescribing information.

ONCE-A-DAY
Trilisate[®] TABLETS/LIQUID
(choline magnesium trisalicylate)

Brief summary of prescribing information

Indications:

Relief of signs and symptoms of rheumatoid arthritis, osteoarthritis, and other arthritides, in both the acute flare and the long-term management of these diseases. Also for alleviating the pain and loss of motion of acute painful shoulder syndrome.

Contraindications: Hypersensitivity to salicylates.

Precautions: Like other salicylates, TRILISATE Tablets/Liquid should be used with caution in patients with chronic renal insufficiency, active erosive gastritis, or peptic ulcer. Reports indicate that when salicylates are given with steroids, the butazones, or alcohol, the risk of gastrointestinal ulceration is increased. Caution should also be exercised in patients requiring coumarin or indandione anticoagulants, or heparin. As with any drug, the usual care should be exercised during pregnancy; use immediately prior to onset of labor is not recommended. TRILISATE Tablets/Liquid have not been studied in rheumatoid arthritis patients in the Functional Class IV, nor in children under 12 years of age.

Adverse Reactions: TRILISATE Tablets/Liquid are generally well tolerated at recommended dosage ranges, particularly by the gastrointestinal system. As with all salicylates, salicylism and/or salicylate intoxication may occur with large doses or with extended therapy. Tinnitus may be regarded as a therapeutic guide; should it develop, reduction of dosage is recommended.

Dosage Summary: For moderate to severe disease, initiate therapy with 2000 mg to 2500 mg once a day (preferably at bedtime) or in divided doses (q. 12 h.). For mild disease, begin therapy with 1500 mg once a day or in divided doses (q. 12 h.). Allow 3-4 days to achieve steady state. Increase or decrease dosage, if necessary.

Supplied: TRILISATE 500 Tablets (pale pink, scored) in bottles of 100 tablets. TRILISATE 750 Tablets (white, scored) in bottles of 100 tablets.

TRILISATE Liquid in bottles of 8 fl. oz. (237 ml). Only on prescription.

Purdue Frederick

The Purdue Frederick Company/Norwalk, CT 06856

B576 PFG-874/81