

Tuss-Ornade advertisement.

[s.l.]: [s.n.], 1979


<https://digital.library.wisc.edu/1711.dl/JFRNE6TNICJOR85>

<http://rightsstatements.org/vocab/InC/1.0/>

The libraries provide public access to a wide range of material, including online exhibits, digitized collections, archival finding aids, our catalog, online articles, and a growing range of materials in many media.

When possible, we provide rights information in catalog records, finding aids, and other metadata that accompanies collections or items. However, it is always the user's obligation to evaluate copyright and rights issues in light of their own use.

Expressly
for kids with
colds that
cough*



orange-pineapple flavored

Tuss-Ornade[®] cough/cold liquid

Each 5 ml. teaspoonful contains 5 mg. caramiphen edisylate; 2 mg. chlorpheniramine maleate; 15 mg. phenylpropanolamine hydrochloride; isopropamide iodide equivalent to 0.75 mg. of isopropamide; and alcohol, 7.5%.

Before prescribing, see complete prescribing information in SK&F literature or PDR. The following is a brief summary.

* Indications

Based on a review of this drug by the National Academy of Sciences—National Research Council and/or other information, FDA has classified the indications as follows: Lacking in substantial evidence of effectiveness as a fixed combination: For relief from coughing, upper respiratory congestion and hypersecretion associated with the common cold, sinusitis, vasomotor rhinitis and allergic rhinitis. Final classification of the less-than-effective indications requires further investigation.

Contraindications: Hypersensitivity to any component; concurrent MAO inhibitor therapy; severe hypertension; bronchial asthma; coronary artery disease; stenosing peptic ulcer; pyloro-duodenal or bladder neck obstruction. Do not use 'Tuss-Ornade' Liquid in children less than six months of age or under 15 lbs. in weight. Do not use 'Tuss-Ornade' Spansule capsules in children under 12 years of age.

Warnings: Warn vehicle or machine operators of possible drowsiness. Warn patients of possible additive effects of alcohol and other C.N.S. depressants.

Usage in Pregnancy: Use in pregnancy, nursing mothers and women who might bear children only when potential benefits

have been weighed against possible hazards. An inhibitory effect on lactation may occur.

Effect on PBI Determination and I¹³¹ Uptake: The iodine in isopropamide iodide may alter PBI test results and will suppress I¹³¹ uptake; use thyroid tests unaffected by exogenous iodides.

Precautions: Use with caution in persons with cardiovascular disease, glaucoma, prostatic hypertrophy, hyperthyroidism.

Adverse Reactions: Drowsiness; excessive dryness of nose, throat or mouth; nervousness, insomnia; nausea, vomiting, diarrhea; rash; dizziness; weakness; tightness of chest; angina pain; abdominal pain; irritability; palpitation; headache; incoordination; tremor; difficulty in urination; thrombocytopenia, leukopenia; convulsions; hypertension, hypotension; anorexia; constipation; visual disturbances; iodine toxicity (acne, parotitis); dysuria; epigastric distress.

Supplied: 'Tuss-Ornade' Liquid: An orange-pineapple flavored liquid in 16 fl. oz. bottles. For patients 12 years or older, 'Tuss-Ornade' Spansule capsules (each capsule contains 20 mg. caramiphen edisylate; 8 mg. chlorpheniramine maleate; 50 mg. phenylpropanolamine HCl; and isopropamide iodide equivalent to 2.5 mg. of isopropamide), in bottles of 50 and 500 capsules.

SK&F
a SmithKline company

Smith Kline & French Laboratories
Div. of SmithKline Corp., Phila., Pa.

©Smith Kline & French Laboratories, 1979