



Bamadex Sequels advertisement.


[s.l.]: [s.n.], 1964

<https://digital.library.wisc.edu/1711.dl/HDM6L6Q3XTIIN8C>

<http://rightsstatements.org/vocab/InC/1.0/>

The libraries provide public access to a wide range of material, including online exhibits, digitized collections, archival finding aids, our catalog, online articles, and a growing range of materials in many media.

When possible, we provide rights information in catalog records, finding aids, and other metadata that accompanies collections or items. However, it is always the user's obligation to evaluate copyright and rights issues in light of their own use.



The Lederle Skinfold Caliper, which measures the thickness of subcutaneous fat, has great psychological value for demonstrating to the patient that he or she is not just stout or heavy but "fat." Such a demonstration, independent of weight-and-height charts, can be an important factor in motivating the patient to follow the physician's weight-reducing instructions more faithfully. *When a patient fails the skinfold test, he usually faces a diet test later. That's when BAMADEX SEQUELS can be helpful.* By combining in sustained release form the appetite-suppressant action of *d*-amphetamine and the tranquilizing action of meprobamate, BAMADEX SEQUELS lend vital support to an overweight patient's efforts to establish a new pattern of eating less—the ultimate aim of weight-reduction therapy.



A single BAMADEX SEQUELS Capsule one-half hour before breakfast curtails appetite all day, while at the same time easing the tension and nervousness that often accompany diet curtailment.

Indications: Dietary adjunct in control of obesity.

Side Effects: Occasional allergic skin reactions may occur with meprobamate, accompanied by fever, nonthrombocytopenic purpura, angioneurotic edema, hypotension, and bronchial spasm. Drowsiness with or without ataxia and/or alteration in visual accommodation may occur. Effects of alcoholic beverages may

be increased by meprobamate. Use with caution in psychoneurotics, in patients with coronary or cardiovascular disease or severe hypertension. Prolonged use may result in dependence. Reactions can occur if drug is not gradually withdrawn.

Available on Request: The Lederle Skinfold Caliper illustrated here is available to physicians free of charge. Write your request on your prescription blank and mail to Lederle Laboratories, Room 408, Bldg. 2, Pearl River, New York.

WHEN YOUR PATIENT FAILS THE SKINFOLD TEST

BAMADEX® SEQUELS®

d-amphetamine sulfate (15 mg.) and meprobamate (300 mg.)

SUSTAINED RELEASE CAPSULES

LEDERLE LABORATORIES • A Division of AMERICAN CYANAMID COMPANY, Pearl River, New York



338-4