



LIBRARIES
UNIVERSITY OF WISCONSIN - MADISON

Actifed-C Expectorant advertisement.


[s.l.]: [s.n.], 1975

<https://digital.library.wisc.edu/1711.dl/VS2GZ4QDMN2QD8L>

<http://rightsstatements.org/vocab/InC/1.0/>


The libraries provide public access to a wide range of material, including online exhibits, digitized collections, archival finding aids, our catalog, online articles, and a growing range of materials in many media.

When possible, we provide rights information in catalog records, finding aids, and other metadata that accompanies collections or items. However, it is always the user's obligation to evaluate copyright and rights issues in light of their own use.



**It's All in the Family...
a family of cough-and-cold benefits:**

- 1.** to reduce excessive coughing without preventing productive expectoration
- 2.** to decongest nasal passages and eustachian tubes



One medication for the whole family.*

ACTIFED-C[®] EXPECTORANT

Each 5 cc teaspoonful contains: codeine phosphate 10 mg (Warning: may be habit-forming); Actidil[®] brand Triprolidine Hydrochloride 2 mg; Sudafed[®] brand Pseudoephedrine Hydrochloride 30 mg; glyceryl guaiacolate 100 mg. Preservatives are methylparaben 0.1% and sodium benzoate 0.1%.

Indications: For symptomatic relief of cough in conditions such as: the common cold, acute bronchitis, allergic asthma, bronchiolitis, croup, emphysema and tracheobronchitis.

Warning: Use in Pregnancy: Experience with this drug in pregnant women is inadequate to determine whether there exists a potential for harm to the developing fetus.

Precautions: Although pseudoephedrine hydrochloride causes virtually no pressor

effect in normotensive patients, it should be used with caution in patients with hypertension. In addition, even though triprolidine hydrochloride produces only a low incidence of drowsiness, appropriate precautions should be observed.

Side effects: The great majority of patients will not have any side effects. Only certain patients, sensitive to one or another of the ingredients, may note mild stimulation or mild sedation.

Supplied: Bottles of 1 pint. Complete literature available on request from Professional Services Dept. PML.

*Not recommended for use in children under 2 years of age.



Burroughs Wellcome Co.
Research Triangle Park
North Carolina 27709