

Obedrin-LA advertisement.

[s.l.]: [s.n.], 1960

<https://digital.library.wisc.edu/1711.dl/6JUQUAMSE5YOG8E>

<http://rightsstatements.org/vocab/InC/1.0/>

The libraries provide public access to a wide range of material, including online exhibits, digitized collections, archival finding aids, our catalog, online articles, and a growing range of materials in many media.

When possible, we provide rights information in catalog records, finding aids, and other metadata that accompanies collections or items. However, it is always the user's obligation to evaluate copyright and rights issues in light of their own use.

*"Why,
Mrs. Nelson
YOU'RE A
SIZE 12
AGAIN"*



She didn't get back into a 12 by merely counting calories. Her doctor prescribed Obedrin-LA and the Obedrin Menu Plan. The all-day trickle release action of Obedrin-LA effectively curbs appetite and reduces anxiety without excessive C.N.S. stimulation. The Obedrin Menu Plan permits flexible meal planning without restrictive calorie counting. Add your supervision to these benefits and the result is proved and acceptable for a weight control regimen.

*Write for
Free Copies of the
Obedrin Menu Plan*

FORMULA: Each OBEDRIN-LA Tablet contains:

Methamphetamine Hydrochloride*	12.5 mg.
Pentobarbital* (Warning: May be habit forming)	50 mg.
Ascorbic Acid	200 mg.
Thiamine Mononitrate	1 mg.
Riboflavin	2 mg.
Nicotinic Acid (Niacin)	10 mg.

DOSAGE: OBEDRIN-LA—1 tablet at 10 a.m. daily. Time of administration may be adjusted to fit individual patient's eating habits.

SUPPLIED: OBEDRIN-LA (long acting) Tablets as two-layer (one pink and the other beaded pink and white) in bottles of 50 on prescription only. OBEDRIN is also available as scored Tablets or Capsules for b.i.d. dosage.

PRECAUTIONS AND CONTRAINDICATIONS: OBEDRIN should be used with caution in patients hypersensitive to sympathomimetic compounds or barbiturates and in cases of coronary, cardiovascular or hypertensive disease.

*U.S. Pat. Nos. 2,736,682; 2,809,916; 2,809,917; 2,809,918; and pat. pend.

Long-Acting

Obedrin®-LA*

TABLETS

FORMULA: Methamphetamine Hydrochloride, 12.5 mg.; Pentobarbital (Warning: May be habit forming), 50 mg.; Ascorbic Acid, 200 mg.; Thiamine Mononitrate, 1 mg.; Riboflavin, 2 mg.; Nicotinic Acid (Niacin), 10 mg.

THE S. E. MASSENGILL COMPANY Bristol, Tennessee • New York • Dallas • Chicago • Kansas City • San Francisco