



LIBRARIES
UNIVERSITY OF WISCONSIN - MADISON

Bendectin advertisement.

[s.l.]: [s.n.], 1958

<https://digital.library.wisc.edu/1711.dl/OY33EYOBOHFPU84>

<http://rightsstatements.org/vocab/InC/1.0/>

The libraries provide public access to a wide range of material, including online exhibits, digitized collections, archival finding aids, our catalog, online articles, and a growing range of materials in many media.

When possible, we provide rights information in catalog records, finding aids, and other metadata that accompanies collections or items. However, it is always the user's obligation to evaluate copyright and rights issues in light of their own use.

stop
 "morning
 sickness"
 the
 night
 before
 with
 timed-
 release
Bendectin
 2 tabs. h.s.



PREVENTS "MORNING SICKNESS" IN 9 OUT OF 10 PREGNANCIES

In 941 cases^{1,2} effective in all but 17. Two timed-release tablets at bedtime start to work in the early morning and reach maximum potency at normal waking hour. BENDECTIN then provides exceptional relief of nausea and vomiting by three distinct and complementary actions. 1. **Antispasmodic**—Bentyl 10 mg.—relaxes G-I smooth-muscle spasm; 2. **Antinauseant**—Decapryn 10 mg.—centrally effective . . . combats histamine-like metabolites often present in blood stream during pregnancy; 3. **Nutritional supplement**—pyridoxine 10 mg.—just the amount necessary to help control "morning sickness."

1. Nulsen, R. O.: Ohio State M. J. 53:665 1957. 2. Personal communications, 1956-57.

Formula: Each tablet contains:
 Bentyl (dicyclomine) Hydrochloride 10 mg.
 Decapryn (doxylamine) Succinate . . . 10 mg.
 Pyridoxine Hydrochloride 10 mg.



THE WM. S. MERRELL COMPANY
 New York • CINCINNATI • St. Thomas, Ontario
 Another Exclusive Product of Original Merrell Research

TRADEMARKS: "BENTYL," "DECAPRYN," BENDECTIN®